

THE $\Pi_1^1 \downarrow$ LÖWENHEIM-SKOLEM-TARSKI PROPERTY OF STATIONARY LOGIC

SEAN COX

ABSTRACT. Fuchino-Maschio-Sakai [7] proved that the Löwenheim-Skolem-Tarski (LST) property of Stationary Logic is equivalent to the Diagonal Reflection Principle on internally club sets (DRP_{IC}) introduced in [4]. We prove that the restriction of the LST property to (downward) reflection of Π_1^1 formulas, which we call the $\Pi_1^1 \downarrow$ -LST property, is equivalent to the *internal* version of DRP from [2]. Combined with results from [2], this shows that the $\Pi_1^1 \downarrow$ -LST Property for Stationary Logic is strictly weaker than the full LST Property for Stationary Logic, though if CH holds they are equivalent.

CONTENTS

1.	Introduction	1
2.	Preliminaries	3
3.	Proof of Theorem 1.2	5
4.	Concluding remarks	8
	References	9

1. INTRODUCTION

Stationary Logic is a relatively well-behaved fragment of Second Order Logic introduced by Shelah [12], and first investigated in detail by Barwise et al [1]. Stationary Logic augments first order logic by introducing a new second order quantifier *stat*; we typically interpret “ $\text{stat}Z \phi(Z, \dots)$ ” to mean that there are stationarily many countable Z such that $\phi(Z, \dots)$ holds.¹ The quantifier *aa* stands for “almost all” or “for club many”; so

$$\text{aa}Z \phi(Z, \dots)$$

is an abbreviation for

$$\neg \text{stat}Z \neg \phi(Z, \dots).$$

2010 *Mathematics Subject Classification.* 03E05, 03E55, 03E35, 03E65 .

The author thanks Hiroshi Sakai (JSPS Kakenhi Grant Number 18K03397) for travel support to the RIMS Set Theory Workshop 2019, and also gratefully acknowledges support from Simons Foundation grant 318467.

¹Other interpretations, e.g. for uncountable Z , or for filters other than the club filter, are often considered too.

Section 2 provides more details.

By *structure* we will always mean a first order structure in a countable signature. The question of whether every structure has a “small” elementary substructure in Stationary Logic was raised already in [1]. One cannot hope to always get countable elementary substructures; e.g. if κ is regular and uncountable, then (κ, \in) satisfies “ \in is a linear order and

$$\text{aa}Z \exists x \text{ } x \text{ is an upper bound of } Z”,$$

but no countable linear order can satisfy that sentence. In a footnote in [1], it was observed that even the statement

(LST) “Every structure has an elementary (w.r.t. Stationary Logic) substructure of size $\leq \omega_1$ ”

carries large cardinal consistency strength.² The quoted statement above is now typically called the *Löwenheim-Skolem-Tarski* (LST) property of Stationary Logic.³

Fuchino et al. recently proved that LST is equivalent to a version of the Diagonal Reflection Principle introduced in Cox [4]:

Theorem 1.1 (Fuchino-Maschio-Sakai [7]). *LST is equivalent to the **Diagonal Reflection Principle on internally club sets** (DRP_{IC}).*

The purpose of the present note is to prove the following variant of Theorem 1.1 involving Π_1^1 formulas in Stationary Logic (defined in Section 2 below) and the principle DRP_{internal} from [2]:

Theorem 1.2. *The $\Pi_1^1\downarrow$ -LST property of Stationary Logic (see Definition 2.2) is equivalent to the principle DRP_{internal} .*

Cox [2] proved that DRP_{IC} is strictly stronger than DRP_{internal} . This was obtained by forcing over a model of a strong forcing axiom in a way that preserved DRP_{internal} while killing DRP_{IC} (in fact killing RP_{IC} ; the argument owed much to Krueger [10]). Furthermore, if CH holds, then DRP_{IC} is equivalent to DRP_{internal} . Combining those results with Theorem 1.2 immediately yields:

Corollary 1.3. *The LST property of Stationary Logic is **strictly stronger** than the $\Pi_1^1\downarrow$ -LST property of Stationary Logic.*


*However, if the Continuum Hypothesis holds, they are equivalent.*⁴

²See Definition 2.2 for precisely what is meant by “elementary substructure” in this context.

³The weaker assertion that every consistent theory (in Stationary Logic) has a model of size ω_1 , on the other hand, is a theorem of ZFC, as proven in [1].

⁴One doesn’t actually need the full continuum hypothesis for this equivalence to hold, but rather a variant of Shelah’s Approachability Property, namely that the class of internally stationary sets is the same (mod NS) as the class of internally club sets. See Cox [2] for more details.

FIGURE 1. An arrow indicates an implication, an arrow with an X indicates a non-implication



We note that while the technical strengthening MM^{++} of Martin's Maximum implies DRP_{IC} (see [4]), recent work of Cox-Sakai [6] shows that Martin's Maximum alone does not imply even the weakest version of DRP. Figure 1 summarizes the relevant implications and non-implications discussed in this introduction.

Section 2 covers the relevant preliminaries, and Section 3 proves Theorem 1.2. Section 4 ends with some concluding remarks.

2. PRELIMINARIES

Recall that $S \subseteq [A]^\omega$ is stationary if it meets every closed, unbounded subset of $[A]^\omega$ (in the sense of Jech [9]). By Kueker [11] this is equivalent to requiring that for every $f : [A]^{<\omega} \rightarrow A$ there is an element of S that is closed under f .

In what follows, we will use uppercase letters to denote second order variables/parameters, and lowercase letters to denote first order variables/parameters. We will also use some standard abbreviations; e.g. if our language includes the \in symbol, v is a first order variable, and Z is a second order variable, " $v = Z$ " is short for

$$\forall x \ x \in v \iff Z(x).$$

Given a structure $\mathfrak{A} = (A, \dots)$ (which we always assume to have a countable signature), the satisfaction relation in Stationary Logic is defined recursively by:

$$\begin{aligned} \mathfrak{A} \models \text{stat}Z \ \phi(Z, U_1, \dots, U_\ell, p_1, \dots, p_k) \\ \iff \\ \{Z \in [A]^\omega : \mathfrak{A} \models \phi(Z, U_1, \dots, U_\ell, p_1, \dots, p_k)\} \text{ is stationary in } [A]^\omega. \end{aligned}$$

We define a hierarchy of formulas in Stationary Logic that mimics the usual hierarchy in Second Order Logic. Since

$$aaZ \phi(Z, \dots)$$

roughly translates as

$$\exists C \text{ } C \text{ is club and } \forall Z \in C \phi(Z, \dots),$$

the *aa* quantifier will correspond to the existential second order quantifier when constructing the hierarchy. Similarly, since

$$statZ \phi(Z, \dots)$$

roughly translates as

$$\forall C \text{ } C \text{ is club } \implies \exists Z \in C \phi(Z, \dots),$$

the *stat* quantifier will correspond to the universal second-order quantifier.

Definition 2.1. *A formula in Stationary Logic without second order quantifiers will be denoted by Σ_0^1 or Π_0^1 . For $n > 0$, a formula of the form*

$$statZ_1 \dots statZ_k \phi(Z_1, \dots, Z_k, \dots)$$

where ϕ is Σ_{n-1}^1 will be called a Π_n^1 formula, and a formula of the form

$$aaZ_1 \dots aaZ_k \psi(Z_1, \dots, Z_k, \dots)$$

where ψ is Π_{n-1}^1 will be called a Σ_n^1 formula.

For example, if $\phi(Z_0, Z_1, v_1, \dots, v_\ell)$ has no *stat* or *aa* quantifiers, then

$$statZ_0 \ aaZ_1 \ \phi(Z_0, Z_1, v_1, \dots, v_\ell)$$

is a Π_2^1 formula.

Definition 2.2. *We say that the **LST property holds for Stationary Logic** iff for every structure $\mathfrak{A} = (A, \dots)$ ⁵ there exists a $W \subseteq A$ of size $\leq \omega_1$ such that for all formulas ϕ in Stationary Logic with no free occurrences of second order variables, and all first order parameters $p_1, \dots, p_k \in W$,*

$$\mathfrak{A} \models \phi[\vec{p}] \text{ if and only if } \mathfrak{A}|W \models \phi[\vec{p}].$$

*We say that the $\Pi_1^1 \downarrow$ **LST property holds for Stationary Logic** iff for every structure $\mathfrak{A} = (A, \dots)$ there exists a $W \subseteq A$ of size $\leq \omega_1$ such that for all Π_1^1 formulas ϕ in Stationary Logic with no free occurrences of second order variables, and all first order parameters $p_1, \dots, p_k \in W$,*

$$\text{if } \mathfrak{A} \models \phi[\vec{p}], \text{ then } \mathfrak{A}|W \models \phi[\vec{p}].$$

Remark 2.3. *Note that in the definition of the $\Pi_1^1 \downarrow$ LST property, we only require that Π_1^1 formulas reflect **downward**. If there is always an ω_1 sized substructure that reflects Π_1^1 formulas both upward and downward, then the full LST property holds. This issue is discussed further in Section 4.*

⁵Recall we always assume countable signature, though for everything discussed in this paper an ω_1 -sized signature would still be fine.

We consider variants of the **Diagonal Reflection Principle** introduced in Cox [4] and [2]. We use the following definition, which by Cox-Fuchs [5] is equivalent to the definitions from [4] and [2]:

Definition 2.4. $DRP_{internal}$ asserts that for every sufficiently large regular θ , there are stationarily many $W \in \wp_{\omega_2}(H_\theta)$ such that:

- $|W| = \omega_1 \subset W$; and
- Whenever $A \in W$ is uncountable and $S \in W$ is a stationary subset of $[A]^\omega$, the set $S \cap W \cap [W \cap A]^\omega$ is stationary in $[W \cap A]^\omega$.

The “internal” part of the definition refers to the fact that we require that $S \cap W \cap [W \cap A]^\omega$ is stationary, not merely that $S \cap [W \cap A]^\omega$ is stationary. Definition 2.4 is simply the diagonal version of an internal variant of WRP introduced in Fuchino-Usuba [8] (see Cox [2] for a discussion).

3. PROOF OF THEOREM 1.2

We prove a slightly stronger variant of Theorem 1.2. The proof below is strongly influenced by Fuchino et al [7].

Theorem 3.1. *The following are equivalent:*

- (1) $DRP_{internal}$.
- (2) For every structure $\mathfrak{A} = (A, \dots)$, there is a $W \subseteq A$ of size at most ω_1 such that for every finite list $p_1, \dots, p_k \in W \cap A$ and every formula ϕ without 2nd order quantifiers,

$$\left(\mathfrak{A} \models \text{stat}Z \phi[Z, \vec{p}] \right) \implies \left(\mathfrak{A}|W \models \text{stat}Z \phi[Z, \vec{p}] \right).$$

- (3) The $\Pi_1^1 \downarrow$ -LST property holds of Stationary Logic (as in Definition 2.2);
- (4) For every structure $\mathfrak{A} = (A, \dots)$, there is a $W \subseteq A$ of size at most ω_1 such that for every formula ψ in 2nd order prenex form with no free occurrences of second order variables, and every finite list p_1, \dots, p_k of elements of W , **if**

$$\mathfrak{A} \models \psi[\vec{p}]$$

then, letting $\hat{\psi}$ be the formula obtained from ψ by changing all aa quantifiers to stat quantifiers,

$$\mathfrak{A}|W \models \hat{\psi}[\vec{p}].$$

Before proving the theorem, we remark that in parts 2, 3, and 4 of Theorem 3.1, we only mentioned first order parameters from $W \cap A$. If the structure \mathfrak{A} is sufficiently rich then it often makes sense to also speak of second-order parameters that are elements of W . But in general (e.g. when \mathfrak{A} is a group) it is more natural to only speak of first order parameters from $W \cap A$.

Proof. (of Theorem 3.1): (4) trivially implies (3), since if ψ is represented as a prenex Π_1^1 formula, then $\hat{\psi} = \psi$ (because there are no *aa* quantifiers in the original formula at all). Similarly, (3) trivially implies (2) because if ϕ has no second order quantifiers,

$$\text{stat}Z \phi$$

is obviously a Π_1^1 formula.

To see that (2) implies (1), assume (2) and suppose θ is a regular cardinal $\geq \omega_2$. We need to find a $W \prec (H_\theta, \in)$ such that $|W| = \omega_1 \subset W$ and for every $s \in W$ that is a stationary collection of countable sets,

$$s \cap W \cap \left[W \cap \bigcup s \right]^\omega \text{ is stationary.}$$

Consider $\mathfrak{A} = (H_\theta, \in)$. Let $W \subset H_\theta$ be as in the statement of (2). Fix any $s \in W$ that is a stationary collection of countable sets. Then

$$\mathfrak{A} \models \text{stat}Z \exists p p = Z \cap \bigcup s \text{ and } p \in s$$

and hence, since $s \in W$ and the only second order quantifier in the (prenex) formula above is a *stat* quantifier,

$$\mathfrak{A}|W \models \text{stat}Z \exists p p = Z \cap \bigcup s \text{ and } p \in s.$$

Unravelling the definition of the satisfaction relation, this means that

$$\{Z \in [W]^\omega : Z \cap \bigcup s \in W \cap s\} \text{ is stationary in } [W]^\omega$$

and it follows that $W \cap s \cap [W \cap \bigcup s]^\omega$ is stationary in $[W \cap \bigcup s]^\omega$.

To see that $\omega_1 \subset W$, it suffices to show that $W \cap \omega_1$ is uncountable (since by first-order elementarity of W in (H_θ, \in) , $W \cap \omega_1$ is transitive). Now

$$\mathfrak{A} \models \text{stat}Z \exists p \exists \alpha (p = Z \cap \omega_1, \alpha < \omega_1, \text{ and } \alpha \text{ is an upper bound of } p),$$

so by assumption on W , this statement is also satisfied by $\mathfrak{A}|W$ (note that the parameter ω_1 is an element of W because ω_1 is first-order definable in \mathfrak{A} and W is at least first-order elementary in \mathfrak{A}). If $W \cap \omega_1$ were countable, say $W \cap \omega_1 = \delta < \omega_1$, it would follow that for stationarily many $Z \in W \cap [W]^\omega$, there is an $\alpha < W \cap \omega_1 = \delta$ such that α is an upper bound of $Z \cap \delta$. This would be a contradiction, since due to the countability of δ , the set of $Z \in [W]^\omega$ such that $\delta \subseteq Z$ is a club.

Finally, to prove that (1) implies (4): fix a structure $\mathfrak{A} = (A, \dots)$ and let θ be a sufficiently large regular cardinal with $\mathfrak{A} \in H_\theta$. By (1) there is a $W \prec (H_\theta, \in, \mathfrak{A})$ witnessing $\text{DRP}_{\text{internal}}$. We prove by induction on complexity of formulas ψ in 2nd order prenex form that if $p_1, \dots, p_k \in W \cap A$ and

$$\mathfrak{A} \models \psi[\vec{p}]$$

then, letting $\hat{\psi}$ be the result of replacing all *aa* quantifiers with *stat* quantifiers,

$$\mathfrak{A}|(W \cap A) \models \hat{\psi}[\vec{p}].$$

We actually need to inductively prove a slightly stronger statement: namely, that whenever ψ is a 2nd order prenex formula, $p_1, \dots, p_k \in W \cap A$, and $Z_1, \dots, Z_\ell \in W \cap [A]^\omega$,

$$(1) \quad \mathfrak{A} \models \psi[\vec{Z}, \vec{p}] \implies \mathfrak{A} \upharpoonright (W \cap A) \models \hat{\psi}[\vec{Z}, \vec{p}].$$

So suppose

$$(2) \quad \mathfrak{A} \models QZ \phi[Z, U_1, \dots, U_k, p_1, \dots, p_\ell]$$

where Q is either the *aa* or *stat* quantifier, U_1, \dots, U_k are each elements of $W \cap [A]^\omega$, $p_1, \dots, p_\ell \in W \cap A$, and the inductive hypothesis holds of the formula ϕ .

Now regardless of whether Q is the *aa* or *stat* quantifier,

$$\widehat{QZ \phi} \equiv \text{stat}Z \hat{\phi}.$$

and by (2) (since the *aa* quantifier is stronger than the *stat* quantifier)

$$\mathfrak{A} \models \text{stat}Z \phi[Z, U_1, \dots, U_k, p_1, \dots, p_\ell].$$

Hence, by the definition of the stationary logic satisfaction relation,

$$s := \{Z \in [A]^\omega : \mathfrak{A} \models \phi[Z, \vec{U}, \vec{p}]\} \text{ is stationary in } [A]^\omega.$$

Note that since \vec{U} , \vec{p} , ϕ , and \mathfrak{A} are elements of W , it follows that $s \in W$. Since W is internally diagonally reflecting,

$$s \cap W \cap [W \cap A]^\omega \text{ is stationary in } [W \cap A]^\omega.$$

Consider for the moment an arbitrary $Z \in s \cap W \cap [W \cap A]^\omega$. Then

$$\mathfrak{A} \models \phi[Z, \vec{U}, \vec{p}]$$

and it follows by the induction hypothesis (and that Z , \vec{U} , and \vec{p} are each elements of W) that:

$$\mathfrak{A} \upharpoonright (W \cap A) \models \hat{\phi}[Z, \vec{U}, \vec{p}].$$

Hence, we have shown that

$$s \cap W \cap [W \cap A]^\omega \subseteq \{Z \in [W \cap A]^\omega : \mathfrak{A} \upharpoonright (W \cap A) \models \hat{\phi}[Z, \vec{U}, \vec{p}]\}.$$

Since the set on the left side is stationary, the set on the right side is too. So by the definition of the satisfaction relation,

$$\mathfrak{A} \upharpoonright (W \cap A) \models \text{stat}Z \hat{\phi}[Z, \vec{U}, \vec{p}].$$

This completes the proof of the (1) \implies (4) direction. □

4. CONCLUDING REMARKS

We remark that it is straightforward to show, in ZFC alone, that:

Lemma 4.1. *For every structure $\mathfrak{A} = (A, \dots)$ there exists a $W \subseteq A$ of size at most ω_1 such that*

$$\mathfrak{A}|W \prec_{\downarrow}^{\Sigma_1^1} \mathfrak{A}$$

(i.e. such that Σ_1^1 formulas satisfied by \mathfrak{A} are also satisfied by $\mathfrak{A}|W$).

In fact, if θ is a regular cardinal such that $\mathfrak{A} \in H_\theta$, and

$$W \prec_{1st\ order} (H_\theta, \in, \mathfrak{A})$$

is such that $|W| = \omega_1$ and

$$(3) \quad W \cap [W \cap A]^\omega \text{ contains a club in } [W \cap A]^\omega$$

(this always holds for stationarily many W , e.g. for those W that are internally approachable), then

$$\mathfrak{A}|(W \cap A) \prec_{\downarrow}^{\Sigma_1^1} \mathfrak{A}.$$

We briefly sketch the proof of the lemma; more details, and other related results, can be found in Cox [3]. One proves by induction on complexity of formulas, making use of (3), that if ϕ is Σ_1^1 , $p_1, \dots, p_k \in W \cap A$, and $Z_1, \dots, Z_\ell \in W \cap [A]^\omega$, then

$$\text{if } \mathfrak{A} \models \phi[\vec{Z}, \vec{p}], \text{ then } \mathfrak{A}|(W \cap A) \models \phi[\vec{Z}, \vec{p}].$$

This was basically part of the proof from Fuchino et al [7] that DRP_{IC} implied the LST for Stationary Logic. See [3] for some other related ZFC theorems.

So by Lemma 4.1 one can always get an ω_1 sized substructure that reflects all Σ_1^1 statements downward. And if $\text{DRP}_{\text{internal}}$ holds, one can *also* get an ω_1 sized substructure that reflects all Π_1^1 statements downward. But it is consistent that both of these are true, yet no *single* ω_1 -sized substructure downward reflects all Π_1^1 *and* all Σ_1^1 statements. In particular, in any model where $\text{DRP}_{\text{internal}}$ holds and DRP_{IC} fails, Theorem 1.2 tells us that there is a structure such that no ω_1 -sized substructure reflects all Π_1^1 and all Σ_1^1 statements (though there are structures that reflect one or the other).

Another way to view this phenomenon, in terms of DRP-like principles, is that $\text{DRP}_{\text{internal}}$ yields stationarily many $W \in \wp_{\omega_2}(H_\theta)$ such that the transitive collapse H_W of W is “correct about stationary sets”; i.e. whenever $s \in H_W$ and $H_W \models$ “ s is a stationary set of countable sets”, then V believes this too. However, if W is not internally club, it is possible (by [2]) that H_W is correct about stationary sets, but is *not* correct about clubs; i.e. there can be a $c \in H_W$ such that $H_W \models$ “ c is a club of countable sets”, but V does not believe this. If, on the other hand, W witnesses DRP_{IC} , then H_W is correct about *both* stationarity *and* clubness.

REFERENCES

- [1] Jon Barwise, Matt Kaufmann, and Michael Makkai, *Stationary logic*, Ann. Math. Logic **13** (1978), no. 2, 171–224, DOI 10.1016/0003-4843(78)90003-7. MR486629
 - [2] Sean Cox, *Forcing axioms, approachability at ω_2 , and stationary set reflection*, Journal of Symbolic Logic, to appear, available at <https://arxiv.org/abs/1807.06129>.
 - [3] ———, *Internal reflection and applications*, in preparation.
 - [4] ———, *The Diagonal Reflection Principle*, Proceedings of the American Mathematical Society **140** (2012), no. 8, 2893–2902.
 - [5] Sean Cox and Gunter Fuchs, in preparation.
 - [6] Sean Cox and Hiroshi Sakai, *Martin's Maximum and the Diagonal Reflection Principle*, under review, available at <https://arxiv.org/abs/1904.00262>.
 - [7] Sakaé Fuchino, André Rodrigues Ottenbreit Maschio, and Hiroshi Sakai, *Strong downward Löwenheim-Skolem theorems for stationary logics, I*, preprint (2018).
 - [8] Sakaé Fuchino and Toshimichi Usuba, *A reflection principle formulated in terms of games*, RIMS Kokyuroku **1895**, 37–47.
 - [9] Thomas Jech, *Set theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded. MR1940513 (2004g:03071)
 - [10] John Krueger, *Internal approachability and reflection*, J. Math. Log. **8** (2008), no. 1, 23–39.
 - [11] David W. Kueker, *Countable approximations and Löwenheim-Skolem theorems*, Ann. Math. Logic **11** (1977), no. 1, 57–103. MR0457191 (56 #15406)
 - [12] Saharon Shelah, *Generalized quantifiers and compact logic*, Trans. Amer. Math. Soc. **204** (1975), 342–364, DOI 10.2307/1997362. MR376334
- Email address:* `scox9@vcu.edu`

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, VIRGINIA COMMONWEALTH UNIVERSITY, 1015 FLOYD AVENUE, RICHMOND, VIRGINIA 23284, USA