

The Cauchy problem for the Navier-Stokes type equations on the Heisenberg group

Yasuyuki Oka

Abstract

The aim of this article is to give a résumé of already published paper [28]. Thus we introduce an existence and uniqueness result of solutions for the Cauchy problem of the Navier-Stokes type equations associated with the sublaplacian provided by the left invariant vector fields on the Heisenberg group. To avoid the difficulty of the non-commutative which is intrinsic in 2-step stratified Lie groups, we construct the solenoidal space by using the right invariant vector fields on the Heisenberg group.

1 2

1 Introduction and main results

In this article, we introduce without the proofs an existence and uniqueness result for the Cauchy problem of the Navier-Stokes type equations on the Heisenberg group in [28] as an analogy of them on the Euclidean space. For the Cauchy problem of the Navier-Stokes equations on the Euclidean space \mathbb{R}^3 ,

$$\begin{cases} \mathbf{u}_t - \Delta \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla \pi = 0, & x \in \mathbb{R}^3, t > 0, \\ \operatorname{div} \mathbf{u} = 0, & x \in \mathbb{R}^3, t > 0, \\ \mathbf{u}(x, 0) = \mathbf{a}, & x \in \mathbb{R}^3, \end{cases} \quad (1.1)$$

T. Kato [18] and Y. Giga [11] gave the following existence and uniqueness of local and global solutions of (1.1):

- If $\mathbf{a} \in L^3_\sigma$, there exists a unique solution $u \in C([0, T], L^3_\sigma)$ for $T > 0$. (1.2)
- Moreover if there exists a constant δ such that $\|\mathbf{a}\|_{L^3} \leq \delta$, we can take $T = \infty$ in (1.2).

¹*Keywords:* Heisenberg group, Navier-Stokes equations, Local solution, Global solution.

²**AMS Subject Classification 2010:** 35A01, 35A02.

After this result, this direction has been deeply investigated (for instance, [4], [20], [21], [22], [25] and so on). On the other hand, many studies of partial differential equations on 2-step stratified Lie groups have been done by many mathematicians (for instance, [1], [2], [3], [6], [7], [8], [9], [10], [12], [14], [15], [17], [24], [26], [27], [31], [32] and so on). Many of them, however, are linear partial differential equations. Recently it has started out even the results of nonlinear evolution equations (for instance, [5], [16], [26], [29], [30] and so on). The studies of partial differential equations on 2-step stratified Lie groups have become more difficult than the Euclidean cases by the non-commutative which is intrinsic in 2-step stratified Lie groups.

By the way, the following two results have led us to this investigation. One is Lewy equations. They strongly suggest that the solvability is characterized by the Lie group structure of the equation. The other is Hörmander's condition. It strongly suggests the existence of a differential operator on the group corresponding to the Laplacian in Euclidean space. Lewy equation was given from the investigation of differential equations without solutions in [23] and Hörmander's condition was given from the investigation of hypoellipticity in [13]. From these two results, we have the following question: What is the relationship between the structure of Lie groups and the solvability for nonlinear evolution equations? The motivation for our investigation is to (partially) clarify how groups affect solvability.

Here we prepare some notations needed to assert our results. Let $g = (x, y, s)$ and $g' = (x', y', s') \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} = \mathbb{R}^{2d+1}$. Then we consider \mathbb{R}^{2d+1} with the group law defined by

$$g \cdot g' = (x, y, s) \cdot (x', y', s') = (x + x', y + y', s + s' + 1/2(x' \cdot y - x \cdot y')), \quad (1.3)$$

where $x \cdot y = \sum_{j=1}^d x_j y_j$. The group \mathbb{R}^{2d+1} with respect to the group law defined by (1.3) is called the Heisenberg group and denoted by \mathbb{H}^d . For $\lambda > 0$, we define the dilation δ_λ by

$$\delta_\lambda(x, y, s) = (\lambda x, \lambda y, \lambda^2 s).$$

The left-invariant vector fields in the Heisenberg group \mathbb{H}^d as \mathbb{R}^{2d+1} are represented by

$$X_j = \frac{\partial}{\partial x_j} + \frac{1}{2} y_j \frac{\partial}{\partial s}, \quad X_{d+j} = \frac{\partial}{\partial y_j} - \frac{1}{2} x_j \frac{\partial}{\partial s} \quad \text{and} \quad X_{2d+1} = \frac{\partial}{\partial s}$$

for $j = 1, 2, \dots, d$ and these make a basis for the Lie algebra of \mathbb{H}^d . On the other hand, the right-invariant vector fields in \mathbb{H}^d are represented by

$$\tilde{X}_j = \frac{\partial}{\partial x_j} - \frac{1}{2} y_j \frac{\partial}{\partial s}, \quad \tilde{X}_{d+j} = \frac{\partial}{\partial y_j} + \frac{1}{2} x_j \frac{\partial}{\partial s} \quad \text{and} \quad \tilde{X}_{2d+1} = \frac{\partial}{\partial s}$$

for $j = 1, 2, \dots, d$ and these also make a basis for the Lie algebra \mathfrak{h} of \mathbb{H}^d . Put

$$V_1 = \{X_1, X_2, \dots, X_{2d}\} \quad (\dim V_1 = 2d (= Q_1)) \quad \text{and} \quad V_2 = \{X_{2d+1}\} \quad (\dim V_2 = 1 (= Q_2)).$$

Then $\mathfrak{h} = V_1 \oplus V_2$ and

$$\text{rank}(\text{Lie}\{X_1, X_2, \dots, X_{2d}\}(g)) = 2d + 1, \quad (\text{Hörmander's condition})$$

for any $g \in \mathbb{H}^d$. The homogeneous dimension N of \mathbb{H}^d is given by $N = 2d + 2$ (If $\mathbb{R}^{2d+1} = \mathbb{R}^{Q_1} \times \mathbb{R}^{Q_2} = \mathbb{R}^{2d} \times \mathbb{R}^1$, then $N = \sum_{i=1}^2 iQ_i$). Define the vector ∇ of the left-invariant vector fields by $\nabla = (X_1, X_2, \dots, X_{2d})$ and the vector $\tilde{\nabla}$ of the right-invariant vector fields by $\tilde{\nabla} = (\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_{2d})$. Moreover the divergence of the left invariant vector fields is denoted by div and the divergence of the right invariant vector fields is denoted by div_R , respectively. The sub-Laplacian \mathcal{L} and \mathcal{L}_R on \mathbb{H}^d are defined by

$$\mathcal{L} = - \sum_{j=1}^{2d} X_j^2$$

and

$$\mathcal{L}_R = - \sum_{j=1}^{2d} \tilde{X}_j^2,$$

respectively.

We consider the following Cauchy problem of the Navier-Stokes type equations,

$$\begin{cases} \mathbf{u}_t + \mathcal{L}\mathbf{u} + (\mathbf{u} \cdot \nabla)\mathbf{u} + \tilde{\nabla}\pi = 0, \\ \text{div}_R \mathbf{u} = 0, \\ \mathbf{u}(g, 0) = \mathbf{a}(g) \end{cases} \quad (1.4)$$

for $g \in \mathbb{H}^d$ and $t > 0$, limited to Lebesgue spaces on 2-step stratified Lie groups, especially Heisenberg group, instead of the Euclidean spaces.

Let $\mathbf{b} = (b_1, \dots, b_{2d}) \in L^p(\mathbb{H}^d)^{2d}$, $1 < p < \infty$, and set

$$\mathbf{P}\mathbf{b} = (P_1\mathbf{b}, \dots, P_{2d}\mathbf{b}), \quad P_j\mathbf{b} = b_j + \sum_{k=1}^{2d} R_j \bar{R}_k b_k$$

and

$$\mathbf{Q}\mathbf{b} = - \sum_{i=1}^{2d} \mathcal{L}_R^{-1} \tilde{X}_i b_i,$$

where $R_l = \tilde{X}_l \mathcal{L}_R^{-\frac{1}{2}}$ and $\bar{R}_l = \mathcal{L}_R^{-\frac{1}{2}} \tilde{X}_l$, $l = 1, \dots, 2d$. Then we decompose \mathbf{b} by

$$\mathbf{b} = \mathbf{P}\mathbf{b} + \tilde{\nabla}\mathbf{Q}\mathbf{b}.$$

If $\pi = -Q(\mathbf{u} \cdot \nabla)\mathbf{u}$, then (1.4) becomes

$$\begin{cases} \mathbf{u}_t + \mathcal{L}\mathbf{u} = -P(\mathbf{u} \cdot \nabla)\mathbf{u}, \\ \operatorname{div}_R \mathbf{u} = 0, \\ \mathbf{u}(g, 0) = \mathbf{a}(g). \end{cases} \quad (1.5)$$

If there exists a solution of (1.5), then by Proposition 5 in [28], a solution \mathbf{u} can be expressed by

$$\mathbf{u}(t) = S(t)\mathbf{a} - \int_0^t S(t-\sigma)P[(\mathbf{u} \cdot \nabla)\mathbf{u}](\sigma)d\sigma, \quad (1.6)$$

where $S(t)\mathbf{a} = (a_1 * h_t, a_2 * h_t, \dots, a_{2d} * h_t)$ and a solution of the integral equations (1.6) satisfies (1.5) in the sense of distributions.

In this article, we introduce the following theorems for an existence and uniqueness of local solutions and global solutions. The basic idea of the proof is similar to [18] and [11]. We refer to [28] for detail proofs.

Theorem 1 ([28]). *Let $N = 2d + 2 < p < \infty$. Then for an initial value $\mathbf{a} \in J^N(\mathbb{H}^d) = \{\mathbf{a} = (a_1, \dots, a_{2d}) \in L^N(\mathbb{H}^d)^{2d}, \operatorname{div}_R \mathbf{a} = 0\}$, there exists a positive constant T and the following holds:*

(i) (Existence) *The integral equation (1.6) has a solution $\mathbf{u}(t)$ satisfying the following conditions:*

$$\begin{aligned} & \cdot \mathbf{u}(t) \in C([0, T], L^N(\mathbb{H}^d)^{2d}), \quad \lim_{t \rightarrow 0^+} \|\mathbf{u}(t) - \mathbf{a}\|_N = 0, \\ & \cdot \operatorname{div}_R \mathbf{u}(t) = 0, \end{aligned} \quad (1.7)$$

$$\cdot \mathbf{u}(t) \in C((0, T), L^p(\mathbb{H}^d)^{2d}), \quad \lim_{t \rightarrow 0^+} \sup_{0 < \sigma < t} \sigma^{\frac{1}{2} - \frac{N}{2p}} \|\mathbf{u}(\sigma)\|_p = 0 \text{ and} \quad (1.8)$$

$$\cdot \nabla \mathbf{u}(t) \in C((0, T), L^N(\mathbb{H}^d)^{2d \times 2d}), \quad \lim_{t \rightarrow 0^+} \sup_{0 < \sigma < t} \sigma^{\frac{1}{2}} \|\nabla \mathbf{u}\|_N = 0. \quad (1.9)$$

(ii) (Uniqueness) *If $\mathbf{v}(t)$ satisfies (1.6), (1.7), (1.8) and (1.9), then $\mathbf{u}(t) = \mathbf{v}(t)$ for $0 \leq t < T$.*

Theorem 2 ([28]). *Let $N = 2d + 2 < p < \infty$.*

(i) (Existence) *If an initial value $\mathbf{a} \in J^N(\mathbb{H}^d)$ satisfies*

$$\|\mathbf{a}\|_N \leq C_{N,p}$$

for a positive constant $C_{N,p}$, the integral equation (1.6) has a solution $\mathbf{u}(t)$ satisfying the following conditions.

$$\begin{aligned} & \cdot \mathbf{u}(t) \in C([0, \infty), L^N(\mathbb{H}^d)^{2d}), \quad \lim_{t \rightarrow 0^+} \|\mathbf{u}(t) - \mathbf{a}\|_N = 0, \\ & \cdot \operatorname{div}_R \mathbf{u}(t) = 0, \end{aligned} \tag{1.10}$$

$$\cdot \mathbf{u}(t) \in C((0, \infty), L^p(\mathbb{H}^d)^{2d}), \quad \lim_{t \rightarrow 0^+} \sup_{0 < \sigma < t} \sigma^{\frac{1}{2} - \frac{N}{2p}} \|\mathbf{u}(\sigma)\|_p = 0, \tag{1.11}$$

$$\cdot \nabla \mathbf{u}(t) \in C((0, \infty), L^N(\mathbb{H}^d)^{2d \times 2d}), \quad \lim_{t \rightarrow 0^+} \sup_{0 < \sigma < t} \sigma^{\frac{1}{2}} \|\nabla \mathbf{u}\|_N = 0 \text{ and} \tag{1.12}$$

· there exists a positive constant C such that

$$\|\mathbf{u}(t)\|_p \leq Ct^{-\frac{1}{2} - \frac{N}{2p}}, \quad \|\nabla \mathbf{u}(t)\|_N \leq Ct^{-\frac{1}{2}}.$$

(ii) (Uniqueness) If $\mathbf{v}(t)$ satisfies the integral equation (1.6), (1.10), (1.11) and (1.12). Then $\mathbf{u}(t) = \mathbf{v}(t)$ for $0 \leq t < \infty$.

We find out the following from the main results:

- The integrability of the space of solutions depends on the dilation δ_λ and the homogeneous dimension N on the Heisenberg group, not the topological dimension.
- The dimension of the space of solutions depends on $\dim V_1$, that is, Hörmander's condition on the Heisenberg group.
- Our idea to avoid the difficulty of the non-commutative coming from the Heisenberg group is to use the right-invariant fields to construct the solenoidal space. Hence we use the left-invariant vector fields to construct the equations and use the right-invariant vector fields to construct the space of the initial value and solutions.

2 Future work

In this article, we have introduced an existence and uniqueness result of solutions for the Cauchy problem of the Navier-Stokes type equations associated with the sublaplacian on the Heisenberg group. As we have seen, they depend on the dilation, homogeneous dimension, Hörmander's condition and non-commutative on the Heisenberg group. What happens if we change groups? It is known that the Heisenberg group has a close relationship with the Gabor transform. On the other hand, it is also known that NA group (or $ax + b$ group) has a close relationship with the wavelet transform (for instance, we refer to [19]). So it is natural to consider the NA group after the consideration on the Heisenberg group from the point of view of time-frequency analysis. Since the Heisenberg group is a nilpotent Lie group, that is, it is a unimodular Lie group, but NA group is a solvable Lie group,

that is, it is a non-unimodular Lie group, the situation on NA group is considerably more complicated than on the Heisenberg group. However, we consider that it is an interesting subject to clarify the relationship between the properties of non-unimodular Lie groups and the solvability of nonlinear evolution equations, which are not encountered as far as nilpotent Lie groups (for example the Heisenberg group, the diamond group, stratified Lie groups and graded Lie groups and so on) are considered.

References

- [1] H. Bahouri, P. Gérard et C. -J. Xu, *Espaces de Besov et estimations de Strichartz généralisées sur le groupe de Heisenberg*, J. Anal. Math., 82 (2000), 93-118.
- [2] H. Bahouri, C. Fermanian and I. Gallagher, *Dispersive estimates for the Schrödinger operator on step-2 stratified Lie groups*, Anal. PDE 9 (2016), no. 3, 545–574.
- [3] A. Bonfiglioli and F. Uguzzoni, *Nonlinear Liouville theorem for some critical problem on H-type groups*, Journal of Functional Analysis 207, (2004), 161-215.
- [4] J. Bourgain and N. Pavlović, *Ill-posedness of the Navier-Stokes equations in a critical space in 3D*, J. Funct. Anal. 225, (2008), 2233–2247.
- [5] T. Bruno, M. M. Peloso, A. Tabacco and M. Vallarino, *Sobolev spaces on Lie groups: embedding theorems and algebra properties*, J. Funct. Anal. 276 (2019), no. 10, 3014–3050.
- [6] G. Furioli and A. Veneruso, *Strichartz inequalities for the Schrödinger equation with the full Laplacian on the Heisenberg group*, Studia Math. 160 (2004), no. 2, 157–178.
- [7] G. B. Folland, *Subelliptic estimates and function spaces on nilpotent Lie group*, Ark. Mat. 13 (1975), 161–207.
- [8] N. Garofalo and E. Lanconelli, *Existence and nonexistence results for semilinear equations on the Heisenberg group*, Indiana Univ. Math. J. 41 (1992), no. 1, 71–98.
- [9] N. Garofalo and D. Vassilev, *Symmetry properties of positive entire solutions of Yamabe-type equations on groups of Heisenberg type*, Duke Math. J. 106 (2001), no. 3, 411–448.
- [10] B. Gaveau, *Principe de moindre action, propagation de la chaleur, et estimées sous elliptiques sur certains groupes nilpotents*, Acta Math. 139, (1977), 95–153.
- [11] Y. Giga, *Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system*, J. Differential Equations 62 (1986), 186–212.

- [12] M. D. Hierro, *Dispersive and Strichartz estimates on H-type groups*, Studia Mathematica 169 (1), (2005), 1–20.
- [13] L. Hörmander, *Hypoelliptic second order differential equations*, Acta Math. 119 (1967), 147–171.
- [14] A. Hulanicki, *The distribution of energy in the Brownian motion in the Gaussian field and analytic hypoellipticity of certain subelliptic operators in the Heisenberg group*, Studia Math. 56, (1976), 165–173.
- [15] D. S. Jerison and A. Sánchez-Calle, *Estimates for the heat kernel for a sum of squares of vector fields*, Indiana Univ. Math. J., 35, (1986), 835–854.
- [16] M. Jleli, M. Kirane and B. Samet, *A Fujita-type theorem for a multitime evolutionary p -Laplace inequality in the Heisenberg group*, Electron. J. Differential Equations (2016), Paper No. 303, 1–8 pp.
- [17] A. Kaplan, *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc., 258, (1980), 147–153.
- [18] T. Kato, *Strong L^p -solutions of the Navier-Stokes equation in \mathbb{R}^m , with applications to weak solutions*, Math. Z. 187, (1984), 471–480.
- [19] T. Kawazoe, Harmonic Analysis on Groups, *Asakura-syoten*, (2000) (in Japanese).
- [20] H. Koch and D. Tataru, *Well-posedness for the Navier-Stokes equations*, Adv. Math. 157, (2001), 22–35.
- [21] H. Kozono, T. Ogawa and T. Taniuchi, *The critical Sobolev inequalities in Besov spaces and regularity criterion to some semi-linear evolution equations*. Math. Z. 242, (2002), 251–278.
- [22] H. Kozono and Y. Shimada, *Bilinear estimates in homogeneous Triebel-Lizorkin spaces and the Navier-Stokes equations*, Math. Nach. 276, (2004), 63–74.
- [23] H. Lewy, *An example of a smooth linear partial differential equation without solution*, Ann. of Math. (2) 66 (1957), 155–158.
- [24] H. Liu and M. Song, *Strichartz inequalities for the Schrödinger equation with the full Laplacian on H-type groups*, arXiv:1402.4311v3, 12 Mar (2015).
- [25] H. Miura, *Remark on uniqueness of mild solutions to the Navier-Stokes equations*, J. Funct. Anal. 218, (2005), 110–129.

- [26] Y. Oka, *Local well-posedness for semilinear heat equations on H type groups*, Taiwanese J. Math. 22 (2018), no. 5, 1091–1105.
- [27] Y. Oka and K. Yoshino, *Solvability of Lewy equation in the space of the tempered ultrahyperfunctions*, Journal of Pseudo-Differential Operators and Applications: Volume 3, Issue 3 (2012), 321–328 ((DOI) 10.1007/s11868-012-0052-y).
- [28] Y. Oka, *An existence and uniqueness result for the Navier-Stokes type equations on the Heisenberg group*, J. Math. Anal. Appl. 473 (2019), no. 1, 382–407.
- [29] S. Pohozaev and L. Véron, *Nonexistence results of solutions of semilinear differential inequalities on the Heisenberg group*, Manuscripta Math. 102 (2000), no. 1, 85–99.
- [30] M. Ruzhansky and N. Yessirkegenov, *Existence and non-existence of global solutions for semilinear heat equations and inequalities on sub-riemannian manifolds, and Fujita exponent on unimodular Lie groups*, arXiv: 1810.01933v2 [math.AP] 6 Dec 2018.
- [31] E. M. Stein, Harmonic Analysis, *Princeton University Press*. Princeton, N.J., (1993).
- [32] N. Th. Varopoulos, L. Saloff-Coste and T. Coulhon, Analysis and geometry on groups, *Cambridge Tracts in Mathematics, 100*. Cambridge University Press, Cambridge, (1992).

School of Liberal Arts and Sciences
Daido University
10-3 Takiharū-cho, Minami-ku, Nagoya 457-8530
JAPAN
E-mail address: y-oka@daido-it.ac.jp