

On triality relations of normal triality algebras and Lie algebras

– ある代数系の三対原理とリー代数–

Noriaki Kamiya (神谷徳昭)

University of Aizu (福島県公立大学法人会津大学)

概要: この小論ではリー代数とジョルダン代数を含むある代数系 (called a normal triality algebra) の三対原理, つまり自己同型群と微分の一般化について論究します. そしてその代数系から root systems, Cartan matrix の concept を用いなくて例外型の単純リー代数をいくつか構成する方法に応用します.

§0. Introduction (motivation), §1. Definitions of triality relations and normal triality algebras, §2. Examples of triality relations of algebras, §3. Lie algebras construction, Appendix (対称的合成代数の基底と三対原理), References.

このような内容について, 具体的実例を含め述べさせていただきます (又, これらの section は独立に読める様に心がけたつもりです).

§0. Introduction (motivation)

我々の三対原理の concept を複素数 \mathbf{C} のとき考えます. \mathbf{C} の普通の積を $*$, 共役元を $\bar{x} = a - ib (x = a + ib, a, b \text{ は実数}, i = \sqrt{-1})$ とするとき, new product を $xy = \bar{x} * \bar{y}$ によって定義します. この new product は非結合的代数系であり

$$g(xy) = (gx)(gy), \quad d(xy) = (dx)y + x(dy)$$

なる自己同型 g と微分 d を考える. \mathbf{C} を $re^{i\theta}$ で表示, 周期 2π で考察すると

$$\text{Aut } \mathbf{C} = \{e^{i\theta} | \theta = \frac{2\pi}{3}n, n : \text{integer}\} = \{1, \frac{-1 + \sqrt{3}i}{2}, \frac{-1 - \sqrt{3}i}{2}\} \cong S_3$$

$$\text{Der } \mathbf{C} = \{d \in \text{End } \mathbf{C} | d = \frac{2\pi}{3}in, n : \text{integer}\} = \{0, \frac{2\pi i}{3}, \frac{4\pi i}{3}\}$$

が成り立ちます. **product (積)** に注意してください.

$j = 0, 1, 2$ に対して (自然数 j を mod 3 で考えます), $\text{End } \mathbf{C}$ の元で,

$$g_j(xy) = (g_{j+1}x)(g_{j+2}y), \quad d_j(xy) = (d_{j+1}x)y + x(d_{j+2}y)$$

となる g_j, d_j を考えると, 次のような $\text{Aut } \mathbf{C}, \text{Der } \mathbf{C}$ の一般化が得られます.

$$\text{Trig } \mathbf{C} = \{(g_0, g_1, g_2) | g_j = \exp(\sqrt{-1} \alpha_j), \alpha_0 + \alpha_1 + \alpha_2 = 0, \alpha_j \in \text{Re } \mathbf{C}\},$$

$$\text{Trid } \mathbf{C} = \{(d_0, d_1, d_2) | d_j = \sqrt{-1} \alpha_j, \alpha_0 + \alpha_1 + \alpha_2 = 0, \alpha_j \in \text{Re } \mathbf{C}\}.$$

次に $n \times n$ 次の行列代数 A で考察すると, $j = 0, 1, 2$ に対して

$$\sigma_j(a)x = a_{j+1} * x * {}^t a_{j+2}, \quad d_j(p)x = p_{j+1} * x - x * p_{j+2}.$$

ただし $a_j \in A_0^* := \{b \in A | b * {}^t b = Id_n\}, p_j \in \text{Alt}(A) := \{c \in A | {}^t c = -c\},$

($x * y$ は standard product of matrix, ${}^t x$ は transpose of matrix),

とします. ここで $xy = {}^t(x * y)$ と **new product** を定義すると, この積 xy は非結合的代数ですが, 自己同型と微分の一般化が得られます;

$\sigma_j(a)(xy) = (\sigma_{j+1}(a)x)(\sigma_{j+2}(a)y)$, $d_j(p)(xy) = (d_{j+1}(p)x)y + x(d_{j+2}(p)y)$, $p_j = (1 - a_j) * (1 + a_j)^{-1}$ (Cayley transformation) が成り立ちます.

Remark. 8 元数の三対原理については ([S],[T],[Tô] and [K.3] 等) を参照.

Remark. Note that for an algebra A , if $\sqrt{3} \in F$ (基礎体), and $\dim_F A = 2$, then $\text{Aut} A \cong S_3$ (3 次の対称群), if $\dim_F A = 1$, then $\text{Trig} A \cong K_4$ (Klein's 4 group) ([K-O.3], [K.2]). これらの概念を一般の非結合的代数で以下考えます.

§1. Definitions of triality relations and normal triality algebras

この章で非結合的代数系における一般的な triality relations (local and global の cases) を述べさせていただきます. (a correspondence of local and global triality relations)

Let A be a nonassociative algebra over $ch F \neq 2, 3$ (algebra は有限次元そして必ずしも単位元をもたない非結合的代数系の場合を考えます).

Following ([K-O.2] or [K-O.3]), suppose that a triple $g = (g_1, g_2, g_3) \in (\text{Epi } A)^3$ satisfies a global triality relation

$$g_j(xy) = (g_{j+1}x)(g_{j+2}y) \quad (1.1)$$

where the index j is defined by modulo 3, so that $g_{j\pm 3} = g_j$ (this is said to be a triality group). We denote

$$\text{Trig } A = \{g = (g_1, g_2, g_3) \in (\text{Epi } A)^3 | g_j(xy) = (g_{j+1}x)(g_{j+2}y), \forall j = 1, 2, 3\} \quad (1.2)$$

This is a generalization of the automorphism group of A .

In contrast to the algebra triality relations (1.1), we may also consider a local triality relation

$$t_j(xy) = (t_{j+1}x)y + x(t_{j+2}y). \quad (1.3)$$

Analogously to (1.2) if $[t_j, t_k]$ is closed, we introduce

$$\text{Trid } A = \{t = (t_1, t_2, t_3) \in (\text{End } A)^3 | t_j(xy) = (t_{j+1}x)y + x(t_{j+2}y), \forall j = 1, 2, 3\}. \quad (1.4)$$

Then, it defines a Lie algebra with component wise commutation relation. Also if $(t_1, t_2, t_3) \in \text{Trid } A$, it is easy to verify that we have for any $\alpha_j \in F$, $\alpha_{j\pm 3} = \alpha_j$

$$t' = (t'_1, t'_2, t'_3) \in \text{Trid } A, \text{ where } t'_j = \sum_{k=1}^3 \alpha_{j-k} t_k \quad (j = 1, 2, 3).$$

Furthermore, if the exponential map $t_j \rightarrow \xi_j$ is given by

$$\xi_j = \exp t_j = \sum_{n=0}^{\infty} \frac{1}{n!} (t_j)^n \quad (1.5)$$

is well-defined, then we can show that

$$\xi_j(xy) = (\xi_{j+1}x)(\xi_{j+2}y), \quad (1.6)$$

provided that $t = (t_1, t_2, t_3) \in \text{Trid } A$ and vice-versa.

Next we introduce multiplication operators of A , $L(x), R(x) \in \text{End } A$ by

$$L(x)y = xy \text{ and } R(x)y = yx.$$

Def.1.1. Let $d_j(x, y) \in \text{End } A$, for $x, y \in A$ ($j = 1, 2, 3$) be to satisfy

(i)

$$d_1(x, y) = R(y)L(x) - R(x)L(y) \quad (1.7a)$$

$$d_2(x, y) = L(y)R(x) - L(x)R(y). \quad (1.7b)$$

(ii) The explicit form for $d_3(x, y)$ is unspecified except for

$$d_3(x, y) = -d_3(y, x), \quad (1.7c)$$

(iii)

$$(d_1(x, y), d_2(x, y), d_3(x, y)) \in \text{Trid } A.$$

We call the algebra A satisfying these conditions to be a *regular triality algebra*.

Remark. Any Lie (resp. Jordan) algebra with product $[x, y]$ (resp. xy) is an example of the regular (in particular, normal) triality algebra with respect to the $L([x, y]) = d_j(x, y)$ (resp. $L(x), L(y)$) for $j = 1, 2, 3$.

Proposition 1.2 ([K-O.3]). *Let A be a regular triality algebra satisfying either the condition (B) or (C); (B) $AA = A$, (C) if some $b \in A$ satisfies either $L(b) = 0$ or $R(b) = 0$, then $b = 0$. Then we obtain the following.*

(i) For any $t = (t_1, t_2, t_3) \in \text{Trid } A$, we have

$$[t_j, d_k(x, y)] = d_k(t_{j-k}x, y) + d_k(x, t_{j-k}y) \quad (1.8a)$$

Especially, if we choose $t_j = d_j(x, y)$ it yields

$$[d_j(u, v), d_k(x, y)] = d_k(d_{j-k}(u, v)x, y) + d_k(x, d_{j-k}(u, v)y). \quad (1.8b)$$

(ii) For any $g = (g_1, g_2, g_3) \in \text{Trig } A$, we have

$$g_j d_k(x, y) g_j^{-1} = d_k(g_{j-k}x, y) + d_k(x, g_{j-k}y). \quad (1.8c)$$

Let A be a regular triality algebra with either (B) or (C), and set

$$\mathfrak{D} = \text{span} \langle d_j(x, y), \forall x, y \in A, j = 1, 2, 3 \rangle. \quad (1.9)$$

Then \mathfrak{D} is a Lie algebra by (1.8b). Moreover, it is an ideal of the large Lie algebra $\text{Trid } A$ by (1.8a), denoted by $\mathfrak{D} \triangleleft \text{Trid } A$. We call an "inner triality derivation" (naming of the author) this \mathfrak{D} as a generalization of derivations.

Def.1.3. If a regular triality algebra satisfies Eqs.(1.7) as well as

$$d_3(x, y)z + d_3(y, z)x + d_3(z, x)y = 0, \quad (1.10a)$$

$$[d_j(u, v), d_k(x, y)] = d_k(d_{j-k}(u, v)x, y) + d_k(x, d_{j-k}(u, v)y), \quad (1.10b)$$

then we call a *pre normal triality algebra* ([K-O.1]). Furthermore, if we have

$$Q(x, y, z) := d_1(z, xy) + d_2(y, zx) + d_3(x, yz) = 0, \quad (1.11)$$

then A is called a *normal triality algebra* ([K-O.2]). Next we introduce the second bilinear product in the same vector space A with involution $x \rightarrow \bar{x}$ by

$$x * y = \overline{xy} = \bar{y} \bar{x}. \quad (1.12)$$

Then the resulting algebra $(A, x * y)$ is said to be a *conjugation algebra* of A , for the new product $x * y$, by means of $\overline{Qx} = \overline{Q} \bar{x}$ and $Q \in \text{End } A$, we have

$$\bar{g}_j(x * y) = (g_{j+1}x) * (g_{j+2}y), \quad \bar{d}_j(x * y) = (d_{j+1}x) * y + x * (d_{j+2}y). \quad (1.13)$$

Remark ([K-O.1]). The conjugation algebra of a structurable algebra which contains an alternative algebra is a normal triality algebra.

Note that the vector space $\mathfrak{A}_0 \otimes \mathfrak{J}_0$ with 182 dimension ([S]) appeared Tits second construction of the Lie algebra of type E_8 is a normal triality algebra (see next section for the details).

Theorem 1.4 ([K-O.2]). *The symmetric composition algebra, Lie and Jordan algebras are a normal triality algebra.*

Theorem 1.5. *For a normal triality algebra A , if we define*

$$\xi_j = \exp d_j \quad (j = 1, 2, 3), \quad (\text{assuming the well - defined})$$

then we have

$$\xi_j(xy) = (\xi_{j+1}x)(\xi_{j+2}y), \quad \text{that is, } (\xi_j, \xi_{j+1}, \xi_{j+2}) \in \text{Trig } A,$$

$$\left[\frac{d}{dt} ((\exp td_j) d_k (\exp td_j)^{-1}) \right]_{t=0} = [d_j, d_k] \in \text{Trid } A.$$

That is, this means that $([d_j, d_k], [d_{j+1}, d_{k+1}], [d_{j+2}, d_{k+2}]) \in \text{Trid } A$.

Corollary. *For the pseudo octonion or para Hurwitz algebras, the same result in Theorem 1.5 holds, as these algebras are a symmetric composition algebra and so a normal triality algebra.*

Remark. In the normal triality algebra A , if we define an endomorphism by $D(x, y) := d_1(x, y) + d_2(x, y) + d_3(x, y)$, then we have the relations $D(x, y) = -D(x, y)$, $D(xy, z) + D(yz, x) + D(zx, y) = 0$ and $D(x, y)$ is a derivation satisfying $[D(x, y), D(u, v)] = D(D(x, y)u, v) + D(u, D(x, y)v)$, thus this algebra A is a generalized structurable algebra ([K.1]).

Remark[K-O.1]. The conjugation algebra $(A, x * y)$ of normal triality algebra (A, xy) with a para unit e (i.e., $ex = xe = \bar{x}$) is a structurable algebra with the unit $e * x = x * e = x$, since $x * y = \overline{xy}$ and $\overline{\bar{x}} = x$.

Remark. Let $(A, x * y)$ be an associative algebra. Then $(A, x \cdot y)$ is a Jordan algebra with new product and involution defined by $x \cdot y = x * y + y * x$ and $\bar{x} = x$, since they satisfy the identities $x \cdot y = y \cdot x$ and $(x \cdot y) \cdot x^2 = x \cdot (y \cdot x^2)$.

Remark. Let A be a normal triality alg. For $(\xi_1, \xi_2, \xi_3) = (\exp d_1, \exp d_2, \exp d_3) \in \exp \mathfrak{D}$, provided that the exponential map is well-defined,

$$\forall g = (g_1, g_2, g_3) \in Trig A \implies$$

$$g_j \xi_k g_j^{-1} = g_j (\exp d_k) g_j^{-1} = \exp(g_j d_k g_j^{-1}) \in \exp \mathfrak{D}. \text{ (by (1.8) and (1.9))}$$

Therefore $G_0 = \langle \xi_1, \xi_2, \xi_3 \rangle_{span}$ is an invariant subgroup of $Trig A$. We call an "inner triality group" (naming of the author) and we obtain $\mathfrak{D} \longleftrightarrow G_0$.

For the definitions of this section, we would like to refer ([K-O.1],[K-O.2] and [K-O.3]), that is, for the concept of normal triality algebras and related topics.

It seems that this concept (called a normal triality algebra) of a generalization of the derivation may be regarded as a generalization in the "principle of triality" of the octonion (or para octonion) algebra due to Tits ([K.3]).

§2. Examples of triality relations of algebras

For several examples of normal triality algebras, in particular, to construct Lie algebras, we will exhibit them in this section.

First, it is known that a symmetric composition algebra A over a field F ($ch F \neq 2$) with a symmetric bilinear form $\langle x, y \rangle$ satisfying the relations

$$(xy)x = x(yx) = \langle x, x \rangle y, \quad \langle x, y \rangle \text{ is non-degenerate,}$$

is a normal triality algebra and the symmetric composition algebra is either a para-Hurwitz algebra (that is, the conjugation algebra is a Hurwitz algebra which contains the Cayley number if $ch F = 0$), or an eight dimensional pseudo octonion algebra due to M.Gell-mann ([G],[K-O.2],[K-O.3], [K3] and [O]).

This symmetric composition algebra A has a triality derivation $(d_0, d_1, d_2) \in Tri d A$ defined by $d_0(x, y) = 2\{[L(x), L(y)] - R([x, y])\}$,

$$d_1(x, y) = R(y)L(x) - R(x)L(y), \text{ and } d_2(x, y) = L(y)R(x) - L(x)R(y). \quad (\spadesuit)$$

The case of the symmetric composition algebra;

$$\langle d_j(a, b) \rangle_{span} \cong D_4 \text{ (Lie algebra of 28 dimension),}$$

since $\langle d_j(a, b)x, y \rangle + \langle x, d_j(a, b)y \rangle = 0, (\forall j = 0, 1, 2)$.

Secondarily, the Lie algebra ($\varepsilon = -1$) and Jordan algebra ($\varepsilon = 1$) are a normal triality algebra equipped with the product $L(x)y = xy = \varepsilon yx = \varepsilon R(x)y$ and the triality derivation $d(x, y) := d_0(x, y) = d_1(x, y) = d_2(x, y) = -\varepsilon[L(x), L(y)]$.

The other examples are the following:

Example a ([K-O.2]). The vector space $O \otimes O$ with 64 dimension induced from two para octonion algebras O (para octonion is the conjugation algebra of octonion \mathbf{O}) is a normal triality algebra with respect to the $D_j(a \otimes x, b \otimes y) := d_j^{(1)}(a, b) \otimes \langle x, y \rangle id + \langle a, b \rangle id \otimes d_j^{(2)}(x, y)$, where $d_j^{(1)}$ and $d_j^{(2)}$ with the triality derivation defined by (\spadesuit) respectively. Note that $D = D_0 + D_1 + D_2$ construct a Lie algebra of type $G_2 \oplus G_2$. This implies that $Der O \cong G_2$ (Lie algebra of 14 dim) in particular.

As we will show in next section, this case is relevant for a construction of so-called Freudenthal's magic square.

Remark. A vector of the tensor product $O_p \otimes O_p$ induced from the pseudo octonion algebra O_p is a normal triality algebra with $D_j(a \otimes x, b \otimes y) = d_j^{(1)}(a, b) \otimes \langle x, y \rangle + id + \langle a, b \rangle id \otimes d_j^{(2)}(x, y)$ as well as the tensor product of the para octonion algebra O . However note that $D = D_0 + D_1 + D_2$ construct a Lie algebra of type $A_2 \oplus A_2$. Hence this means that $Der O_p \cong A_2$ (Lie algebra of 8 dim).

Example b) ([K-O.1], [K-O.3]). For the octonion algebra \mathbf{O} and a para Zorn vector matrix with 56 dimension induced from the vector space

$$\begin{pmatrix} \alpha & \mathbf{a} \\ \mathbf{b} & \beta \end{pmatrix} \quad (\clubsuit)$$

is a normal triality algebra, where $\forall \mathbf{a}, \mathbf{b} \in H_3(\mathbf{O})(= \mathfrak{J})$ is a exceptional Jordan algebra of degree three with 27 dimension over a field F , and $\alpha, \beta \in F$ (the base field). For the details, see section 5 in ([K-O.3]). This concept was appeared the meta symplectic geometry due to H. Freudenthal (for example, see ([K-O.4])).

Example c). The vector space $\mathfrak{A}_0 \otimes \mathfrak{J}_0$ with 182 dimension appeared Tits second construction of $E_8([S], [T\delta])$ is a normal triality algebra with the (triality) derivation $D_0 = D_1 = D_2 (= D)$. Here this derivation of $\mathfrak{A}_0 \otimes \mathfrak{J}_0$ means that $D = (d_0 + d_1 + d_2) \otimes \langle x, y \rangle + id + \langle a, b \rangle id \otimes [R(x), R(y)]$, where d_j is the triality derivation of the normal triality algebra \mathfrak{A} (as the para octonion algebra) and $[R(x), R(y)]$ is the derivation of the normal triality algebra \mathfrak{J} (as the exceptional Jordan algebra with $dim \mathfrak{J} = 27$). Here denote by $\mathfrak{A}_0 = \{a \in \mathfrak{A} | trace(a) = 0\}$, $\mathfrak{J}_0 = \{x \in \mathfrak{J} | Trace(x) = 0\}$, and $dim \mathfrak{A}_0 = 7$, $dim \mathfrak{J}_0 = 26$ respectively.

§3. Lie algebras construction

In this section, we will discuss a construction of Lie algebras associated with the normal triality algebras of Examples a), b) and c) in section 2.

Following ([K-O.2]), let A be a normal triality algebra and consider linear maps:

$$\rho_j : A \rightarrow V, \text{ and } T_j : A \otimes A \rightarrow V \quad (3.1)$$

for $j = 0, 1, 2$, where V is an unspecified algebras with skew symmetric bi-linear product $[\circ, \circ]$. We set now

$$T(A, A) = span \langle T_j(x, y), \forall x, y \in A \rangle, \forall j = 0, 1, 2. \quad (3.2)$$

$$L(A) = \rho_0(A) \oplus \rho_1(A) \oplus \rho_2(A) \oplus T(A, A). \quad (3.3)$$

Let (i, j, k) be a cyclic permutation of indices $(0, 1, 2)$, and assume the following anti-commutative multiplication relations:

$$[\rho_i(x), \rho_i(y)] = -[\rho_i(y), \rho_i(x)] = T_{3-i}(x, y) \quad (3.4a)$$

$$[\rho_i(x), \rho_j(y)] = -[\rho_j(y), \rho_j(x)] = -\rho_k(xy) \quad (3.4b)$$

$$[T_i(x, y), \rho_j(z)] = -[\rho_j(z), T_i(x, y)] = \rho_j(d_{i+j}(x, y)z) \quad (3.4c)$$

$$\begin{aligned}
[T_l(u, v), T_m(x, y)] &= T_m(d_{l-m}(u, v)x, y) + T_m(x, d_{l-m}(u, v)y) \\
&= -T_l(d_{m-l}(x, y)u, v) - T_l(u, d_{m-l}(x, y)v)
\end{aligned} \tag{3.4d}$$

for $l, m = 0, 1, 2$. Hence, we introduce the Jacobian in $L(A)$ by

$$J(X, Y, Z) = [[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] \tag{3.5}$$

for $X, Y, Z \in L(A)$.

Lemma 3.1. $T(A, A)$ and $T_j(A, A)$ for $j = 0, 1, 2$ are Lie algebras. Also $T_j(A, A)$ is an ideal of $T(A, A)$, where $T_j(A, A) = \text{span} \langle T_j(x, y), \forall x, y \in A \rangle$.

Condition (D) Suppose that we have $\rho_j(x) = 0$ for some $x \in A$ and for some value of $j = 0, 1, 2$, we then have $x = 0$.

Proposition 3.2. Let A be a pre-normal triality algebra. If we have

$$J(x, y, z) = T_0(x, yz) + T_1(z, xy) + T_2(y, zx) = 0, \tag{3.6}$$

then $L(A)$ is a Lie algebra. Moreover, if the condition (D) holds, then A is a normal triality algebra. Conversely, if $L(A)$ is a Lie algebra and if the condition (D) holds, then A is a normal triality algebra with the validity of Eq.(3.6).

Theorem 3.3. Let A be a normal triality algebra. Then, the quotient algebra $\tilde{L}(A) = L(A)/J$ is a Lie algebra, where $J = \text{span} \langle J(x, y, z) \rangle$.

この定理 3.3 を適用して 5-graded $g_{-2} \oplus g_{-1} \oplus g_0 \oplus g_1 \oplus g_2$ の例外型リー代数を構成します. 特に normal triality algebra A の inner triality derivation d_j を T_j とみなす.

$$L(A) = \rho_1(A) \oplus \rho_2(A) \oplus \rho_3(A) \oplus T(A, A), \text{ and}$$

$$T(A, A) = \langle T_j(a, b), \forall j = 0, 1, 2, \forall a, b \in A \rangle_{\text{span}},$$

and we identify $T_j(a, b)$ with the triple $T_j(a, b) = (d_j(a, b), d_{j+1}(a, b), d_{j+2}(a, b))$.

§2 の Example a), b), c) の場合に付随した Lie algebras を考察する. 以後, 基礎体 F は標数 0 の代数閉体とする.

a) $A = \mathbf{O} \otimes \mathbf{O}$ の場合 (tensor product case, and $\dim g_{-2} = \dim g_2 = 14$): $A = A_1 \otimes A_2$, $\dim A_1, \dim A_2$ なる記号を用いると, それぞれ construct される Lie algebra は以下の様になります, ただし \mathbf{O} は octonion algebra.

$$L(A) \cong E_8, L_j = \rho_j(A) \oplus T_{3-j}(A, A) \cong D_8, T_j(A, A) \cong D_4 \oplus D_4.$$

$\dim A_2 \backslash \dim A_1$	1	2	4	8
1	A_1	A_2	C_3	F_4
2	A_2	$A_2 \oplus A_2$	A_5	E_6
4	C_3	A_5	D_6	E_7
8	F_4	E_6	E_7	E_8

E_8 を Extended Dynkin diagram で表すと $L(A)/L_j(A)$ は 128 次元の対称空間;

$\odot - \circ - \circ - \circ - \circ - \circ - \circ - \circ - \square$ \square omitted $\cong D_8$, and \odot is highest root.

|
○

Appendix (対称的合成代数の基底と三対原理).

Let A be a symmetric composition algebra over the field F and $\sqrt{3} \in F$.

$$\Sigma = \{a = (a_1, a_2, a_3) \in A^3 \mid a_j a_{j+1} = a_{j+2}, \langle a_j, a_j \rangle = 1, \forall j = 1, 2, 3\}. \quad (A.1)$$

For a given $a = (a_1, a_2, a_3) \in \Sigma$, we introduce a notation

$$\Lambda(a) = \{p = (p_1, p_2, p_3) \in A^3 \mid a_j p_{j+1} + p_j a_{j+1} = p_{j+2}, \langle p_j, a_j \rangle = 0, \forall j = 1, 2, 3\} \quad (A.2)$$

Note that $\Lambda(a)$ is a vector space over F .

Moreover, we define $q_j \in A$ by $q_j = a_{j+1} p_{j+2} = p_j - p_{j+1} a_{j+2}$.

From (Th.2.5 and Th.3.2 in [K-O.3]) and the notation being as above, we have the following theorems for global and local triality relations.

Theorem A.1. For $\forall a, b \in \Sigma$, we have $\sigma_j(a)$ and $\theta_j(a) \in \text{Trig } A$, $G = \langle \theta_j(a) \sigma_j(b) \text{ and } \sigma_j(a) \theta_j(b) \rangle_{\text{span}}$ is an invariant subgroup of $\text{Trig } A$, and $\sigma_{j+2}(a) \sigma_{j+1}(a) \sigma_j(a) = \text{Id}$, $\theta_{j+2}(a) \theta_{j+1}(a) \theta_j(a) = \text{Id}$, (inner triality group) where $\sigma_j(a) = R(a_{j+1}) R(a_{j+2})$ and $\theta_j(a) = L(a_{j+2}) L(a_{j+1})$.

Theorem A.2. For any $a \in \Sigma$ and $p \in \Lambda(a)$, if we introduce $D_j(a, p) \in \text{End } A$ by $D_j(a, p)x = (p_{j+1}x) a_{j+1} + a_j(x q_j)$, then this $D_j(a, p)$ satisfies

$$D_j(a, p)(xy) = (D_{j+1}(a, p)x)y + x(D_{j+2}(a, p)y) \quad (\text{inner triality derivation}).$$

Furthermore, we obtain that $T_j(A, A) = \langle D_j(a, p) \rangle_{\text{span}}$ is an ideal of $\text{Trid } A$.

また 8 次元の symmetric composition algebra の場合, u, v を適当に選ぶと

$D_j(a, p) \langle \dots \rangle d_j(u, v)$ (normal triality algebra における local triality derivation) の対応が知られています ([K-O.3]). Hence, $\langle d_j(u, v) \rangle_{\text{span}} \cong D_4$ (28 次元のリー代数) なので $D_j(a, p)$ を構成する 28 個の $a = (a_1, a_2, a_3)$ and $p = (p_1, p_2, p_3)$ を具体的に以下列挙させていただきます. これらの基底に関する乗積表は para and pseudo octonion algebras はそれぞれ次の表で, 内積は $x = \sum_j x_j e_j$, $y = \sum_j y_j e_j$ のとき $\langle x, y \rangle = \sum_j x_j y_j$ で定義されます.

(*) para octonion の基底: e_1, \dots, e_7 の積の例, $e_1 e_2 = -e_3, e_7 e_1 = -e_6, e_6 e_7 = -e_1$.

para case	e_1	e_2	e_3	e_4	e_5	e_6	e_7
e_1	-1	$-e_3$	e_2	$-e_5$	e_4	$-e_7$	e_6
e_2	e_3	-1	$-e_1$	e_6	$-e_7$	$-e_4$	e_5
e_3	$-e_2$	e_1	-1	$-e_7$	$-e_6$	e_5	e_4
e_4	e_3	$-e_6$	e_7	-1	$-e_1$	e_2	$-e_3$
e_5	$-e_4$	e_7	e_6	e_1	-1	$-e_3$	$-e_2$
e_6	e_7	e_4	$-e_5$	$-e_2$	e_3	-1	$-e_1$
e_7	$-e_6$	$-e_5$	$-e_4$	e_3	e_2	e_1	-1

(AI) Examples of triality group and derivation in the para octonion algebra (the conjugation algebra of Cayley algebra), この場合 $e_0 = 1$ なので省略.

$a = (a_1, a_2, a_3), p = [p_1, p_2, p_3]$ と表示する.

(AI):(para octonion algebra), with respect to basis e_1, \dots, e_7 .

(1) $a = (e_1, e_2, -e_3), p = [e_2, -e_1, 0]$,

Remark. $a = (a_1, a_2, a_3) \in \Sigma$ に対して $p_j = a_{j+1}p_{j+2} + p_{j+1}a_{j+2}$ を満たす $p = [p_1, p_2, p_3]$ の選び方はいくつか存在する. 例えば, $p = [e_3, 0, -e_1]$. $p = [0, e_3, -e_3]$ がある (これはある関数における微分係数の様に理解できます).

(2)($e_1, e_4, -e_5$), [$e_4, -e_1, 0$] (3)($e_1, e_6, -e_7$), [$e_6, -e_1, 0$] (4)(e_2, e_4, e_6), [$e_4, -e_2, 0$]
(5)($e_2, e_5, -e_7$), [$e_5 - e_2, 0$] (6)($e_3, e_4, -e_7$), [$e_4, -e_3, 0$] (7)($e_3, e_5, -e_6$), [$e_5, -e_3, 0$]
(8)($\frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2, \frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2, -1$), [$\frac{\sqrt{3}}{2}e_1 - \frac{1}{2}e_2, \frac{\sqrt{3}}{2}e_1 - \frac{1}{2}e_2, 0$], (i, j) = (1, 2) を
除いて (9)~(28): $i \neq j$, ($\frac{1}{2}e_i + \frac{\sqrt{3}}{2}e_j, \frac{1}{2}e_i + \frac{\sqrt{3}}{2}e_j, -1$), [$\frac{\sqrt{3}}{2}e_i - \frac{1}{2}e_j, \frac{\sqrt{3}}{2}e_i - \frac{1}{2}e_j, 0$]
つまり ($\frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7, \frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7, -1$), [$\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7, \frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7, 0$] etc. (8)~(28):
 ${}_{7}C_2 = 21$ 通り. where $e_0e_i = e_ie_0 = \bar{e}_i = -e_i$ ($i \neq 0$), e_0 is a para unit.

Note that the product $[e_j, e_k] := e_je_k - e_ke_j$, ($j, k = 1, \dots, 7$) makes a Malcev algebra with seven dimension ([O]).

(**) pseudo octonion の基底: e_1, \dots, e_8 の積の例, $e_1e_2 = e_3$, $e_8e_1 = e_1$, $e_8e_8 = -e_8$.

pseudo case	e_1	e_2	e_3	e_4
e_1	e_8	e_3	$-e_2$	$\frac{\sqrt{3}}{2}e_6 + \frac{1}{2}e_7$
e_2	$-e_3$	e_8	e_1	$\frac{1}{2}e_6 - \frac{\sqrt{3}}{2}e_7$
e_3	e_2	$-e_1$	e_8	$\frac{\sqrt{3}}{2}e_4 + \frac{1}{2}e_5$
e_4	$\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$	$-\frac{1}{2}e_6 - \frac{\sqrt{3}}{2}e_7$	$\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$	$\frac{\sqrt{3}}{2}e_3 - \frac{1}{2}e_8$
e_5	$\frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7$	$\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$	$\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	$-\frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8$
e_6	$\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$	$\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	$-\frac{\sqrt{3}}{2}e_6 + \frac{1}{2}e_7$	$\frac{\sqrt{3}}{2}e_1 - \frac{1}{2}e_2$
e_7	$\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	$-\frac{\sqrt{3}}{2}e_4 + \frac{1}{2}e_5$	$-\frac{1}{2}e_6 - \frac{\sqrt{3}}{2}e_7$	$-\frac{1}{2}e_1 - \frac{\sqrt{3}}{2}e_2$
e_8	e_1	e_2	e_3	$-\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$

pseudo case	e_5	e_6	e_7	e_8
e_1	$-\frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7$	$\frac{\sqrt{3}}{2}e_4 + \frac{1}{2}e_5$	$-\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	e_1
e_2	$\frac{\sqrt{3}}{2}e_6 + \frac{1}{2}e_7$	$-\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	$-\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$	e_2
e_3	$-\frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5$	$-\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$	$\frac{1}{2}e_6 - \frac{\sqrt{3}}{2}e_7$	e_3
e_4	$\frac{1}{2}e_3 + \frac{\sqrt{3}}{2}e_8$	$\frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2$	$\frac{1}{2}e_1 - \frac{\sqrt{3}}{2}e_2$	$-\frac{1}{2}e_4 - \frac{\sqrt{3}}{2}e_5$
e_5	$\frac{\sqrt{3}}{2}e_3 - \frac{1}{2}e_8$	$-\frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$	$\frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2$	$\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$
e_6	$\frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$	$-\frac{\sqrt{3}}{2}e_3 - \frac{1}{2}e_8$	$-\frac{1}{2}e_3 + \frac{\sqrt{3}}{2}e_8$	$-\frac{1}{2}e_6 - \frac{\sqrt{3}}{2}e_7$
e_7	$\frac{\sqrt{3}}{2}e_1 - \frac{1}{2}e_2$	$\frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8$	$-\frac{\sqrt{3}}{2}e_3 - \frac{1}{2}e_8$	$\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$
e_8	$-\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$	$-\frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7$	$-\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$	$-e_8$

(AII) Examples of triality group and derivation in the pseudo octonion algebra (Gell-Mann's eight dimensional matrix) この代数は単位元をもたない.

(AII):(pseudo octonion algebra), with respect to basis e_1, \dots, e_8 .

(1)(e_1, e_2, e_3), [$e_2, -e_1, 0$], (2)($e_1, e_4, \frac{\sqrt{3}}{2}e_6 + \frac{1}{2}e_7$), $p = [e_4, \frac{1}{2}e_1 - \frac{\sqrt{3}}{2}e_2, 0]$,
(3)($e_1, e_5, -\frac{1}{2}e_6 + \frac{\sqrt{3}}{2}e_7$), [$e_5, \frac{1}{2}e_1 - \frac{\sqrt{3}}{2}e_2, 0$], (4)($e_1, e_6, \frac{1}{2}e_5 + \frac{\sqrt{3}}{2}e_4$), [$e_6, \frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2, 0$], (5)($e_1, e_7, \frac{\sqrt{3}}{2}e_5 - \frac{1}{2}e_4$), [$e_7, \frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2, 0$], (6)($e_2, e_4 - \frac{\sqrt{3}}{2}e_7 + \frac{1}{2}e_6$),
 $[e_4, \frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2, 0]$, (7)($e_2, e_5, \frac{1}{2}e_7 + \frac{\sqrt{3}}{2}e_6$), [$e_5, \frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2, 0$], (8)($e_2, e_6, \frac{\sqrt{3}}{2}e_5 - \frac{1}{2}e_4$),
 $[e_6, -\frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2, 0]$ (9)($e_2, e_7, -\frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5$), [$e_7, -\frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2, 0$],
(10)($e_3, e_4, \frac{\sqrt{3}}{2}e_4 + \frac{1}{2}e_5$), [$e_4, \frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8, 0$], (11)($e_3, e_5, \frac{\sqrt{3}}{2}e_5 - \frac{1}{2}e_4$), [$e_5, \frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8, 0$],
(12)($e_3, e_6, -\frac{\sqrt{3}}{2}e_6 - \frac{1}{2}e_7$), [$e_6, \frac{1}{2}e_3 + \frac{\sqrt{3}}{2}e_8, 0$], (13)($e_3, e_7, -\frac{\sqrt{3}}{2}e_7 + \frac{1}{2}e_6$),
 $[e_7, \frac{1}{2}e_3 + \frac{\sqrt{3}}{2}e_8, 0]$, (14)($e_4, e_4, -\frac{1}{2}e_8 + \frac{\sqrt{3}}{2}e_3$), [$e_3, -\frac{1}{2}e_3 + \frac{\sqrt{3}}{2}e_8, 0$]

(15) $(e_4, e_5, \frac{\sqrt{3}}{2}e_8 + \frac{1}{2}e_3), [e_5, -e_4, 0]$, (16) $(e_4, e_6, \frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2), [e_6, \frac{\sqrt{3}}{2}e_5 + \frac{1}{2}e_4, 0]$,
(17) $(e_4, e_7, -\frac{\sqrt{3}}{2}e_2 + \frac{1}{2}e_1), [e_7, \frac{1}{2}e_4 + \frac{\sqrt{3}}{2}e_5, 0]$, (18) $(e_4, e_8, -\frac{\sqrt{3}}{2}e_5 - \frac{1}{2}e_4), [e_5, -e_3, 0]$,
(19) $(e_5, e_5, \frac{\sqrt{3}}{2}e_3 - \frac{1}{2}e_8), [e_3, \frac{\sqrt{3}}{2}e_8 - \frac{1}{2}e_3, 0]$, (20) $(e_5, e_6, \frac{\sqrt{3}}{2}e_2 - \frac{1}{2}e_1), [e_6, \frac{1}{2}e_5 - \frac{\sqrt{3}}{2}e_4, 0]$,
(21) $(e_5, e_7, \frac{\sqrt{3}}{2}e_1 + \frac{1}{2}e_2), [e_7, -\frac{\sqrt{3}}{2}e_4 + \frac{1}{2}e_5, 0]$,
(22) $(e_5, e_8, \frac{\sqrt{3}}{2}e_4 - \frac{1}{2}e_5), [e_4, e_3, 0]$, (23) $(e_6, e_6, -\frac{1}{2}e_8 - \frac{\sqrt{3}}{2}e_3), [e_3, -\frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8, 0]$,
(24) $(e_6, e_7, \frac{\sqrt{3}}{2}e_8 - \frac{1}{2}e_3), [e_7, -e_6, 0]$, (25) $(e_6, e_8, -\frac{\sqrt{3}}{2}e_7 - \frac{1}{2}e_6), [e_7, e_3, 0]$,
(26) $(e_7, e_7, \frac{1}{2}e_8 + \frac{\sqrt{3}}{2}e_3), [e_3, -\frac{1}{2}e_3 - \frac{\sqrt{3}}{2}e_8, 0]$, (27) $(e_7, e_8, -\frac{1}{2}e_7 + \frac{\sqrt{3}}{2}e_6), [e_6, -e_3, 0]$,
(28) $(-e_8, -e_8, -e_8), [e_1, -e_1, 0]$. where $e_8e_8 = -e_8$. (e_8 が单位元的な役割です).
Note that if we define a product by $[\frac{1}{2}e_j, \frac{1}{2}e_k] = \sum_{l=1}^8 f_{jkl}(\frac{1}{2}e_l)$, where the notation f_{jkl} refer ([G], [O], [K.3]), then it makes the Lie algebra of type A_2 .

Acknowledgment. This work was supported by the Research Institute for Mthmactical Science, an International Joint usage/Research Center located Kyoto Univeristy.

References

- [G] M.Gell-mann, Symmetry of Baryons and Mesons, Phys.Rev.125(1962), 1067-1084.
[K.1] N.Kamiya, On generalized structurable algebras and Lie related triple, Adv.Clifford Algebras,5(1995)127-140.
[K.2] N.Kamiya, A construction of Quandles associated with Quadratic algebras II, RIMS Kokyuroku(Kyoto Univ.) vol.2188,(2021) 34-44.
[K.3] N.Kamiya, On triality relations of certain eight dimensional algebras, RIMS, Kokyuroku (Kyoto Univ.) vol.2229, (2022) 52-63.
[K-O.1] N.Kamiya and S.Okubo, Triality of structurable and pre-structurable algebras, J.Alg.vol.416,(2014)58-83.
[K-O.2] N.Kamiya and S.Okubo, Algebras satisfying triality and S_4 symmetry, Arxiv.1503.00614, A.G.G.vol.33,(2016)1-92
[K-O.3] N.Kamiya and S.Okubo, A triality group of nonassociative algebras with involution, Arxiv.1609.05892, A.G.G.vol 35,(2018)113-168.
[K-O.4] N.Kamiya and S.Okubo, Symmetry of Lie algebras associated with (ε, δ) Freudenthal-Kantor triple systems. Proc.Edinb.Math.Soc. vol.89,(2016)169-192.
[O] S.Okubo, *Introduction to Octonion and other nonassociative algebras in Physics*, Cambridge Univ. press.(1995).
[S] R.D.Schafer, *An introduction to nonassociative algebras*, Academic press, (1966).
[T] J.Tits, Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles, I. construction, Indag. Math. 28(1966),223-237.
[Tô] S.Tôgô, リー代数, 槇書店,(1984).

Current address;

Noriaki Kamiya
shigekamiya@outlook.jp
Japan, Chigasaki city chigasaki 1-2-47-201