

## RIMS 共同研究「非圧縮性粘性流体の数理解析」

日時： 2020 年 12 月 7 日 (月) 14:00 ~ 12 月 9 日 (水) 12:00

研究代表者：前川 泰則 (京都大学)

副代表者：柴田 良弘 (早稲田大学)

本研究集会は Zoom によるオンライン開催となります。ご参加の方は以下から事前参加登録をお願い致します。

<https://forms.gle/xFp9NnLbEjUVfJq9>

参加登録締切日：12 月 4 日 (金)

Zoom のミーティング情報は、開催日が近づきましたら、参加登録をされた方にお知らせ致します。

### プログラム

#### 12 月 7 日 (月)

14:00 - 14:50 鈴木 政尋 (名古屋工業大学)

Stationary solutions to the Euler–Poisson equations in a perturbed half-space

15:10 - 16:00 小池 開 (京都大学)

Refined pointwise estimates for the solutions to the one-dimensional barotropic compressible Navier-Stokes equations: An application to the analysis of the long-time behavior of a moving point mass

16:20 - 16:50 石垣 祐輔 (東京工業大学)

Diffusion wave phenomena and  $L^p$  decay estimates of solutions of compressible viscoelastic system

12月8日(火)

10:00 - 10:50 谷内 靖 (信州大学)

On uniqueness of mild solutions on the whole time axis to the Boussinesq equations in unbounded domains

11:10 - 12:00 久保 隆徹 (お茶の水女子大学)

Analysis of non-stationary Navier-Stokes equations approximated by the pressure stabilization method

14:00 - 14:50 Xin Zhang (Tongji University, Shanghai)

The decay property of the multidimensional compressible flow in the exterior domain

15:10 - 16:00 相木 雅次 (東京理科大学)

On the head-on collision of coaxial vortex rings

16:20 - 16:50 清水 雄貴 (京都大学)

Current-valued solutions of the Euler-Arnold equation on surfaces and its applications

12月9日(水)

10:00 - 10:50 三浦 英之 (東京工業大学)

Estimates of the regular set for Navier-Stokes flows in terms of initial data

11:10 - 12:00 橋本 伊都子 (金沢大学 / OCAMI)

Existence of radially symmetric stationary solutions for the compressible Navier-Stokes equation



RIMS 共同研究  
「非圧縮性粘性流体の数理解析」

2020年12月7日—12月9日 (Online via Zoom)

研究代表者：前川泰則 (京都大学)  
副代表者：柴田良弘 (早稲田大学)

Abstract 集

# Stationary solutions to the Euler–Poisson equations in a perturbed half-space

Masahiro Suzuki

Department of Computer Science and Engineering, Nagoya Institute of Technology  
masahiro@nitech.ac.jp

The purpose of this talk is to mathematically investigate the formation of a plasma sheath near the surface of walls immersed in a plasma. The motion of plasma is governed by the Euler–Poisson equations:

$$\rho_t + \nabla \cdot (\rho \mathbf{u}) = 0, \quad \mathbf{u}_t + (\mathbf{u} \cdot \nabla) \mathbf{u} + K \nabla (\log \rho) = \nabla \phi, \quad \Delta \phi = \rho - e^{-\phi}, \quad (1a)$$

where unknown functions  $\rho$ ,  $\mathbf{u} = (u_1, u_2, u_3)$ , and  $-\phi$  represent the density and velocity of the positive ions and the electrostatic potential, respectively. Furthermore,  $K$  is a positive constant. We study an initial–boundary value problem of (1a) in a perturbed half-space  $\Omega := \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1 > M(x_2, x_3)\}$  with  $M \in \cap_{k=1}^{\infty} H^k(\mathbb{R}^2)$ . The initial and boundary data are prescribed as

$$(\rho, \mathbf{u})(0, x) = (\rho_0, \mathbf{u}_0)(x), \quad (1b)$$

$$\lim_{x_1 \rightarrow \infty} (\rho, u_1, u_2, u_3, \phi)(t, x_1, x_2, x_3) = (1, u_+, 0, 0, 0), \quad (1c)$$

$$\phi(t, M(x_2, x_3), x_2, x_3) = \phi_b \quad \text{for } (x_2, x_3) \in \mathbb{R}^2, \quad (1d)$$

where  $u_+ < 0$  and  $\phi_b \in \mathbb{R}$  are constants. The initial data  $(\rho_0, \mathbf{u}_0)$  are supposed to satisfy

$$\inf_{x \in \Omega} \rho_0(x) > 0, \quad \inf_{x \in \partial \Omega} \frac{\mathbf{u}_0(x) \cdot \nabla (M(x_2, x_3) - x_1)}{\sqrt{1 + |\nabla M(x_2, x_3)|^2}} - \sqrt{K} > 0. \quad (2)$$

For the end state  $u_+$ , we assume the Bohm criterion and the supersonic outflow condition:

$$u_+^2 > K + 1, \quad u_+ < 0, \quad (3)$$

$$\inf_{x \in \partial \Omega} \frac{-u_+}{\sqrt{1 + |\nabla M(x_2, x_3)|^2}} - \sqrt{K} > 0. \quad (4)$$

The second condition in (2) is necessary for the well-posedness of the problem (1). We remark that (4) is required if solutions to problem (1) are established in a neighborhood of the constant state  $(\rho, u_1, u_2, u_3, \phi) = (1, u_+, 0, 0, 0)$ .

In the case of planar wall  $M = 0$ , Bohm proposed a criterion on the velocity of the positive ion for the formation of sheath [1], and several mathematical results validated the Bohm criterion (3) and defined the fact that the sheath corresponds to the stationary solution of (1a). It is of greater interest to analyze the criterion for nonplanar walls. In this talk, we study the existence and stability of stationary solutions of (1) for  $M \neq 0$ .

To state our main results, let us introduce the existence theorem of stationary solutions  $(\tilde{\rho}, \tilde{u}, \tilde{\phi})(x_1)$  in a one-dimensional half-space. The stationary solutions solve

$$(\tilde{\rho} \tilde{u})' = 0, \quad \tilde{u} \tilde{u}' + K (\log \tilde{\rho})' = \tilde{\phi}', \quad \tilde{\phi}'' = \tilde{\rho} - e^{-\tilde{\phi}}, \quad x_1 > 0, \quad (5a)$$

$$\tilde{\phi}(0) = \phi_b, \quad \lim_{x_1 \rightarrow \infty} (\tilde{\rho}, \tilde{u}, \tilde{\phi})(x_1) = (1, u_+, 0), \quad \inf_{x_1 \in \mathbb{R}_+} \tilde{\rho}(x_1) > 0. \quad (5b)$$

**Theorem 1** ([2]). *Let  $u_+$  satisfy (3). There exists a constant  $\delta > 0$  such that if  $|\phi_b| < \delta$ , a unique monotone solution  $(\tilde{\rho}, \tilde{u}, \tilde{\phi}) \in C^\infty(\overline{\mathbb{R}_+})$  to (5) exists.*

We have constructed stationary solutions  $(\rho^s, \mathbf{u}^s, \phi^s)$  in the domain  $\Omega$  by regarding it as a perturbation of  $(\tilde{\rho}, \tilde{u}, 0, 0, \tilde{\phi})(\tilde{M}(x))$ , where  $\tilde{M}(x) := x_1 - M(x_2, x_3)$ . Furthermore, we use the weighted Sobolev space  $H_\alpha^k(\Omega)$  for  $k = 1, 2, 3, \dots$  and  $\alpha > 0$ :

$$H_\alpha^k(\Omega) := \left\{ f \in H^k(\Omega) \mid \|f\|_{k,\alpha}^2 < \infty \right\}, \quad \|f\|_{k,\alpha}^2 := \sum_{j=0}^k \int_{\Omega} e^{\alpha x_1} |\nabla^j f|^2 dx.$$

The existence and stability of stationary solutions are summarized in the following theorems. It is worth pointing out that we do not require any smallness assumptions for the function  $M$  representing the boundary  $\partial\Omega$ .

**Theorem 2** ([3]). *Let  $m \geq 3$ , and  $u_+$  satisfy (3) and (4). There exists a positive constant  $\delta$  such that if  $\beta + |\phi_b| \leq \delta$ , the problem (1) has a unique stationary solution  $(\rho^s, \mathbf{u}^s, \phi^s)$  as*

$$\begin{aligned} & (\rho^s, u_1^s, u_2^s, u_3^s, \phi^s) - (\tilde{\rho} \circ \tilde{M}, \tilde{u} \circ \tilde{M}, 0, 0, \tilde{\phi} \circ \tilde{M}) \in [H_\beta^m(\Omega)]^4 \times H_\beta^{m+1}(\Omega), \\ & \|(\rho^s - \tilde{\rho} \circ \tilde{M}, u_1^s - \tilde{u} \circ \tilde{M}, u_2^s, u_3^s)\|_{m,\beta}^2 + \|\phi^s - \tilde{\phi} \circ \tilde{M}\|_{m+1,\beta}^2 \leq C|\phi_b|, \end{aligned}$$

where  $C$  is a positive constant independent of  $\phi_b$ .

**Theorem 3** ([3]). *Let  $u_+$  satisfy (3) and (4). There exists a positive constant  $\delta$  such that if  $\beta + \|(\rho_0 - \rho^s, \mathbf{u}_0 - \mathbf{u}^s)\|_{3,\beta} + |\phi_b| \leq \delta$  the problem (1) has a unique time-global solution  $(\rho, \mathbf{u}, \phi)$  in the following space:*

$$(\rho - \rho^s, \mathbf{u} - \mathbf{u}^s, \phi - \phi^s) \in \left[ \bigcap_{i=0}^1 C^i([0, \infty); H_\beta^{3-i}(\Omega)) \right]^4 \times C([0, \infty); H_\beta^4(\Omega)).$$

Moreover, there holds for certain positive constants  $C$  and  $\gamma$  independent of  $\phi_b$  and  $t$ ,

$$\sup_{x \in \Omega} |(\rho - \rho^s, \mathbf{u} - \mathbf{u}^s, \phi - \phi^s)(t, x)| \leq C e^{-\gamma t}, \quad t \in [0, \infty).$$

We can conclude from Theorems 2 and 3 that (3) and (4) guarantee the sheath formation as long as the shape of walls is drawn by a graph.

**Acknowledgments.** This talk is based on a joint work with Prof. Masahiro Takayama (Keio Univ.).

## Reference

- [1] D. Bohm, *Minimum ionic kinetic energy for a stable sheath*, The characteristics of electrical discharges in magnetic fields, A. Guthrie and R.K. Wakerling eds., McGraw-Hill, New York, (1949), 77–86.
- [2] M. Suzuki, *Asymptotic stability of stationary solutions to the Euler–Poisson equations arising in plasma physics*, *Kinetic and Related Models* **4** (2010), pp569–588.
- [3] M. Suzuki and M. Takayama, *Stability and existence of stationary solutions to the Euler–Poisson equations in a domain with a curved boundary*, to appear in *Arch. Ration. Mech. Anal.*

# Refined pointwise estimates for the solutions to the one-dimensional barotropic compressible Navier–Stokes equations: An application to the analysis of the long-time behavior of a moving point mass

Kai Koike

Graduate School of Engineering, Kyoto University<sup>1</sup>  
koike.kai.42r@st.kyoto-u.ac.jp

The objective of this study is to understand the long-time behavior of a point mass moving inside a one-dimensional viscous compressible fluid. In a previous work [2], we showed that the velocity of the point mass  $V(t)$  satisfies a decay estimate  $V(t) = O(t^{-3/2})$ ; in this work, we give a necessary and sufficient condition (under some regularity and smallness assumptions) for the corresponding lower bound  $C^{-1}(t+1)^{-3/2} \leq |V(t)|$  to hold for large enough  $C$  and  $t$ .

The system of equations we consider is, in the Lagrangian mass coordinate, the following:

$$\begin{cases} v_t - u_x = 0, & x \in \mathbb{R}_*, t > 0, \\ u_t + p(v)_x = \nu \left( \frac{u_x}{v} \right)_x, & x \in \mathbb{R}_*, t > 0, \\ u(0_{\pm}, t) = V(t), & t > 0, \\ mV'(t) = \llbracket -p(v) + \nu u_x/v \rrbracket(t), & t > 0, \\ V(0) = V_0; v(x, 0) = v_0(x), u(x, 0) = u_0(x), & x \in \mathbb{R}_*. \end{cases} \quad (1)$$

Here,  $v = v(x, t)$  is the specific volume,  $u = u(x, t)$  is the velocity,  $p(v)$  is the pressure, and the constant  $\nu > 0$  is the viscosity of the fluid. We assume that the fluid is barotropic, so that  $p$  is a known function; we assume that  $p$  is smooth and satisfies  $p'(v) < 0$  and  $p''(v) \neq 0$  for  $v > 0$ . In the Lagrangian mass coordinate, the location of the point mass — whose mass and velocity are denoted by  $m$  and  $V(t)$  — is always  $x = 0$ , and  $\mathbb{R}_* := \mathbb{R} \setminus \{0\}$  is the domain in which the fluid flows. The double brackets  $\llbracket f \rrbracket(t)$  denote the jump of a function  $f = f(x, t)$  at  $x = 0$ , that is,  $\llbracket f \rrbracket(t) := f(0_+, t) - f(0_-, t)$ , where  $f(0_{\pm}, t) = \lim_{x \rightarrow \pm 0} f(x, t)$ . The first two equations in (1) are the barotropic compressible Navier–Stokes equations written in the Lagrangian mass coordinate, the so-called  $p$ -system; the third one is the Dirichlet boundary condition for the first two equations, which just says that the fluid does not penetrate through the point mass; the fourth equation is Newton’s second law; the final set of equations are initial conditions. In what follows, we set  $m = 1$  for simplicity.

We consider small solutions around a steady state  $(v, u, V) = (1, 0, 0)$ . We denote by  $\|\cdot\|_k$  the  $H^k(\mathbb{R}_*)$ -norm. The following theorem is obtained as a corollary to a theorem on the pointwise estimates of the fluid variables [1, Theorem 1.2]; due to the page limitation, we only state a corollary, which reads as follows [1, Corollary 1.2].<sup>2</sup>

**Theorem 1.** *Let  $v_0 - 1, u_0 \in H^6(\mathbb{R}_*)$ , and  $V_0 \in \mathbb{R}$ . Assume that they satisfy suitable compatibility conditions. Then there exist  $\delta_0 > 0$  and  $C > 1$  such that if*

$$\delta := \|v_0 - 1\|_6 + \|u_0\|_6 + \sup_{x \in \mathbb{R}^*} \left[ (|x| + 1)^{9/4} \{ |(v_0 - 1)(x)| + |u_0(x)| \} \right] \leq \delta_0 \quad (2)$$

<sup>1</sup>This work was supported by Grant-in-Aid for JSPS Research Fellow (Grant Number 20J00882).

<sup>2</sup>The assumption on the spatial decay can be slightly weakened as is stated in [1].

and

$$\left[ \int_{-\infty}^{\infty} (v_0 - 1)(x) dx \right] \cdot \left[ \int_{-\infty}^{\infty} u_0(x) dx + V_0 \right] \neq 0, \quad (3)$$

then the unique global-in-time solution  $(v, u, V)$  to (1) exists and satisfies

$$C^{-1} \delta^2 (t + 1)^{-3/2} \leq |V(t)| \quad (t \geq T(\delta)) \quad (4)$$

for some  $T(\delta) > 0$

**Remark 1.** (i) The upper bound  $|V(t)| = O(t^{-3/2})$  was obtained with less stringent assumptions [2], but we need to make these stronger to prove the lower bound.

(ii) By (4), we see that  $V(t)$  does not change its sign after sufficiently long time has elapsed. We can also predict the final sign of  $V(t)$  in terms of the initial data: it is the opposite sign of the left-hand side of (3).

(iii) When the left-hand side of (3) is zero, an improved decay estimate  $V(t) = O(t^{-7/4})$  can be proved [1, Corollary 1.3]. From some numerical simulations for the corresponding Cauchy problem, we conjecture that the rate  $-7/4$  is optimal under the condition that the left-hand side of (3) is zero, but to prove this would require much more work.

(iv) The presence of the point mass does not introduce additional technical difficulties compared to [2]; the main technical advancement lies in the analysis of the corresponding Cauchy problem.

The idea of the proof is to improve the previously known pointwise estimates for the fluid variables, see [3, Theorem 2.6] for the Cauchy problem and [2, Theorem 1.2] for our system, by a refined choice of leading order terms of the solution; we also need to analyze nonlinear interactions more precisely than in the previous works. We also note that we need to make use of finer space-time structure of fundamental solution, obtained in [4], compared to those presented in [3]. These allow us to understand the behavior of the solution around the origin  $x = 0$  more precisely and lead us to prove lower bound (4) for  $V(t) = u(0_{\pm}, t)$ .

## References

- [1] K. Koike, *Refined pointwise estimates for the solutions to the one-dimensional barotropic compressible Navier–Stokes equations: An application to the analysis of the long-time behavior of a moving point mass*, arXiv:2010.06578 (2020).
- [2] ———, *Long-time behavior of a point mass in a one-dimensional viscous compressible fluid and pointwise estimates of solutions*, J. Differential Equations **271** (2021), 356–413.
- [3] T.-P. Liu and Y. Zeng, *Large time behavior of solutions for general quasilinear hyperbolic-parabolic systems of conservation laws*, Mem. Amer. Math. Soc. **125** (1997), no. 599.
- [4] ———, *On Green’s function for hyperbolic-parabolic systems*, Acta Math. Sci. Ser. B (Engl. Ed.) **29B** (2009), 1556–1572.

# Diffusion wave phenomena and $L^p$ decay estimates of solutions of compressible viscoelastic system<sup>1</sup>

Yusuke Ishigaki

Department of Mathematics, Institute of Tokyo Technology

e-mail: ishigaki.y.aa@m.titech.ac.jp

## 1 Introduction

This talk is concerned with the following compressible viscoelastic system in  $\mathbb{R}^3$ :

$$(1.1) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho v) = 0, \\ \rho(\partial_t v + v \cdot \nabla v) - \nu \Delta v - (\nu + \nu') \nabla \operatorname{div} v + \nabla P(\rho) = \beta^2 \operatorname{div}(\rho F^\top F), \\ \partial_t F + v \cdot \nabla F = \nabla v F, \\ \operatorname{div}(\rho^\top F) = 0, \\ (\rho, v, F)|_{t=0} = (\rho_0, v_0, F_0), \quad \operatorname{div}(\rho_0^\top F_0) = 0. \end{cases}$$

Here  $\rho = \rho(x, t)$ ,  $v = {}^\top(v^1(x, t), v^2(x, t), v^3(x, t))$  and  $F = (F^{jk}(x, t))_{1 \leq j, k \leq 3}$  denote the unknown density, velocity field and deformation tensor, respectively, at time  $t \geq 0$  and position  $x \in \mathbb{R}^3$ .  $P(\rho)$  is the pressure that is a smooth function of  $\rho$  satisfying  $P'(1) > 0$ .  $\nu$ ,  $\nu'$  and  $\beta$  are constants satisfying  $\nu > 0$ ,  $2\nu + 3\nu' \geq 0$ ,  $\beta > 0$ . Here  $\nu$  and  $\nu'$  are the viscosity coefficients;  $\beta$  is the strength of elasticity. If we set  $\beta = 0$  formally, we obtain the compressible Navier-Stokes equations. Here and in what follows  ${}^\top \cdot$  stands for the transposition.

The aim of this talk is to investigate the large time behavior of solutions of the problem (1.1) around a motionless state  $(\rho, v, F) = (1, 0, I)$ . Here  $I$  is the  $3 \times 3$  identity matrix.

In the case  $\beta = 0$ , Hoff-Zumbrun[1] derived the following  $L^p$  ( $1 \leq p \leq \infty$ ) decay estimates and asymptotic properties:

$$\|(\phi(t), m(t))\|_{L^p} \leq \begin{cases} C(1+t)^{-\frac{3}{2}(1-\frac{1}{p})-\frac{1}{2}(1-\frac{2}{p})}, & 1 \leq p < 2, \\ C(1+t)^{-\frac{3}{2}(1-\frac{1}{p})}, & 2 \leq p \leq \infty. \end{cases}$$

$$\left\| \left( (\phi(t), m(t)) - \left( 0, \mathcal{F}^{-1} \left( e^{-\nu|\xi|^2 t} \hat{\mathcal{P}}(\xi) \hat{m}_0 \right) \right) \right) \right\|_{L^p} \leq C(1+t)^{-\frac{3}{2}(1-\frac{1}{p})-\frac{1}{2}(1-\frac{2}{p})}, \quad 2 \leq p \leq \infty,$$

where  $(\phi(t), m(t)) = (\rho(t) - 1, \rho(t)v(t))$  and  $\hat{\mathcal{P}}(\xi) = I - \frac{\xi^\top \xi}{|\xi|^2}$ ,  $\xi \in \mathbb{R}^3$ . The authors of [1] showed that the hyperbolic aspect of sound wave plays a role of the spreading effect of the wave equation, and the decay rate of the solution becomes slower than the heat kernel when  $1 \leq p < 2$ . On the other hand, if  $2 < p \leq \infty$ , the compressible part of the solution  $(\phi(t), m(t)) - \left( 0, \mathcal{F}^{-1} \left( e^{-\nu|\xi|^2 t} \hat{\mathcal{P}}(\xi) \hat{m}_0 \right) \right)$  converges to 0 faster than the heat kernel.

In the case  $\beta > 0$ , Hu-Wu[2] and Li-Wei-Wao[5] established the following  $L^p$  ( $2 \leq p \leq \infty$ ) decay estimates:

$$\|u(t)\|_{L^p} \leq C(1+t)^{-\frac{3}{2}(1-\frac{1}{p})},$$

where  $u(t) = (\phi(t), w(t), G(t)) = (\rho(t), w(t), F(t)) - (1, 0, I)$ . However the hyperbolic aspects of elastic shear wave and sound wave does not appear. We will clarify the diffusion wave phenomena caused by interaction of three properties; sound wave, viscous diffusion and elastic shear wave and improve the results obtained in [2, 5].

<sup>1</sup>This work was partially supported by JSPS KAKENHI Grant Number 19J10056.

We consider the nonlinear problem for  $u(t) = (\phi(t), w(t), G(t))$ :

$$(1.2) \quad \begin{cases} \partial_t \phi + \operatorname{div} w = g_1, \\ \partial_t w - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w + \gamma^2 \nabla \phi - \beta^2 \operatorname{div} G = g_2, \\ \partial_t G - \nabla w = g_3, \\ \nabla \phi + \operatorname{div}^\top G = g_4, \\ u|_{t=0} = u_0 = (\phi_0, w_0, G_0). \end{cases}$$

Here  $\tilde{\nu} = \nu + \nu'$ ;  $g_j$ ,  $j = 1, 2, 3, 4$  are nonlinear terms.

## 2 Main Result

We have the following result.

**Theorem 2.1.** ([3]) *Let  $1 < p \leq \infty$ . Assume that  $\phi_0$ ,  $G_0$ , and  $F_0^{-1}$  satisfy  $\nabla \phi_0 - \operatorname{div}^\top(I + G_0)^{-1} = 0$  and  $F_0^{-1} = \nabla X_0$  for some vector field  $X_0$ . If  $u_0 = (\phi_0, w_0, G_0)$  satisfies  $\|u_0\|_{H^3} \ll 1$  and  $u_0 \in L^1$ , then there exists a unique solution  $u(t) \in C([0, \infty); H^3)$  of the problem (1.2), and  $u(t) = (\phi(t), w(t), G(t))$  satisfies*

$$\|u(t)\|_{L^p} \leq C(1+t)^{-\frac{3}{2}(1-\frac{1}{p})-\frac{1}{2}(1-\frac{2}{p})} (\|u_0\|_{L^1} + \|u_0\|_{H^3}), \quad t \geq 0.$$

Here  $C(p)$  is a positive constant depending only on  $p$ .

**Outline of the proof.** We note that the solenoidal part of the solution of linearized system  $(w_s, \tilde{G}_s) = (\mathcal{F}^{-1}(\hat{\mathcal{P}}(\xi)\hat{w}), \beta\mathcal{F}^{-1}(\hat{\mathcal{P}}(\xi)\hat{G}))$  satisfies the following linear symmetric parabolic-hyperbolic system:

$$\begin{cases} \partial_t w_s - \nu \Delta w_s - \beta \operatorname{div} \tilde{G}_s = 0, \\ \partial_t \tilde{G}_s - \beta \nabla w_s = 0. \end{cases}$$

It follows from [4, 6] that if  $p > 2$ , then the  $L^p$  norm of the solution of the linearized problem decays faster than the case  $\beta = 0$ . In the case of the nonlinear problem, we use a nonlinear transform  $\tilde{\psi}$  defined by  $\psi = \tilde{\psi} - (-\Delta)^{-1} \operatorname{div}^\top(\phi \nabla \tilde{\psi} + (1 + \phi)h(\nabla \tilde{\psi}))$  instead of  $G$ . Here  $\tilde{\psi}$  is a displacement vector and  $(-\Delta)^{-1} = \mathcal{F}^{-1}|\xi|^{-2}\mathcal{F}$ . We then see that the nonlinear constraint  $\operatorname{div}(\rho^\top F) = 0$  becomes the linear condition  $\phi + \operatorname{tr}(\nabla \psi) = \phi + \operatorname{div} \psi = 0$  and straightforward application of the semigroup theory works well. Here  $h(\nabla \tilde{\psi})$  is a nonlinear term satisfying  $h(\nabla \tilde{\psi}) = O(|\nabla \tilde{\psi}|^2)$ ,  $|\nabla \tilde{\psi}| \ll 1$ .

## References

- [1] D. Hoff and K. Zumbrun: Multi-dimensional diffusion waves for the Navier-Stokes equations of compressible flow, *Indiana Univ. Math. J.* **44** (1995), pp. 603-676.
- [2] X. Hu, G. Wu, Global existence and optimal decay rates for three-dimensional compressible viscoelastic flows, *SIAM J. Math. Anal.*, **45** (2013), pp. 2815-2833.
- [3] Y. Ishigaki, Diffusion wave phenomena and  $L^p$  decay estimates of solutions of compressible viscoelastic system, *J. Differential Equations*, **269** (2020), pp. 11195-11230.
- [4] T. Kobayashi, Y. Shibata, Remark on the rate of decay of solutions to linearized compressible Navier-Stokes equation, *Pacific J. Math.* **207** (2002), pp. 199-234.
- [5] Y. Li, R. Wei, Z. Yao, Optimal decay rates for the compressible viscoelastic flows, *J. Math. Phys.*, **57**, 111506, (2016).
- [6] Y. Shibata, On the rate of decay of solutions to linear viscoelastic equation, *Math. Methods Appl. Sci.*, **23** (2000), pp. 203-226.

# On uniqueness of mild solutions on the whole time axis to the Boussinesq equations in unbounded domains

Yasushi TANIUCHI

Shinshu University  
taniuchi@math.shinshu-u.ac.jp

In this talk, we consider the Boussinesq equations in 3-dimensional unbounded domains  $\Omega$ . The Boussinesq equations describe the heat convection in a viscous incompressible fluid.

$$(B) \begin{cases} \partial_t u - \Delta u + u \cdot \nabla u + \nabla p = g\theta, & t \in \mathbb{R}, x \in \Omega, \\ \partial_t \theta - \Delta \theta + u \cdot \nabla \theta = S, & t \in \mathbb{R}, x \in \Omega, \\ \nabla \cdot u = 0, & t \in \mathbb{R}, x \in \Omega, \\ u|_{\partial\Omega} = 0, \quad \theta|_{\partial\Omega} = \xi, \end{cases}$$

where  $u = (u^1(x, t), u^2(x, t), u^3(x, t))$ ,  $\theta = \theta(x, t)$  and  $p = p(x, t)$  denote the velocity vector, the temperature and the pressure, respectively, of the fluid at the point  $(x, t) \in \Omega \times \mathbb{R}$ . Here  $\xi = \xi(x, t)$  is a given boundary temperature,  $S$  is a given external heat source and  $g$  denotes the acceleration of gravity. When  $\Omega$  is some unbounded domain (e.g. the half-space  $\mathbb{R}_+^3$ ), we can show the existence theorem of small mild solutions on whole time axis to (B). Typical examples of solutions on whole time axis are stationary, time-periodic and almost time-periodic solutions. In this talk, we consider the uniqueness of such solutions. Very roughly speaking, we will show that if there are two solutions  $(u, \theta_1)$  and  $(v, \theta_2)$  in some function spaces with the same data and if we assume that  $\theta_1$  is small in some sense, then  $(u, \theta_1) = (v, \theta_2)$ . Here, we do not need to assume any smallness condition on  $u, v, \theta_2$ .

# Analysis of non-stationary Navier-Stokes equations approximated by the pressure stabilization method

Takayuki Kubo

Ochanomizu Univ.

kubo.takayuki@ocha.ac.jp

The results of this talk are joint works with Dr. R. Matsui (Ushiku high school affiliated with Toyo University) and with Dr. H. Kikuchi (Univ. of Tsukuba).

The mathematical description of fluid flow is given by the Navier-Stokes equations:

$$\left\{ \begin{array}{ll} \partial_t u - \Delta u + (u \cdot \nabla)u + \nabla \pi = f & t > 0, x \in \Omega, \\ \nabla \cdot u = 0 & t > 0, x \in \Omega, \\ u(0, x) = a & x \in \Omega, \\ u(t, x) = 0 & x \in \partial\Omega, \end{array} \right. \quad (\text{NS})$$

where the fluid vector fields  $u = u(t, x)$  and the pressure  $\pi = \pi(t, x)$  are unknown function, the external force  $f = f(t, x)$  is a given vector function, the initial data  $a$  is a given solenoidal function and  $\Omega$  is a bounded domain with smooth boundary. It is well-known that one of the difficulty of analysis for Navier Stokes equations (NS) is the pressure term  $\nabla \pi$  and incompressible condition  $\nabla \cdot u = 0$ .

In order to overcome this difficulty, we often use Helmholtz decomposition. The Helmholtz decomposition means that for  $1 < p < \infty$ ,  $L^p(\Omega)^n = L_{p,\sigma}(\Omega) \oplus G_p(\Omega)$ , where  $L_{p,\sigma}(\Omega) = \overline{\{u \mid u_j \in C_0^\infty, \nabla \cdot u = 0\}}^{\|\cdot\|_{L^p}}$  and  $G_p(\Omega) = \{\nabla \pi \in L^p(\Omega)^n \mid \pi \in L_{p,\text{loc}}(\Omega)\}$ . On the other hand, in numerical analysis, some penalty methods are employed as the method to overcome this difficulty. They are methods that eliminate the pressure by using approximated incompressible condition. For example, setting  $\alpha > 0$  as a perturbation parameter, we use  $\nabla \cdot u = -\pi/\alpha$  in the penalty method,

$$(u, \nabla \varphi)_\Omega = \alpha^{-1}(\nabla \pi, \nabla \varphi)_\Omega \quad (\varphi \in \widehat{W}_q^1(\Omega)) \quad (\text{wi})$$

in the pressure stabilization method and  $\nabla \cdot u = -\partial_t \pi/\alpha$  in the pseudocompressible method.

In this talk, we consider (wi) instead of incompressible conditions  $\nabla \cdot u = 0$  in (NS). Namely we consider the following equations:

$$\left\{ \begin{array}{ll} \partial_t u_\alpha - \Delta u_\alpha + (u_\alpha \cdot \nabla)u_\alpha + \nabla \pi_\alpha = f & t > 0, x \in \Omega, \\ u_\alpha(0, x) = a_\alpha & x \in \Omega, \\ u_\alpha(t, x) = 0, \quad \partial_n \pi_\alpha(t, x) = 0 & x \in \partial\Omega. \end{array} \right. \quad (\text{NSa})$$

under the approximated weak incompressible condition (wi) in  $L^q$ -framework ( $n/2 < q < \infty$ ). We shall use the maximal regularity theorem for linearized problem for (NSa) in order to prove the local in time existence theorem and the error estimate in the  $L_p$  in time and the  $L_q$  in space framework with  $n/2 < q < \infty$  and  $\max\{1, n/q\} < p < \infty$ .

Main result in this talk is concerned with the local in time existence theorem for (NSa) with approximated weak incompressible condition (wi).

**Theorem 1.** Let  $n \geq 2$ ,  $n/2 < q < \infty$  and  $\max\{1, n/q\} < p < \infty$ . Let  $\alpha > 0$  and  $T_0 \in (0, \infty)$ . For any  $M > 0$ , assume that the initial data  $a_\alpha \in B_{q,p}^{2(1-1/p)}(\Omega) = (L_q(\Omega), W_q^2(\Omega))_{1-1/p,p}$  and the external force  $f \in L_p((0, T_0), L_q(\Omega)^n)$  satisfy

$$\|a_\alpha\|_{B_{q,p}^{2(1-1/p)}(\Omega)} + \|f\|_{L_p((0, T_0), L_q(\Omega)^n)} \leq M.$$

Then, there exists  $T^*$  depending on only  $M$  such that (NSa) under (wi) has a unique solution  $(u_\alpha, \pi_\alpha)$  of the following class:

$$\begin{aligned} u_\alpha &\in W_p^1((0, T^*), L_q(\Omega)^n) \cap L_p((0, T^*), W_q^2(\Omega)^n), \\ \pi_\alpha &\in L_p((0, T^*), \widehat{W}_q^1(\Omega)). \end{aligned}$$

Moreover the following estimate holds:

$$\|u_\alpha\|_{L_\infty((0, T^*), L_q(\Omega))} + \|(\partial_t u_\alpha, \nabla^2 u_\alpha, \nabla \pi_\alpha)\|_{L_p((0, T^*), L_q(\Omega))} + \|\nabla u_\alpha\|_{L_r((0, T^*), L_q(\Omega))} \leq C$$

for  $1/p - 1/r \leq 1/2$ , where  $C$  is the positive constant depend on  $n, p, q$  and  $T^*$ .

Next we consider the error estimate between the solution  $(u, \pi)$  to (NS) under the weak incompressible condition  $(u, \nabla \varphi)_\Omega = 0$  for  $\varphi \in \widehat{W}_q^1(\Omega)$  and solution  $(u_\alpha, \pi_\alpha)$  to (NSa) under (wi). To this end, setting  $u_e = u - u_\alpha$  and  $\pi_e = \pi - \pi_\alpha$ , we see that  $(u_e, \pi_e)$  enjoys that

$$\begin{cases} \partial_t u_e - \Delta u_e + \nabla \pi_e + N(u_e, u_\alpha) = 0, & t > 0, x \in \Omega, \\ u_e(0, x) = a_e, & x \in \Omega, \\ u_e(t, x) = 0, & x \in \partial\Omega, \end{cases} \quad (\text{PE})$$

where  $N(u_e, u_\alpha) = (u_e \cdot \nabla)u_e + (u_e \cdot \nabla)u_\alpha + (u_\alpha \cdot \nabla)u_e$  and  $a_e = a - a_\alpha$  under the approximated weak incompressible condition

$$(u_e, \nabla \varphi)_\Omega = \alpha^{-1}(\nabla \pi_e, \nabla \varphi)_\Omega + \alpha^{-1}(\nabla \pi, \nabla \varphi)_\Omega \quad \varphi \in \widehat{W}_q^1(\Omega) \quad (\text{wie})$$

for  $1 < q < \infty$ . In a similar way to Theorem 1, we obtain the following theorems:

**Theorem 2.** Let  $n \geq 2$ ,  $n/2 < q < \infty$ ,  $\max\{1, n/q\} < p < \infty$  and  $\alpha > 0$ . Let  $T^*$  be a positive constant obtained in Theorem 1 and  $(u_\alpha, \pi_\alpha)$  be a solution obtained in Theorem 1. For any  $M > 0$ , assume that  $a_e \in B_{q,p}^{2(1-1/p)}(\Omega)$  satisfies

$$\|a_e\|_{B_{q,p}^{2(1-1/p)}(\Omega)} \leq M\alpha^{-1}.$$

Then there exists  $T^\flat$  such that (PE) has a unique solution  $(u_e, \pi_e)$  which satisfies

$$\|u_e\|_{L_\infty((0, T^\flat), L_q(\Omega))} + \|\nabla u_e\|_{L_r((0, T^\flat), L_q(\Omega))} + \|(\nabla^2 u_e, \partial_t u_e, \nabla \pi_e)\|_{L_p((0, T^\flat), L_q(\Omega))} \leq C\alpha^{-1}$$

for  $1/p - 1/r \leq 1/2$ .

Furthermore, in this talk, we will introduce the estimates for error  $(u_e, \pi_e)$  derived from the maximal regularity theorem for the linearized problem for (NSa) under the approximated weak incompressible condition (wi).

# The decay property of the multidimensional compressible flow in the exterior domain

Xin Zhang

School of Mathematical Sciences, Tongji University  
xinzhang2020@tongji.edu.cn

This talk discusses about the decay property of the compressible flow in the exterior domain in the general  $L_p$  framework. Here, we consider the motion of the gases with some free surface  $\Gamma_t$ , which can be described by the following (barotropic) compressible Navier-Stokes system in the exterior domain  $\Omega_t \subset \mathbb{R}^N (N \geq 3)$  :

$$\begin{cases} \partial_t \rho + \operatorname{div}((\rho_e + \rho)\mathbf{v}) = 0 & \text{in } \Omega_t, \\ (\rho_e + \rho)(\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v}) - \operatorname{Div}(\mathbf{S}(\mathbf{v}) - P(\rho_e + \rho)\mathbf{I}) = 0 & \text{in } \Omega_t, \\ (\mathbf{S}(\mathbf{v}) - P(\rho_e + \rho)\mathbf{I})\mathbf{n}_{\Gamma_t} = P(\rho_e)\mathbf{n}_{\Gamma_t}, \quad V_{\Gamma_t} = \mathbf{v} \cdot \mathbf{n}_{\Gamma_t} & \text{on } \Gamma_t, \\ (\rho, \mathbf{v}, \Omega_t)|_{t=0} = (\rho_0, \mathbf{v}_0, \Omega) & \end{cases} \quad (1)$$

Given the initial data and the reference density  $\rho_e > 0$ , we seek for the velocity field  $\mathbf{v}$ , the mass density  $\rho + \rho_e$  and the pattern of  $\Omega_t$ . In (1), the Cauchy stress tensor

$$\mathbf{S}(\mathbf{v}) = \mu \mathbf{D}(\mathbf{v}) + (\nu - \mu) \operatorname{div} \mathbf{v} \mathbf{I} \text{ for constants } \mu, \nu > 0,$$

and the doubled deformation tensor  $\mathbf{D}(\mathbf{v}) = \nabla \mathbf{v} + (\nabla \mathbf{v})^\top$ . Moreover, the  $(i, j)$ th entry of the matrix  $\nabla \mathbf{v}$  is  $\partial_i v_j$ ,  $\mathbf{I}$  is the  $N \times N$  identity matrix, and  $\mathbf{M}^\top$  is the transposed the matrix  $\mathbf{M} = [M_{ij}]$ . In addition,  $\operatorname{Div} \mathbf{M}$  denotes an  $N$ -vector of functions whose  $i$ -th component is  $\sum_{j=1}^N \partial_j M_{ij}$ ,  $\operatorname{div} \mathbf{v} = \sum_{j=1}^N \partial_j v_j$ , and  $\mathbf{v} \cdot \nabla = \sum_{j=1}^N v_j \partial_j$  with  $\partial_j = \partial / \partial x_j$ . On the moving boundary  $\Gamma_t$  of  $\Omega_t$ ,  $\mathbf{n}_{\Gamma_t}$  is the outer unit normal vector to the boundary  $\Gamma_t$  of  $\Omega_t$ , and  $V_{\Gamma_t}$  stands for the normal velocity of the moving surface  $\Gamma_t$ .

To study (1), we transfer (1) to some system in the fixed (or initial) domain  $\Omega$ . Assume that  $\Omega \subset \mathbb{R}^N$  with the boundary  $\Gamma$  is an exterior domain such that  $\mathcal{O} = \mathbb{R}^N \setminus \Omega$  is a subset of the ball  $B_R$ , centred at origin with radius  $R > 1$ . Let  $\kappa$  be a  $C^\infty$  functions which equals to one for  $x \in B_R$  and vanishes outside of  $B_{2R}$ . Then we define the partial Lagrangian coordinates

$$x = X_{\mathbf{w}}(y, T) = y + \int_0^T \kappa(y) \mathbf{w}(y, s) ds \quad (\forall y \in \Omega), \quad (2)$$

for some vector field  $\mathbf{w} = \mathbf{w}(\cdot, s)$  defined in  $\Omega$ . By using (2), denoting  $(\gamma_1, \gamma_2) = (\rho_e, P'(\rho_e))$  and neglecting the nonlinear forms, we obtain the following linearized equations \*

$$\begin{cases} \partial_t \rho + \gamma_1 \operatorname{div} \mathbf{v} = 0 & \text{in } \Omega \times \mathbb{R}_+, \\ \gamma_1 \partial_t \mathbf{v} - \operatorname{Div}(\mathbf{S}(\mathbf{v}) - \gamma_2 \rho \mathbf{I}) = 0 & \text{in } \Omega \times \mathbb{R}_+, \\ (\mathbf{S}(\mathbf{v}) - \gamma_2 \rho \mathbf{I})\mathbf{n}_\Gamma = 0 & \text{on } \Gamma \times \mathbb{R}_+, \\ (\rho, \mathbf{v})|_{t=0} = (\rho_0, \mathbf{v}_0) & \text{in } \Omega. \end{cases} \quad (3)$$

---

\*In fact, our result can be extended to more general linear equations with variable coefficients.

Now, our main result of (3) reads as follows:

**Theorem 1** ( $L_p$ - $L_q$  type decay estimate). *Let  $\Omega$  be a  $C^3$  exterior domain in  $\mathbb{R}^N$  with  $N \geq 3$ . Assume that  $(\rho_0, \mathbf{v}_0) \in L_q(\Omega)^{1+N} \cap H_p^{1,0}(\Omega)$  with  $H_p^{1,0}(\Omega) = H_p^1(\Omega) \times L_p(\Omega)^N$  for  $1 \leq q \leq 2 \leq p < \infty$ , and  $\{T(t)\}_{t \geq 0}$  is the semigroup associated to (3) in  $H_p^{1,0}(\Omega)$ . For convenience, we set  $\mathcal{P}_v(\rho, \mathbf{v}) = \mathbf{v}$  and*

$$\|(\rho_0, \mathbf{v}_0)\|_{p,q} = \|(\rho_0, \mathbf{v}_0)\|_{L_q(\Omega)} + \|(\rho_0, \mathbf{v}_0)\|_{H_p^{1,0}(\Omega)}.$$

Then for  $t \geq 1$ , there exists a positive constant  $C$  such that

$$\begin{aligned} \|T(t)(\rho_0, \mathbf{v}_0)\|_{L_p(\Omega)} &\leq Ct^{-(N/q - N/p)/2} \|(\rho_0, \mathbf{v}_0)\|_{p,q}, \\ \|\nabla T(t)(\rho_0, \mathbf{v}_0)\|_{L_p(\Omega)} &\leq Ct^{-\sigma_1(p,q,N)} \|(\rho_0, \mathbf{v}_0)\|_{p,q}, \\ \|\nabla^2 \mathcal{P}_v T(t)(\rho_0, \mathbf{v}_0)\|_{L_p(\Omega)} &\leq Ct^{-\sigma_2(p,q,N)} \|(\rho_0, \mathbf{v}_0)\|_{p,q}, \end{aligned}$$

where the indices  $\sigma_1(p, q, N)$  and  $\sigma_2(p, q, N)$  are given by

$$\begin{aligned} \sigma_1(p, q, N) &= \begin{cases} (N/q - N/p)/2 + 1/2 & \text{for } 2 \leq p \leq N, \\ N/(2q) & \text{for } N < p < \infty, \end{cases} \\ \sigma_2(p, q, N) &= \begin{cases} 3/(2q) & \text{for } N = 3, \\ (N/q - N/p)/2 + 1 & \text{for } N \geq 4 \text{ and } 2 \leq p \leq N/2, \\ N/(2q) & \text{for } N \geq 4 \text{ and } N/2 < p < \infty. \end{cases} \end{aligned}$$

The proof of Theorem 1 relies on the spectral analysis and the local energy method. This is a joint work with Yoshihiro Shibata from Waseda University.

# On the Head-on Collision of Coaxial Vortex Rings

Masashi Aiki

Department of Mathematics, Faculty of Science and Technology,  
Tokyo University of Science, 2641 Yamazaki, Noda, Chiba 278-8510, Japan

## 1 Introduction

The study of the interaction of coaxial vortex rings dates back to the pioneering paper by Helmholtz [2]. In [2], Helmholtz considered vortex motion in an incompressible and inviscid fluid based on the Euler equations. His study includes the motion of circular vortex filaments, and he observed that motion patterns such as head-on collision may occur. Since then, many researches have been done on head-on collision of coaxial vortex rings, and interaction of coaxial vortex rings in general. Most of these researches are either experiments conducted in a laboratory or numerical simulations of the Navier–Stokes equations, and the rigorous mathematical treatment of head-on collision of vortex rings are very scarce.

In light of this, we consider the head-on collision of two coaxial vortex rings, which have circulations of opposite sign, described as the motion of two coaxial circular vortex filaments under the localized induction approximation. A vortex filament is a space curve on which the vorticity of the fluid is concentrated. In our present work, we approximated thin vortex structures, such as vortex rings, by vortex filaments, and described the motion as the motion of a curve in the three-dimensional Euclidean space. In this formulation, a vortex ring is a space curve in the shape of a circle. We prove the existence of solutions to a system of nonlinear partial differential equations modelling the interaction of two vortex filaments proposed by the speaker [1] which exhibit head-on collision. We also give a necessary and sufficient condition for the initial configuration and parameters of the filaments for head-on collision to occur. Our results suggest that there exists a critical value  $\gamma_* > 1$  for the ratio  $\gamma$  of the magnitude of the circulations satisfying the following. When  $\gamma \in [1, \gamma_*]$ , two approaching rings will collide, and when  $\gamma \in (\gamma_*, \infty)$ , the ring with the larger circulation passes through the other and then separate indefinitely. As far as the speaker knows, the existence of such threshold  $\gamma_*$  is only indirectly suggested via numerical investigations of the head-on collision of coaxial vortex rings, for example by Inoue, Hattori, and Sasaki [3]. Hence, our result is the first to obtain the threshold in a way that is possible to numerically calculate  $\gamma_*$ , as well as prove that the threshold exists in a framework of a mathematical model.

## 2 Problem Setting and Main Results

We consider the motion of two interacting vortex filaments. In [1], under physically intuitive assumptions, we derived the following system of partial differential equations which describes the interaction of two vortex filaments.

$$(1) \quad \begin{cases} \mathbf{X}_t = \beta \frac{\mathbf{X}_\xi \times \mathbf{X}_{\xi\xi}}{|\mathbf{X}_\xi|^3} - \alpha \frac{\mathbf{Y}_\xi \times (\mathbf{X} - \mathbf{Y})}{|\mathbf{X} - \mathbf{Y}|^3}, \\ \mathbf{Y}_t = \frac{\mathbf{Y}_\xi \times \mathbf{Y}_{\xi\xi}}{|\mathbf{Y}_\xi|^3} - \alpha\beta \frac{\mathbf{X}_\xi \times (\mathbf{Y} - \mathbf{X})}{|\mathbf{X} - \mathbf{Y}|^3}, \end{cases}$$

where  $\mathbf{X}(\xi, t) = {}^t(X_1(\xi, t), X_2(\xi, t), X_3(\xi, t))$  and  $\mathbf{Y}(\xi, t) = {}^t(Y_1(\xi, t), Y_2(\xi, t), Y_3(\xi, t))$  are the position vectors of the vortex filaments parametrized by  $\xi$  at time  $t$ ,  $\times$  is the exterior product in the three-dimensional Euclidean space, subscripts denote partial differentiation with the respective variables,  $\beta \in \mathbf{R} \setminus \{0\}$  is the quotient of the vorticity strengths of the filaments, and  $\alpha > 0$  is a constant which is introduced in the course of the derivation of the system. We consider the case  $\beta < 0$  to describe colliding vortex rings. To make things more simple, we set  $\gamma = -\beta$  and consider  $\gamma > 0$ .

In this talk, we introduce recent results on the existence of solutions to system (1) which correspond to head-on collision. We give necessary and sufficient conditions on the initial data and parameters for head-on collision to occur. This, in particular, gives the threshold for  $\gamma$  mentioned in the introduction.

## References

- [1] M. Aiki, On the existence of leapfrogging pair of circular vortex filaments, *Stud. Appl. Math.*, **143** (2019), no.3, pp.213–243.
- [2] H. HELMHOLTZ, Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen, *J. Reine Angew. Math.* **55**:25–55 (1858).
- [3] O. Inoue, Y. Hattori, and T. Sasaki, Sound generation by coaxial collision of two vortex rings, *J. Fluid Mech.* **424** (2000), pp.327–365.
- [4] T. T. Lim and T. B. Nickels, Instability and reconnection in the head-on collision of two vortex rings, *Nature* **357** (1992), pp.225–227.
- [5] Y. Oshima, Head-on Collision of Two Vortex Rings, *J. Phys. Soc. Jpn.* **44** (1978), no.1, pp.328–331.

# Current-valued solutions of the Euler-Arnold equation on surfaces and its applications

Yuuki Shimizu

Kyoto university

shimizu@math.kyoto-u.ac.jp

The motion of incompressible and inviscid fluids in the Euclidean plane is governed by the Euler equation and its solution is called an Euler flow. For the Euler flow, the vorticity is a Lagrange invariance. Then, since the fluid velocity and the pressure can be recovered from the vorticity, an Euler flow is determined by a solution of the vorticity equation. Namely, if  $\omega_t$  is a solution of the vorticity equation,

$$\partial_t \omega_t + (-\mathcal{J} \text{grad}\langle G, \omega_t \rangle \cdot \nabla) \omega_t = 0,$$

$(v_t, p_t)$  is an Euler flow, defined by

$$v_t = -\mathcal{J} \text{grad}\langle G, \omega_t \rangle, \quad p_t = \langle G, \text{div}(v_t \cdot \nabla)v_t \rangle, \quad (1)$$

where  $\mathcal{J}$  is the symplectic matrix and  $G$  is the Green function for the Laplacian. On the other hand, the formulae (1) still make sense in the sense of distributions when we give a time-dependent distribution  $\Omega_t$  by a linear combination of delta functions centered at  $q_n(t)$  for  $n = 1, \dots, N$  with the linear coefficient  $\Gamma_n \in \mathbb{R}$ . Then, replacing  $\omega_t$  by  $\Omega_t$  in (1), we formally obtain a fluid velocity  $V_t$  and a pressure  $P_t$ . However, we can not define the dynamics of  $q_n(t)$  from the vorticity equation. Instead, to determine the evolution of  $q_n(t)$  by  $V_t$ , Helmholtz considered the following regularized equation for  $q_n(t)$  [6].

$$\dot{q}_n = \lim_{q \rightarrow q_n} [V_t(q) + \mathcal{J} \text{grad}\langle G, \Gamma_n \delta_{q_n(t)} \rangle(q)] = -\mathcal{J} \text{grad} \sum_{\substack{m=1 \\ m \neq n}}^N \Gamma_m G(q_n, q_m) \equiv v_n(q_n). \quad (2)$$

It is called the *point vortex equation*, and the solution of (2) is called the *point vortex dynamics*. Then, there arises a natural question; How can we interpret  $(V_t, P_t)$  as an Euler flow in an appropriate mathematical sense? In other words, we need to determine a space of solutions of the Euler equation which contains  $(V_t, P_t)$ . Since  $L^p$  space does not contain  $(V_t, P_t)$ , a more sophisticated space is to be considered. This is one of the central problems in the analysis of 2D Euler equation as discussed in [2, 3, 4]. From the viewpoint of the application, the point vortex dynamics is sometimes considered in the presence of a time-dependent vector field  $X_t \in \mathfrak{X}^r(\mathbb{R}^2)$ , called the *point vortex dynamics in the background field  $X_t$* . Then, the evolution of  $q_n(t)$  is governed by the following equation.

$$\dot{q}_n(t) = \beta_X X_t(q_n(t)) + \beta_\omega v_n(q_n(t)), \quad n = 1, \dots, N \quad (3)$$

for a given  $(\beta_X, \beta_\omega) \in \mathbb{R}^2$ . Some experimental studies confirm the importance of background fields in two-dimensional turbulence [1, 7].

The purpose of this research is justifying the point vortex dynamics in background fields as an Euler flow mathematically. To this end, we establish a weak formulation of the Euler equation in the space of currents, which is developed in the theory of geometric analysis and geometric measure theory. Since the notion of currents is defined not only for the Euclidean plane but also general curved surface, the formulation established here can be naturally generalized for surfaces. From the viewpoint of the application, it is of a great significance to justify the point vortex dynamics in a background field on curved surfaces as an Euler-Arnold flow, which is a generalization of the Euler equation to the case of the surfaces, since the point vortex dynamics in the rotational vector field on the unit sphere is adapted as a mathematical model of a geophysical flow in order to take effect of the Coriolis force on inviscid flows into consideration [5] for instance.

The main results consist of two theorems. For a current-valued solution of the Euler-Arnold equation with a regular-singular decomposition, we first prove that, if the singular part of the vorticity is given by a linear combination of delta functions centered at  $q_n(t)$  for  $n = 1, \dots, N$ ,  $q_n(t)$  is a solution of (3). Conversely, we next prove that, if  $q_n(t)$  is a solution of (3), there exists a current-valued solution of the Euler-Arnold equation with a regular-singular decomposition such that the singular part of the vorticity is given by a linear combination of delta functions centered at  $q_n(t)$ . Therefore, we conclude that the point vortex dynamics in a background field on a surfaces is a current-valued solution of the Euler-Arnold equation with a regular-singular decomposition.

## References

- [1] R.P.J. Kunnen, R.R. Trieling, and G.J.F. Heijst, van. Vortices in time-periodic shear. *Theoretical and Computational Fluid Dynamics*, 24(1-4):315–322, 2010.
- [2] A. J. Majda and A. L. Bertozzi. *Vorticity and incompressible flow*. Cambridge University Press, Cambridge, 2002.
- [3] C. Marchioro and M. Pulvirenti. Euler evolution for singular initial data and vortex theory. *Comm. Math. Phys.*, 91(4):563–572, 1983.
- [4] C. Marchioro and M. Pulvirenti. *Mathematical theory of incompressible nonviscous fluids*. Springer-Verlag, New York, 1994.
- [5] P. K. Newton and H. Shokraneh. The  $N$ -vortex problem on a rotating sphere. I. Multi-frequency configurations. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 462(2065):149–169, 2006.
- [6] P. G. Saffman. *Vortex dynamics*. Cambridge Monographs on Mechanics and Applied Mathematics. Cambridge University Press, New York, 1992.
- [7] R. R. Trieling, C. E. C. Dam, and G. J. F. van Heijst. Dynamics of two identical vortices in linear shear. *Physics of Fluids*, 22(11):117104, 2020/05/25 2010.

# Estimates of the regular set for Navier-Stokes flows in terms of initial data

Hideyuki Miura

Tokyo Institute of Technology  
miura@is.titech.ac.jp

This talk is based on a joint work with Kyungkeun Kang (Yonsei university) and Tai-Peng Tsai (University of British Columbia). We consider the regularity of weak solutions for the incompressible Navier-Stokes equations

$$\partial_t v - \Delta v + v \cdot \nabla v + \nabla p = 0, \quad \operatorname{div} v = 0 \quad (\text{NS})$$

associated with the initial value  $v|_{t=0} = v_0$  with  $\operatorname{div} v_0 = 0$ . In [2], Caffarelli, Kohn, and Nirenberg established local regularity theory for suitable weak solutions. As an application of their  $\epsilon$ -regularity theorem, they showed the following result:

**Theorem 1** ([2]). *There exists  $\epsilon_0 > 0$  such that if  $v_0 \in L^2(\mathbb{R}^3)$  satisfies*

$$\|v_0\|_{L^{2,-1}}^2 = \epsilon < \epsilon_0, \quad (1)$$

*then there exists a suitable weak solution which is regular in the set  $\Pi_{\epsilon_0-\epsilon}$ , where*

$$\begin{aligned} \|v_0\|_{L^{2,\alpha}} &:= \| |x|^{\frac{\alpha}{2}} v_0 \|_{L^2(\mathbb{R}^3)}, \\ \Pi_\delta &:= \left\{ (x, t) \in \mathbb{R}^3 \times (0, \infty) : t > \frac{|x|^2}{\delta} \right\}. \end{aligned}$$

This theorem asserts that smallness of initial data in a weighted space implies regularity of the solution above a paraboloid with vertex at the origin. There are at least two interesting features in this result: No regularity condition (better than  $L^2$ ) is assumed away from the origin and the regularity around the origin is propagated globally in time. We also note that if the size of  $v_0$  tends to 0,  $\Pi_{\epsilon_0-\epsilon}$  increases and converges to a limit set  $\Pi_{\epsilon_0}$ . This observation leads to the following questions: (a) Can the size of regular set  $\Pi_\delta$  be enlarged? (b) Can the assumptions of the initial data be relaxed in terms of regularity, decay, and smallness? The question (a) is addressed by D'Ancona and Luca [3], where it is shown that there exists  $\delta_0 > 0$  such that if  $v_0 \in L^2(\mathbb{R}^3)$  satisfies

$$\|v_0\|_{L^{2,-1}} < \delta_0 e^{-4L^2} \quad (2)$$

for some  $L > 1$ , (NS) has a suitable weak solution which is regular in the set  $\Pi_{L\delta_0}$  invading the whole half space  $\mathbb{R}^3 \times (0, \infty)$  when  $v_0$  tends to zero, though (2) still assumes smallness of the data. The aim of this talk is trying to answer questions (a) and (b). To this end, we recall the notion of the local energy solutions introduced by [4] and later modified by [1]. The local energy solution is a suitable weak solution of (NS) defined in  $\mathbb{R}^3$  which satisfies certain uniformly local energy bounds and pressure representation.

In this context, let us also recall the uniformly local  $L^q$  spaces for  $1 \leq q < \infty$ . We say  $f \in L^q_{\text{uloc}}$  if  $f \in L^q_{\text{loc}}(\mathbb{R}^3)$  and

$$\|f\|_{L^q_{\text{uloc}}} = \sup_{x \in \mathbb{R}^3} \|f\|_{L^q(B_1(x))} < \infty.$$

The following result shows estimates of the regular set for the local energy solution for arbitrary initial data in  $L^{2,-1}(\mathbb{R}^3)$  and the data with small scaled energy.

**Theorem 2.** *Let  $(v, p)$  be a local energy solution in  $\mathbb{R}^3 \times (0, \infty)$  for the initial data  $v_0 \in L^2_{\text{uloc}}(\mathbb{R}^3)$ .*

(i) *For any  $v_0 \in L^{2,-1}(\mathbb{R}^3)$  there exist positive constants  $T(v_0)$  and  $c(v_0)$  such that  $v$  is regular in the set*

$$\{(x, t) \in \mathbb{R}^3 \times (0, \infty) : c(v_0)|x|^2 \leq t < T(v_0)\}.$$

(ii) *There exist positive absolute constants  $\epsilon_*$  and  $c$  such that if  $v_0$  satisfies*

$$\dot{N}_0 := \sup_{r>0} \frac{1}{r} \int_{B_r} |v_0(x)|^2 dx \leq \epsilon_*$$

and

$$\sup_{x_0 \in \mathbb{R}^3} \sup_{r \geq 1} \frac{1}{r} \int_{B_r(x_0)} |v_0(x)|^2 dx < \infty,$$

then  $v$  is regular in the set

$$\{(x, t) \in \mathbb{R}^3 \times (0, \infty) : c\dot{N}_0^2|x|^2 \leq t\}.$$

It is not difficult to see that the assumptions of (ii) are slightly weaker than those in Theorem 1. (ii) also refines the convergence rate of the regular set to  $\mathbb{R}^3 \times (0, \infty)$  in [3] as  $\|v_0\|_{L^{2,-1}}$  tends to zero.

## References

- [1] Bradshaw, Z. and Tsai, T.-P., Global existence, regularity, and uniqueness of infinite energy solutions to the Navier-Stokes equations, *Comm. Partial Differential Equations* (to appear).
- [2] Caffarelli, L., Kohn, R., and Nirenberg, L., Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Comm. Pure Appl. Math.* 35 (1982), no. 6, 771-831.
- [3] D'Ancona, P. and Lucá R., On the regularity set and angular integrability for the Navier-Stokes equation. *Arch. Ration. Mech. Anal.* 221 (2016), no. 3, 1255–1284.
- [4] Lemarié-Rieusset, P. G., *Recent developments in the Navier-Stokes problem*. Chapman Hall/CRC Research Notes in Mathematics, 431. Chapman Hall/CRC, Boca Raton, FL, 2002.

# Existence of radially symmetric stationary solutions for the compressible Navier-Stokes equation

Itsuko Hashimoto

Kanazawa university / OCAMI

itsuko@se.kanazawa-u.ac.jp

(joint work with Akitaka Matsumura of Osaka university)

In this talk, we consider the existence of radially symmetric problems in  $\mathbb{R}^n (n \geq 2)$  to the compressible Navier-Stokes equation:

$$\begin{cases} (r^{n-1}\rho)_t + (r^{n-1}\rho u)_r = 0, \\ (\rho u)_t + (\rho u^2 + p(\rho))_r + (n-1)\frac{\rho u^2}{r} = \mu\left(\frac{(r^{n-1}u)_r}{r^{n-1}}\right)_r, \end{cases} \quad t > 0, \quad r > r_0, \quad (1)$$

where  $\rho = \rho(t, r) > 0$  is the mass density,  $u = u(t, r)$  is the fluid velocity, and  $p = p(\rho)$  is the pressure given by a smooth function of  $\rho$  satisfying  $p'(\rho) > 0$  ( $\rho > 0$ ), and  $\mu$  is a positive constant. We consider the initial boundary value problems to (1) under the initial condition

$$(\rho, u)(0, r) = (\rho_0, u_0)(r), \quad r > r_0,$$

the far field condition

$$\lim_{r \rightarrow \infty} (\rho, u)(t, r) = (\rho_+, u_+), \quad t > 0,$$

and also the following two types of boundary conditions depending on the sign of the velocity on the boundary

$$\begin{cases} (\rho, u)(t, r_0) = (\rho_-, u_-), & t > 0, \quad (u_- > 0), \\ u(t, r_0) = u_-, & t > 0, \quad (u_- \leq 0), \end{cases}$$

where  $\rho_{\pm} > 0, u_{\pm}$  are given constants. The case  $u_- > 0$  is known as ‘‘inflow problem’’, the case  $u_- = 0$  as ‘‘impermeable wall problem’’, and the case  $u_- < 0$  as ‘‘outflow problem’’. The equation (1) is given by letting

$$\rho(t, x) = \rho(t, r), \quad U(t, x) = \frac{x}{r} u(t, r), \quad r = |x|,$$

for the compressible Navier-Stokes equation which describes a barotropic motion of viscous gas in the exterior domain  $\Omega$  to a ball in  $\mathbb{R}^n (n \geq 2)$ :

$$\begin{cases} \rho_t + \operatorname{div}(\rho U) = 0, \\ (\rho U)_t + \operatorname{div}(\rho U \otimes U) + \nabla p = \nu \Delta U + (\nu + \lambda) \nabla(\operatorname{div} U), \end{cases} \quad t > 0, \quad x \in \Omega,$$

where  $\Omega = \{x \in \mathbb{R}^n (n \geq 2); |x| > r_0\}$  ( $r_0$  is a positive constant),  $U = (u_1(t, x), \dots, u_n(t, x))$  is the fluid velocity and  $\nu$  and  $\lambda$  are constants satisfying  $\nu > 0, 2\nu + n\lambda > 0$ .

We show the existence of a unique radially stationary solution in a suitably small neighborhood of the far field state for both inflow and outflow problems for the problem (1). Furthermore, it is shown that the boundary layer of the density appears as the velocity data tend to zero in the inflow problem, but not in the outflow problem. The stationary problem corresponding to the problem (1) is written as

$$\begin{cases} (r^{n-1}\rho u)_r = 0, \\ \rho u u_r + p(\rho)_r = \mu \left( \frac{(r^{n-1}u)_r}{r^{n-1}} \right)_r, & r > r_0, \\ \lim_{r \rightarrow \infty} (\rho, u)(r) = (\rho_+, u_+), \\ (\rho, u)(r_0) = (\rho_-, u_-) \quad (u_- > 0), \quad u(r_0) = u_- \quad (u_- \leq 0). \end{cases} \quad (2)$$

The main theorem of this talk is as follows.

**Theorem 1.** *Let  $n \geq 2$  and  $u_+ = 0$ . Then, for any  $\rho_+ > 0$ , there exist positive constants  $\epsilon_0$  and  $C$  satisfying the following:*

(I) *Let  $u_- > 0$ . If  $|u_-| + |\rho_- - \rho_+| \leq \epsilon_0$ , there exists a unique smooth solution  $(\rho, u)$  of the problem (2) satisfying*

$$\begin{aligned} |\rho(r) - \rho_+| &\leq Cr^{-(n-1)}(|u_-|^2 + |\rho_- - \rho_+|), \\ C^{-1}r^{-(n-1)}|u_-| &\leq |u(r)| \leq Cr^{-(n-1)}|u_-|, \quad r \geq r_0. \end{aligned}$$

Furthermore, for any positive constant  $h$ , there exists a positive constant  $C_h$  such that it holds

$$\sup_{r \geq r_0+h} |\rho(r) - \rho_+| \leq C_h |u_-|^2.$$

(II) *Let  $u_- \leq 0$ . If  $|u_-| \leq \epsilon_0$ , there exists a unique smooth solution  $(\rho, u)$  of the problem (2) satisfying*

$$\begin{aligned} |\rho(r) - \rho_+| &\leq Cr^{-2(n-1)}|u_-|^2, \\ C^{-1}r^{-(n-1)}|u_-| &\leq |u(r)| \leq Cr^{-(n-1)}|u_-|, \quad r \geq r_0. \end{aligned}$$

## References

- [1] S. Jiang, Global spherically symmetric solutions to the equations of a viscous polytropic ideal gas in an exterior domain, *Comm. Math. Phys.* 178 (1996), 339-374.
- [2] T. Nakamura, S. Nishibata, and S. Yanagi, Large-time behavior of spherically symmetric solutions to an isentropic model of compressible viscous fluid in a field of potential forces, *Math. Models Methods Appl. Sci.* 14 (2004), 1849-1879.