

pole

$$\frac{\nabla}{ds} T_\alpha + \frac{1}{2} \sum_{\beta, \gamma} \text{expr}[T_\beta, T_\gamma] = 0$$

Suppose $T_\alpha = \frac{\alpha_\alpha}{s} + \text{higher}$

$$\Rightarrow -\frac{\alpha_\alpha}{s^2} + \frac{1}{2} \sum \text{expr}\left[\frac{\alpha_\beta}{s}, \frac{\alpha_\gamma}{s}\right] + \dots = 0$$

$$\therefore \alpha_\alpha = \frac{1}{2} \text{expr}[\alpha_\beta, \alpha_\gamma] \quad \alpha_1 = [\alpha_2, \alpha_3] \text{ etc}$$

$$SU(2) \cong Sp(1) \cong \text{Im } \mathbb{H} \quad \frac{1}{2}i = \left(\frac{1}{2}j, \frac{1}{2}k\right) \text{ etc}$$


$$\alpha \frac{1}{2} \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$$

$\therefore i, j, k \mapsto 2\alpha_1, 2\alpha_2, 2\alpha_3$ defines a Lie alg. from.

complexification $\rho: SU(2) \rightarrow U(k)$
 $\rho: SL(2) \rightarrow GL(k)$

$$e = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, f = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, h = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

$s_- \quad s_+$



can put P_\pm at s_\pm

For Nahm's equations, corresponding to monopoles
 \Rightarrow Both P_\pm irreducible for s_\pm

But one can put **any** P_\pm

Classification of \mathfrak{p}_\pm

$p: \mathfrak{su}(2) \rightarrow \mathfrak{u}(\mathbb{R})$ ($p: \mathfrak{sl}(2) \rightarrow \mathfrak{sl}(\mathbb{R})$)
 \leftrightarrow \mathbb{R} -dim'l representation of $SL(2)$
 \leftrightarrow partition of \mathbb{R} $(1^{n_1} 2^{n_2} \dots)$ s.t. $\sum i n_i = k$

$$\mathcal{V} = \bigoplus V(i)^{\otimes n_i}$$

↑
i-dim'l irr. rep.

multiplicity

\leftrightarrow nilpotent elements of $\mathfrak{sl}(\mathbb{R})$ up to conjugacy
Jordan normal form

This correspondence holds for arbitrary simple cpx Lie algebra / \mathbb{C}

Th. (Jacobson - Morozov + Kostant)

$$p: \mathfrak{sl}(2) \rightarrow \mathfrak{g} / \text{conj.} \leftrightarrow \text{nilpotent element } e \in \mathfrak{g} / \text{conj.}$$

$$p \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = e$$

ex

$$e = \begin{bmatrix} 0 & 1 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix}, f = \begin{bmatrix} 0 & & & \\ n-1 & 0 & & \\ \vdots & \vdots & \ddots & 0 \\ 2(n-2) & \dots & \dots & n-1 \end{bmatrix}, h = \begin{bmatrix} n-1 & & & 0 \\ n-3 & \dots & & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 1 & \dots & 0 & n \end{bmatrix}$$

$$[e, f] = \begin{bmatrix} n-1 & & & 0 \\ \vdots & \vdots & \ddots & 0 \\ 2(n-2) & \dots & \dots & n-1 \\ 0 & \dots & 0 & n \end{bmatrix} - \begin{bmatrix} 0 & n-1 & & 0 \\ 0 & 2(n-2) & \dots & 0 \\ \vdots & \vdots & \ddots & n-1 \\ 0 & 0 & \dots & n \end{bmatrix}$$

$$k(n-k) - \frac{(k-1)(n-k+1)}{'' k(n-k) - n+k+k-1} = n-2k+1$$

$$[h, e] = 2e$$

Last time : both p_{\pm} = trivial ($= 1^k$)
 \Rightarrow moduli sp. $\cong T^*GL(k)$

This is true for any G^*

Th. (Bielański)

p_- : arbitrary , p_+ = trivial
 \Rightarrow moduli space $\cong G^* \times S(p_-)$

where $S(p_-) =$ Slodowy slice
= $e + Z_{G^*}(f)$

$$e = p_- \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$f = p_- \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

p_- : trivial $\Rightarrow f = 0 \Rightarrow S(p_-) = G^*$
recover the previous theorem

On Slodowy slice


$$g^C \ni e \quad T_e g^C = g^C$$

$O(e) =$ nilpotent orbit through e

$$T_e O(e) = [g^C, e] = \text{Im}(ad e)$$

$$g^C = T_e O(e) \oplus Z_{g^C}(f) = \text{Im}(ad e) \oplus \text{Ker}(ad f)$$

∴ sl₂-rep. theory \Rightarrow irr. rep of sl₂ looks like



$$\Rightarrow S(\rho_-) \cap O(e) = \{e\} \quad \text{in a nbd of } e$$

$$\mathbb{C}^* \rightarrow S(\rho_-) \quad t^2 \underbrace{(\text{Ad}(t^{-\tilde{h}})(e+s))}_{\substack{\parallel \\ \rho \begin{bmatrix} t^{-1} & 0 \\ 0 & t \end{bmatrix}}}$$

$$[\tilde{h}, e] = 2e \Rightarrow e: \text{fixed}$$

$t^{-\tilde{h}}|_{Z_{g^C}(f)}$: nonpositive eigenvalues

\Rightarrow above action is contracting.

∴ Any pt in $S(\rho)$ can be moved to a n.b.d of e .

$$\therefore S(\rho_-) \cap O(e) = \{e\} \quad \text{everywhere}$$

sketch of the proof of Birulański's Thm

cpx description $\left\{ \begin{array}{l} \alpha = \frac{1}{2}(P_0 + iP_1) \\ \beta = \frac{1}{2}(P_2 + iP_3) \end{array} \right.$

cpx eqn. : $\frac{d}{ds}\beta + 2[\alpha, \beta] = 0$ $- \frac{e}{2} + 2\left[\frac{\hbar}{4}, \frac{e}{2}\right] = 0$
 $[e, e] = 2e \checkmark$

Step 1^o $\left\{ \text{solutions of } \begin{array}{l} \text{Nahm's eqn} \\ \text{cpx equation} \end{array} \right\} / g_{00}^C \cong \left\{ \text{solutions of } \begin{array}{l} \text{Donaldson's thm (taking limit)} \end{array} \right\} / g_{00}^C$

Step 2^o $\left(\text{Res } \alpha = \frac{i}{2} \text{Res } P_1 = \frac{i}{4} P \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix} = \frac{\hbar}{4} \right)$
 $\text{Res } \beta = \frac{1}{2}(\text{Res } P_2 + i \text{Res } P_3) = \frac{P}{2} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \frac{e}{2}$

Put $\left\{ \begin{array}{l} \overline{\alpha} = 2s\alpha \\ \overline{\beta} = 2s\beta \end{array} \right.$ $s = \frac{1}{2}e^{-2t}$ $s=0 \dots t=\infty$
 $s=s_+ \dots t: \text{finite}$

$$\frac{d\overline{\beta}}{dt} = -2\overline{\beta} + 2[\overline{\alpha}, \overline{\beta}]$$

After a cpx gauge transform,
we may assume

$$\overline{\alpha} = \frac{\hbar}{2} \quad (\text{i.e. } \alpha = \frac{\hbar}{4s}) \quad \text{in a nbd of } t=\infty$$

$$\Rightarrow \overline{\beta} = e^{-2t} \text{Ad}(e^{t\frac{\hbar}{2}})(\overline{\beta}_0) \quad \overline{\beta}_0 := \overline{\beta}|_{t=0} \quad \text{constant}$$

(i.e. $\beta = \text{Ad}((2s)^{-\frac{\hbar}{2}})(\overline{\beta}_0)$)

$$\left(\begin{aligned} \text{LHS} &= 2\left(\frac{ds}{dt}\beta + s\frac{ds}{dt}\frac{d\beta}{ds}\right) \\ &= 2 \cdot \underbrace{\left(-e^{-2t}\right)}_{2s} \left(\beta + s\frac{d\beta}{ds}\right) \\ &= \underbrace{RHS}_{-2[\alpha, \beta]} \end{aligned} \right)$$

Decompose :

$$g^C = \bigoplus g^C_{(n)} \quad , \quad g^C_{(n)} = \bigoplus_m g^C_{(m)}$$

\uparrow eigenvalue of $\text{ad } h$

Need $\overline{\beta}_{(t \rightarrow \infty)} \rightarrow 2e \Rightarrow \overline{\beta}_0 \in 2e + \mathcal{G}^C(<2)$

We can further make a cpx gauge transform $\gamma(t)$ s.t. it does not change $\overline{\alpha} = h$

$$\overline{\alpha}^\gamma = \gamma^{-1} \overline{\alpha} \gamma - \gamma^{-1} \frac{d\gamma}{dt} \quad \therefore \frac{d\gamma}{dt} = [h, \gamma]$$

$\overline{\alpha}$ γ $\frac{d\gamma}{dt}$
|| γ $-h$ $-h$: connection form

$$\therefore \gamma = e^{th} \gamma_0 e^{-th}$$

Need $\gamma(t) \rightarrow 1 (t \rightarrow \infty) \Rightarrow \gamma_0 \in \exp(\mathcal{G}^C(<0))$

Then $\overline{\beta}^\gamma = \gamma^{-1} \overline{\beta} \gamma = e^{-2t} \text{Ad}(e^{th})(\gamma_0^{-1} \overline{\beta} \gamma_0)$

$$\therefore \overline{\beta}_0 \mapsto \gamma_0^{-1} \overline{\beta}_0 \gamma_0$$

Prop. $\exp(\mathcal{G}^C(<0)) \times S(p_-) \xrightarrow[\Phi]{\cong} e + \mathcal{G}^C(<2)$

∴ Take differential at $(1, e)$

$$d\Phi(\bar{z}, z) = [\bar{z}, e] + z$$


$$\mathcal{G}^C(<0) \stackrel{\Phi}{\cong} \mathcal{G}^C(f) \Rightarrow \Phi: \text{local isom.}$$

Use \mathbb{C}^* -action as before //

$$\therefore \text{moduli sp. } \cong \overset{\uparrow}{G^C} \times S(p)$$

value of the gauge transf. g s.t. $(\alpha, \beta)^g$ is of the above form at the other end S_+

Finally we consider




P_- P_+ : irreducible
rep. of $\mathrm{SL}(2)$

Th. (Donaldson)

moduli sp. \cong the space of based maps $P \xrightarrow{\text{deg } k} P'$
 $\infty \mapsto \infty$

We divide



solve ODE from here
 (X, v)

Lemma. $G^c \times S(P) \xrightarrow{\cong} \{(X, v) \in gl(\ell) \times \mathbb{C}^\ell \mid v: \text{cyclic vector}$ for X

$$\mathbb{C}^\ell = \text{Span}\langle v, Xv, \dots, X^{\ell-1}v \rangle$$


$$(g, e+z) \mapsto (-g(e+z)g^{-1}, g \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix})$$

Rmk. $\begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$: cyclic vector for $e \Rightarrow$ for $e+z$: small z
 \Rightarrow for any $e+z$

$\therefore \text{LHS} \rightarrow \text{RHS}$ is defined.

Using this, one can show:

moduli sp. $\cong \{(X, v_-, v_+) \mid \begin{array}{l} v_-: \text{cyclic} \\ t v_+: \text{cyclic for } tX \end{array}\} / GL(\ell)$



Prop $\{(X, v_-, v_+) : \text{as above}\} /_{\text{GL}(k)} \longrightarrow \{\text{based maps } \mathbb{P}^1 \rightarrow \mathbb{P}^1 \text{ of deg } = k\}$

$$f(z) = v_+ (z - X)^{-1} v_-$$

$$(f(\infty) = 0)$$

quasimaps : $\{(X, v_-, v_+) : \text{arbitrary}\} //_{\text{GL}(k)}$
 $\uparrow \text{set of closed orbits}$