

# Coulomb branches and cyclotomic rational Cherednik algebras

Hiraku Nakajima (RIMS, Kyoto)

2016-09-26 Skoltech (Moscow)

- 1503.03676 Nakajima
- 1601.03586 Braverman-Finkelberg-Nakajima
- 1604.03625 \_\_\_\_\_
- its appendix Braverman-Finkelberg-Kamnitzer-Kodera-Nakajima-Webster-Weekes
- 1608.00875 Kodera-Nakajima

# Coulomb branches of 3D N=4 SUSY gauge theories

Take  $G_c$ : a compact Lie group,  $G$  its complexification

$M$ : a quaternionic representation of  $G_c$

Physics (a symplectic representation of  $G$ )

$\Rightarrow$  3D N=4 SUSY gauge theory (4D N=2 as well)

$\rightsquigarrow \mathcal{M}_c \equiv \mathcal{M}_c(G, M)$  : Coulomb branch

a noncompact hyperKähler manifold

possibly with singularities

and  $SU(2)$ -action rotating complex structures

Very roughly

gauge theory  $\cong$  3D  $\sigma$ -model with target  $\mathcal{M}_C$   
(when  $\mathcal{M}_H = \{0\}$  and  $\mathcal{M}_C$  is smooth)

 Higgs branch = hyperKähler quotient  $M // G_C$

1996 Seiberg-Witten

$\mathcal{M}_C(G_C, M)$  = Atiyah-Hitchin manifold

$\mathcal{M}_C(\mathbb{Z}, 0)$  = moduli of centered  $SU(2)$  charge 2  
magnetic monopoles on  $\mathbb{R}^3$

Then many subsequent works computing  $\mathcal{M}_C$  in  
examples.....

But the definition of  $\mathcal{M}_C$  was not clear to mathematicians (e.g., me).

17 years later

2013 Cremonesi-Hanany-Zaffaroni

combinatorial expression (monopole formula) of

$$\text{ch}_{\mathbb{C}^*}(\underline{\mathbb{C}[\mathcal{M}_C]})$$

coordinate ring of  $\mathcal{M}_C$  (chiral ring)


The formula is mathematically meaningful!

It motivated me to look for a mathematical definition.

## A proposal of a mathematical definition

Idea : Suppose we have a TQFT given by twisting of a SUSY gauge theory.

Then  $Z(S^2) = (\text{Hilbert space for } S^2)$  is



a commutative ring acting on  $Z(\Sigma_g)$  as

$$\partial M = S^2 \cup \Sigma_g \cup -\Sigma_g$$

$$\therefore Z(M) \in \text{Hom}(Z(S^2) \otimes Z(\Sigma_g), Z(\Sigma_g))$$

$$\text{Hence } \mathbb{C}[\mathcal{M}_c] = Z(S^2)$$

Moreover, in mathematical works,  $Z(\Sigma_g)$  is defined by cohomology groups of moduli spaces  $\Big|_{G_c = SU(2), M=0} \Big|_{G_c = U(1), M=\mathbb{H}}$  of a nonlinear PDE (e.g., flat connection, Seiberg-Witten)

Combining it with heuristic consideration and  
monopole formula, we arrive at the following:

Assume  $M = N \oplus N^*$ .


$\mathcal{R}$  = the space of Hecke correspondences with sections  
= moduli stack of  $(P_1, P_2, \varphi, s)$

$P_i$  : holomorphic  $G$ -b'dle on  $D$  (formal disk)

$\varphi : P_1|_{D \setminus \{0\}} \xrightarrow{\cong} P_2|_{D \setminus \{0\}}$  isomorphism

$s$  : holomorphic section of  $P_1 \times_G N$

s.t.  $\varphi(s)$  has no pole at 0





## Lemma

commutative

$$\mathbb{C}[\mathcal{M}_c] \stackrel{\text{def.}}{=} H_*^{\text{BM}}(\mathcal{R}) + \text{convolution product}$$

\* This gives a definition of  $\mathcal{M}_c$  as an affine algebraic variety

- \*  $\mathbb{C}^*$ -action is given by homological degrees with correction
- \*  $ch_{\mathbb{C}^*} \mathbb{C}[\mathcal{M}_c] = \text{monopole formula (when the degree } \geq 0)$

Example  $G = \mathbb{C}^*, N = 0$

$P_2 \cong$  trivial line bundle  $\varphi = \sum^k \quad (k \in \mathbb{Z})$

$$\therefore \mathcal{R} = \dots \overset{-2}{x} \overset{-1}{x} \overset{0}{x} \overset{1}{x} \overset{2}{x} \dots$$

$\nearrow \nearrow$  point /  $\mathbb{C}^* = B\mathbb{C}^*$

$$\therefore H_*^{BM}(\mathcal{R}) = \bigoplus_{k \in \mathbb{Z}} H_*(B\mathbb{C}^*) \cong \mathbb{C}[u, x, x^{-1}]$$

$$\therefore \mathcal{M}_C = \mathbb{C} \times \mathbb{C}^* (= T^*\mathbb{C}^*)$$

Remark 1) When  $G =$  torus,  $N =$  arbitrary,

$\mathbb{C}[\mathcal{M}_C]$  has an explicit presentation.

$$2) \quad \pi_0(\mathcal{R}) \cong \pi_1(G) \quad (\text{in general})$$

$\Rightarrow \mathcal{M}_C$  has an additional  $\pi_1(G)^{\wedge}$ -action.  $\nwarrow$  Pontryagin dual




## Quantization

Consider  $\mathbb{C}^*$ -action on the formal disk  $\mathbb{D}$

The equivariant homology  $H_*^{\mathbb{C}^*}(\mathcal{R})$

- is a deformation of  $H_*(\mathcal{R})$  over  $\mathrm{Spec} H_{\mathbb{C}^*}^*(pt) = \mathbb{C}$
- has a convolution product, but **noncommutative!**



↻ cannot be swapped!

Def.  $\mathcal{A}_\hbar = H_*^{\mathbb{C}^*}(\mathcal{E})$  : quantized Coulomb branch

Example  $G = \mathbb{C}^*, N = 0$

$$\mathcal{A}_\hbar = \langle w, x, x^{-1} \rangle \quad [x^\pm, w] = \hbar x^\pm$$

So  $x^\pm = \mathcal{Q}_w^\pm : f(w) \mapsto f(w \pm \hbar)$  **difference operators**

Lemma  $\mathcal{A}_\hbar(G, N) \xrightarrow{\star} \mathcal{A}_\hbar(T, 0)[\text{root}^{-1}]$

difference operators with  
poles along root hyperplanes

Q. How to compute the image ?

When  $G = GL(\mathbb{R})$  (more generally  $G = \prod GL(\mathbb{R}_i)$ )

$$\mathcal{R} \supset \{ (E_1, E_2, \varphi, s) \mid E_1 \subset E_2 \subset E_1(0) \}$$

or  $E_2 \subset E_1 \subset E_2(0)$




**smooth** closed subvariety

The image of its fundamental class under  $\star$   
can be computed by the fixed point formula !

cf. Bullimore, Dimofte, Gaiotto 1503.04817

Example  $G = GL(k)$ ,  $N = \mathfrak{gl}(k) \oplus (\mathbb{C}^k)^{\oplus 2}$

$$1 \leq n \leq k$$



$$F_n = \sum_{\substack{I \subset \{1, \dots, k\} \\ \#I = n}} \prod_{\substack{i \in I \\ j \notin I}} \frac{w_i - w_j + 1}{w_i - w_j} \prod_{i \in I} \prod_{a=1}^2 (w_i - z_a) \mathcal{D}_{w_i}^{-1}$$

(generalized) Macdonald operators

Moreover, these operators generate  $\mathcal{A}_k$ .



Th.

spherical part of

$A_h \cong$  cyclotomic rational Cherednik algebra

in this example

$$\mathbb{C}S_R \rtimes (\mathbb{Z}/l\mathbb{Z})^R$$

Remark 1)  $\mathcal{U}_C \cong \text{Sym}^k(\mathbb{C}^2/(\mathbb{Z}/l\mathbb{Z}))$  de Boer, Hori, Ooguri, Oz 1997

2) finite ADE quiver  $\Rightarrow A_h$  truncated shifted Yangian