

Quasi-homomorphism Rigidity with Noncommutative Targets

OZAWA, Narutaka

RIMS at Kyoto University

Sendai Symposium, August 2011

Research partially supported by JSPS and HIM

Quasi-homomorphism Rigidity with Noncommutative Targets

OZAWA, Narutaka

RIMS at Kyoto University

Sendai Symposium, August 2011

Research partially supported by JSPS and HIM

Prologue: Zimmer's program



All groups are assumed second countable & locally compact.
All maps are assumed locally bounded & Borel measurable.

A “large” group should not act on a “small” manifold

Problem

Let Γ be a lattice in $SL(n, \mathbb{R})$.

Is every action of Γ on a (compact) mfd M of dimension $< n - 1$ finite?

The known results are mostly for $M = S^1$ (a circle), or $M = \mathbb{R}$ (a line).

Theorem

- (Witte 1994)

YES to the above Problem for finite index subgroups of $SL(n \geq 3, \mathbb{Z})$.

- (Ghys, Burger–Monod 1999)

Let Γ be a lattice in $SL(n \geq 3, \mathbb{R})$. Then, every action $\Gamma \curvearrowright S^1$ has at least one finite orbit, and every C^1 -action $\Gamma \curvearrowright S^1$ is finite.

- (Navas 2002)

Let Γ be a property (T) group. Then, every C^2 -action $\Gamma \curvearrowright S^1$ is finite.

Lattices of $SL(n \geq 3, \mathbb{R})$ have **property (T)** of Kazhdan.

Quasimorphisms

Let $\Gamma \curvearrowright S^1$. Each $g \in \Gamma$ has a lift $\tilde{g} \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$ with $\tilde{g}(0) \in [0, 1)$. Then,

$$c(g, h) = (\tilde{g}h)^{-1}\tilde{g}\tilde{h} \in \{0, 1\}$$

defines the Euler class e in the **bounded** cohomology $H_b^2(\Gamma, \mathbb{Z})$.

Theorem (Ghys 1987)

The Euler class $e \in H_b^2(\Gamma, \mathbb{Z})$ determines $\Gamma \curvearrowright S^1$ up to semi-conjugacy.

Under certain assumption (e.g., $H^2(\Gamma, \mathbb{R}) = 0$), the Euler cocycle c is a coboundary of a not-necessarily bounded map $q: \Gamma \rightarrow \mathbb{R}$.

The map q is a **quasimorphism**:

$$\sup_{g, h \in \Gamma} |q(gh) - (q(g) + q(h))| < +\infty.$$

\rightsquigarrow Want to show every quasimorphism on $\Gamma \leq \text{SL}(3, \mathbb{R})$ is bounded.

Property (T) and what it is good for



All groups are assumed second countable & locally compact.
All maps are assumed locally bounded & Borel measurable.

Kazhdan's property (T)

Definition/Theorem (Kazhdan '67, Delorme '77, Guichardet '72)

G has **property (T)** if it satisfies one of the following equiv conditions:

- The trivial representation is isolated in the unitary dual of G .
- For every unitary representation $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$, every cocycle $b: G \rightarrow \mathcal{H}$ is bounded. Here, a cocycle is a map b satisfying

$$\forall g, h \in G \quad b(gh) = b(g) + \pi(g)b(h).$$

Note: A cocycle b is bounded iff $\exists \xi \in \mathcal{H}$ s.t. $b(g) = \pi(g)\xi - \xi$.

Example

- Simple Lie groups of real rank ≥ 2 have property (T).
- A lattice Γ in G has property (T) iff G has property (T).
- $\mathrm{SL}(n, \mathbb{Z})$ has property (T) iff $n \geq 3$.
- Many hyperbolic groups, e.g. lattices in $\mathrm{Sp}(n, 1)$, have property (T).

Some consequences of Kazhdan's property (T)

Theorem (Kazhdan)

For a discrete group Γ with property (T), the following hold true.

- Γ is finitely generated.
- Γ has finite abelianization.
- For each n , Γ has only finitely many n -dimensional unitary reps, up to unitary equivalence.

Sketchy proof of the last statement.

By property (T), \exists a finite subset $E \subset \Gamma$ and $C > 0$ such that

$$\forall b \text{ cocycle} \quad \sup_{g \in G} \|b(g)\| \leq C \max_{s \in E} \|b(s)\|.$$

For unitary reps $\pi, \sigma: \Gamma \rightarrow \mathcal{U}(n)$, consider the unitary rep $\pi \otimes \bar{\sigma}$ on HS_n defined by $X \mapsto \pi(g)X\sigma(g)^*$, and the cocycle $b(g) = \pi(g)\sigma(g)^* - I_n$.

\leadsto If π and σ are close on E , then they are unitarily equivalent.



Beef up Kazhdan's property (T)

Definition (Kazhdan, Delorme, Guichardet, Burger–Monod,)

A group G has *property (TT)* if every **quasi**-cocycle on G is bounded. Here, a **quasi**-cocycle is a map $b: G \rightarrow \mathcal{H}$, together with $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$, which satisfies

- π is a representation, and
- b satisfies the ~~cocycle identity~~ rough cocycle inequality

$$\sup_{g,h} \| \cancel{b(gh)} - (\cancel{b(g)} + \pi(g)\cancel{b(h)}) \| < +\infty.$$

A quasimorphism is a quasi-cocycle with the trivial representation.

Theorem (Burger–Monod 1999, 2002)

The group $\mathrm{SL}(n, \mathbb{R})$ and its lattices have property (TT) for $n \geq 3$.

Results of Burger and Monod

Theorem (Burger–Monod 1999, 2002)

The group $\mathrm{SL}(n, \mathbb{R})$ and its lattices have property (TT) for $n \geq 3$.

There are groups having property (T), but not (TT): Hyperbolic groups do not have property (TT), because they have **proper** quasi-cocycles.

A cocycle $b: G \rightarrow \mathcal{H}$ is said to be proper if for any $C > 0$, the subset $\{g \in G : \|b(g)\| \leq C\}$ is relatively compact.

Corollary

Every quasimorphism on a lattice Γ in $\mathrm{SL}(n \geq 3, \mathbb{R})$ is bounded.

Corollary (Ghys, Burger–Monod)

Every action $\Gamma \curvearrowright S^1$ has at least one finite orbit.

Property (TTT) and what it is good for

Property (TTT) and what it is good for



All groups are assumed second countable & locally compact.
All maps are assumed locally bounded & Borel measurable.

Beef up Kazhdan's property (T) further

Definition (Kazhdan, Delorme, Guichardet, Burger–Monod, Oz.)

A group G has *property* (TTT) if every **wq**-cocycle on G is bounded. Here, a **wq**-cocycle is a map $b: G \rightarrow \mathcal{H}$, together with $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$, which satisfies

- ~~π is a representation, and~~
- b satisfies the ~~cocycle identity~~ rough cocycle inequality

$$\sup_{g,h} \| \cancel{b(gh)} - \cancel{b(g)} - \pi(g)\cancel{b(h)} \| < +\infty.$$

Theorem (Oz. 2009)

The group $\mathrm{SL}(n, \mathbb{R})$ and its lattices have property (TTT) for $n \geq 3$.

Quasi-homomorphisms

Definition

A map $q: G \rightarrow H$ is called a *quasi-homomorphism* if

$$\{q(gh)^{-1}q(g)q(h) : g, h \in G\}$$

is relatively compact in H .

If $b: H \rightarrow \mathcal{H}$ is a wq-cocycle and $q: G \rightarrow H$ is a quasi-homomorphism, then $b' = b \circ q$ is a wq-cocycle, because

$$b'(gh) = b(q(g)q(h) \square) \approx b'(g) + \pi'(g)b'(h).$$

⚡ Even if π is multiplicative, $\pi' = \pi \circ q$ is not.

Definition

A group H is called *a-TTT-menenable* if there is a proper wq-cocycle on H .

Examples: Abelian groups, solvable groups, amenable groups,
a-T-menenable (a.k.a. Haagerup) groups, hyperbolic groups. . .

Quasi-homomorphisms

Definition

A map $q: G \rightarrow H$ is called a *quasi-homomorphism* if

$$\{q(gh)^{-1}q(g)q(h) : g, h \in G\}$$

is relatively compact in H .

Definition

A group H is called *a-TTT-menenable* if there is a proper wq-cocycle on H .


Examples: Abelian groups, solvable groups, amenable groups,
a-T-menenable (a.k.a. Haagerup) groups, hyperbolic groups. . .

Corollary

If G has property (TTT) and H is a-TTT-menenable, then every quasi-homomorphism from G into H has relatively compact image.

Examples of quasi-homomorphisms. $q(gh)^{-1}q(g)q(h)$

- $\widetilde{\text{Homeo}}(S^1) = \{f \in \text{Homeo}(\mathbb{R}) : f(x+1) = f(x) + 1\}$
and $q: f \mapsto f(0) \in \mathbb{R}$.
 \rightsquigarrow Application to $\Gamma \curvearrowright S^1$ (Burger–Monod, Ghys).
- $q: \mathbb{F}_2 = \langle a, b \rangle \rightarrow \mathbb{Z}$,
 $q(w) = (\# \text{ of } ab \text{ occurs in } w) - (\# \text{ of } b^{-1}a^{-1} \text{ occurs in } w)$



Generalizes to
hyperbolic groups
(Epstein–Fujiwara).

⚡ Defect usually occurs around the joining area: $q(g)^{-1}q(gh)q(h)^{-1}$.
It's difficult to have quasi-homomorphisms with noncommutative targets.

Definition

For $\varepsilon > 0$, a (unitary) ε -representation of a group G on a Hilbert space \mathcal{H} is a map $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$ which satisfies

$$\sup_{g,h \in G} \|\pi(g)\pi(h) - \pi(gh)\| \leq \varepsilon.$$

Problem [S. M. Ulam, A collection of mathematical problems (1960).]

Is an ε -representation π close to a unitary representation?

Kazhdan (1982): **YES!** for amenable groups, and **NO!** in general.

\rightsquigarrow **NO!** for any group which contains \mathbb{F}_2 .

Example (From a quasimorphism to a quasi-character)

Let $q: \Gamma \rightarrow \mathbb{R}$ be a quasimorphism with $\sup |q(gh) - (q(g) + q(h))| \leq 1$. Then $\pi(g) := \exp(i\varepsilon q(g))$ is an ε -character. For ε sufficiently small, π is close to a character iff q is a bounded distance from a homomorphism.

ε -representations and property (TTT)

Definition

For $\varepsilon > 0$, a (unitary) ε -representation of a group G on a Hilbert space \mathcal{H} is a map $\pi: G \rightarrow \mathcal{U}(\mathcal{H})$ which satisfies


$$\sup_{g,h \in G} \|\pi(g)\pi(h) - \pi(gh)\| \leq \varepsilon.$$

An example of ε -representations (Rolli 2009)

Let $\mathbb{F}_2 = \langle a, b \rangle$ and $B(\varepsilon/3) = \{u \in \mathcal{U}(\mathcal{H}) : \|u - 1\| \leq \varepsilon/3\}$.

Fix symmetric functions $\sigma_a, \sigma_b: \mathbb{Z} \rightarrow B(\varepsilon/3)$ and set

$$\pi(a^{m_1} b^{n_1} \cdots a^{m_k} b^{n_k}) = \sigma_a(m_1) \sigma_b(n_1) \cdots \sigma_a(m_k) \sigma_b(n_k).$$



Theorem (B.O.T.; Dimension dependent Ulam stability)

Let Γ be a property (TTT) group. Then, any ε -representation $\pi: \Gamma \rightarrow \mathcal{U}(d)$ with $\varepsilon < \kappa(d)$ is close to a unitary representation.

Theorem (Kazhdan and Burger–Oz.–Thom)

- If Γ is amenable, then every ε -repn is 2ε -close to a unitary repn.
- If $\mathbb{F}_2 \hookrightarrow \Gamma$, then for each $\varepsilon > 0$, \exists ε -repn which is not close to any unitary repn.
- If Γ has property (TT), then every 1-dim ε -repn is $\delta(\varepsilon)$ -close to a unitary repn.
- If Γ has property (TTT), then every d -dim ε -repn is $\delta_d(\varepsilon)$ -close to a unitary repn.
- If $\Gamma = \mathrm{SL}(n \geq 3, \mathbb{Z})$, then every finite-dim ε -repn is $\delta(\varepsilon)$ -close to a unitary repn. The same thing for certain $\mathrm{SL}(2, A)$.

Are two ε -close unitary reps of Γ necessarily unitarily equivalent?


YES if Γ amenable (or unitarizable), and **NO** if $\mathbb{F}_2 \hookrightarrow \Gamma$.

Proof of Property (TTT)



All groups are assumed second countable & locally compact.
All maps are assumed locally bounded & Borel measurable.

Proof of Property (TTT)



All groups are assumed second countable & locally compact.
All maps are assumed locally bounded & Borel measurable.

Relative property (TTT)

Definition

A subgroup $A \leq G$ has *relative property* (TTT) if every wq-cocycle on G is bounded on A .

Theorem

Let A be abelian and $G = G_0 \rtimes A$. Then, for $A \leq G$,
relative property (TTT) \iff *relative property* (T)

The proof is à la Burger, but goes with positive definite kernels

$$\theta_t(g, h) = \exp(-t\|\delta(g) - \delta(h)\|^2)$$

instead of positive type functions.

Bounded generation and property (TTT) for $\mathrm{SL}(n, \mathbb{K})$

Theorem

Let A be abelian and $G = G_0 \rtimes A$. Then, for $A \leq G$,
relative property (TTT) \iff relative property (T)

Corollary

For $n \geq 3$, the group $\mathrm{SL}(n, \mathbb{K})$ has property (TTT).

Proof for $n = 3$.

By relative property (T) for $\mathbb{K}^2 \leq \mathrm{SL}(2, \mathbb{K}) \rtimes \mathbb{K}^2$, every wq-cocycle b on $\mathrm{SL}(3, \mathbb{K})$ is bounded on $\begin{pmatrix} 1 & 0 & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix}$, and on any other elementary matrices. Since every element of $\mathrm{SL}(3, \mathbb{K})$ is a product of at most 10 elementary matrices, the wq-cocycle b is bounded on $\mathrm{SL}(3, \mathbb{K})$. \square

$$\sup_{g,h} \|b(gh) - (b(g) + \pi(g)b(h))\| < +\infty.$$

Lattices


Let G be a property (TTT) group and $\Gamma \leq G$ be a *cocompact* lattice.
Let $X = G/\Gamma$ and choose a section $\sigma: X \rightarrow G$.

Define the Borel cocycle $\beta: X \times G \rightarrow \Gamma$ by

$$\beta(x, g) = \sigma(x)^{-1} g \sigma(g^{-1}x).$$

It satisfies the cocycle identity:

$$\beta(x, gh) = \beta(x, g)\beta(g^{-1}x, h).$$



To prove that Γ has property (TTT), let a wq-cocycle $b: \Gamma \rightarrow \mathcal{H}$ be given, and $\tilde{b}: G \rightarrow L^2(X, \mathcal{H})$ be the induced wq-cocycle on G defined by

$$\tilde{b}(g)(x) = b(\beta(x, g)),$$

together with $\tilde{\pi}: G \rightarrow \mathcal{U}(L^2(X, \mathcal{H}))$, $(\tilde{\pi}(g)\xi)(x) = \pi(\beta(x, g))\xi(g^{-1}x)$.

Problem

If we know \tilde{b} is bounded on G , does it follow b is bounded on Γ ?

Lattices and semi-length functions

Problem

If we know \tilde{b} is bounded on G , does it follow b is bounded on Γ ?

Burger–Monod: The answer is **YES!** if b is a quasi-cocycle, because the L^2 -induction $H_b^2(\Gamma, \mathcal{H}) \rightarrow H_{cb}^2(G, L^2(X, \mathcal{H}))$ is injective.

In general,

let $C := \sup \|b(gh) - (b(g) + \pi(g)b(h))\|$ and $\ell(g) := \|b(g)\| + C$.

Then, ℓ is a semi-length function: $\ell(gh) \leq \ell(g) + \ell(h)$.

The induced semi-length function $L: G \rightarrow \mathbb{R}_{\geq 0}$ is given by

$$L(g) = \int_X \ell(\beta(x, g)) \, dx.$$

The above problem generalizes to

Problem

If we know L is bounded on G , does it follow ℓ is bounded on Γ ?

Lattices and semi-length functions

Problem

If we know \tilde{b} is bounded on G , does it follow b is bounded on Γ ?

Burger–Monod: The answer is **YES!** if b is a quasi-cocycle, because the L^2 -induction $H_b^2(\Gamma, \mathcal{H}) \rightarrow H_{cb}^2(G, L^2(X, \mathcal{H}))$ is injective.

In general,

let $C := \sup \|b(gh) - (b(g) + \pi(g)b(h))\|$ and $\ell(g) := \|b(g)\| + C$.

Then, ℓ is a semi-length function: $\ell(gh) \leq \ell(g) + \ell(h)$.

The induced semi-length function $L: G \rightarrow \mathbb{R}_{\geq 0}$ is given by

$$L(g) = \int_X \ell(\beta(x, g)) \, dx.$$

The above problem generalizes to

Problem

If we know L is bounded on G , does it follow ℓ is bounded on Γ ?

Semi-length functions and nonlinear cohomology?

Theorem

Let $G \curvearrowright X$ be a probability measure preserving action,
and $\ell: X \times G \rightarrow \mathbb{R}_{\geq 0}$ be a groupoid semi-length function:

$$\ell(x, gh) \leq \ell(x, g) + \ell(g^{-1}x, h) \text{ a.e.}$$

If $\operatorname{ess-sup}_{g \in G} \int_X \ell(x, g) dx < +\infty$, then $\exists h \in L^1(X)$ such that

$$\ell(x, g) \leq h(x) + h(g^{-1}x) \text{ a.e.}$$

This theorem acts for the injectivity of $H_b^2(\Gamma, \mathcal{H}) \rightarrow H_{cb}^2(G, L^2(X, \mathcal{H}))$.

Corollary

Let $\ell: \Gamma \rightarrow \mathbb{R}_{\geq 0}$ be a semi-length function and $L: G \rightarrow \mathbb{R}_{\geq 0}$ be the induced semi-length function. If L is bounded, then so is ℓ .

In particular, property (TTT) passes to a cocompact lattice.

Semi-length functions and nonlinear cohomology?

Theorem

Let $G \curvearrowright X$ be a probability measure preserving action,
and $\ell: X \times G \rightarrow \mathbb{R}_{\geq 0}$ be a groupoid semi-length function:

$$\ell(x, gh) \leq \ell(x, g) + \ell(g^{-1}x, h) \text{ a.e.}$$

If $\operatorname{ess-sup}_{g \in G} \int_X \ell(x, g) dx < +\infty$, then $\exists h \in L^1(X)$ such that

$$\ell(x, g) \leq h(x) + h(g^{-1}x) \text{ a.e.}$$

This theorem acts for the injectivity of $H_b^2(\Gamma, \mathcal{H}) \rightarrow H_{cb}^2(G, L^2(X, \mathcal{H}))$.

Corollary

Let $\ell: \Gamma \rightarrow \mathbb{R}_{\geq 0}$ be a semi-length function and $L: G \rightarrow \mathbb{R}_{\geq 0}$ be the induced semi-length function. If L is bounded, then so is ℓ .

In particular, property (TTT) passes to a cocompact lattice.