

The Action of the Steenrod Algebra on the Cohomology of p -Compact Groups

By

Yutaka HEMMI*, Hirokazu NISHINOBU** and Kuniyuki OGI***

Abstract

In the present paper, we determine the action of the Steenrod operations on the cohomology of simply connected p -compact groups with no p -torsion in the integral homologies for an odd prime p . To do so we study simple simply connected p -compact groups since any simply connected p -compact group with no p -torsion in the integral homology is a product of such p -compact groups.

§1. Introduction

A p -compact group is a space X with finite mod p cohomology $H^*(X; \mathbb{F}_p)$ together with a homotopy equivalence $X \rightarrow \Omega(BX)$ to the loop space of a path connected p -complete space BX which is called the classifying space of X ([9]). In the present paper, we determine the action of the Steenrod operations on the cohomology of simply connected p -torsion free p -compact groups for an odd prime p . Here, by “ p -torsion free” we mean “to have no p -torsion in the integral homology”.

Typical examples of p -compact groups are the p -completion of connected compact Lie groups. For simply connected compact Lie groups, it is well known

Communicated by S. Mukai. Received November 27, 2007. Revised May 7, 2008.

2000 Mathematics Subject Classification(s): Primary 55S05; Secondary 57T25, 55P35, 55P60.

Key words: p -compact group, cohomology operation, simple p -compact group.

*Department of Mathematics, Faculty of Science, Kochi University, Kochi 780–8520, Japan.

e-mail: hemmi@math.kochi-u.ac.jp

**Department of Mathematics, Faculty of Science, Kochi University, Kochi 780–8520, Japan.

e-mail: nishinobu@math.kochi-u.ac.jp

***Anritsu Engineering Co., Ltd., 5–1–1 Onna, Atsugi-City, Kanagawa 243–0032, Japan.

e-mail: kuniyuki.Ogi@anritsu.com

how the Steenrod operations act on their cohomologies. In particular, if X is one of the p -completion of the Lie groups $SU(n)$, $Spin(2n+1)$, $Sp(n)$, and p -torsion free exceptional Lie groups for an odd prime p , then the following property holds:

($\mathcal{P}$) X has primitive mod p cohomology generators x_i ($1 \leq i \leq k$) with

$$H^*(X; \mathbb{F}_p) \cong \Lambda(x_1, \dots, x_k), \quad \deg x_i = 2n_i - 1 \quad (n_1 \leq \dots \leq n_k)$$

such that if $n_j - n_i = t(p-1)$ with $t > 0$, then

$$\mathcal{P}^t x_i = \binom{n_i - 1}{t} x_j.$$

On the other hand, ($\mathcal{P}$) is not necessarily true for $X = Spin(2n)_p^\wedge$, where $Y_p^\wedge$ denotes the p -completion of a space Y . However, $Spin(2n)_p^\wedge$ splits as

$$Spin(2n)_p^\wedge \simeq Spin(2n-1)_p^\wedge \times (S^{2n-1})_p^\wedge,$$

and ($\mathcal{P}$) holds for $Spin(2n-1)_p^\wedge$ and for an H -space $(S^{2n-1})_p^\wedge$. Thus, we can say that $Spin(2n)_p^\wedge$ splits as a product of H -spaces satisfying ($\mathcal{P}$).

Our first purpose is to show that the similar result holds for simple simply connected p -torsion free p -compact groups for any odd prime p . It is proved that if p is an odd prime, then a simply connected p -compact group X is simple if and only if its classifying space BX realizes a p -adic pseudoreflexion group in the list given by Clark and Ewing ([7], [8]). We will show that ($\mathcal{P}$) holds for almost all simple simply connected p -torsion free p -compact groups. Our proof uses the Clark-Ewing list. Only the exceptions are the loop spaces of generalized Grassmanians.

The generalized Grassmanian $BX(m, q; n)$ is a p -complete space defined for any positive integers m, q and n with $q|m$, $m|p-1$, $m > 1$ and $n > 1$. The space $BX(m, q; n)$, which realizes the p -adic pseudoreflexion group of No. 2a in the Clark-Ewing list, satisfies

$$\begin{aligned} H^*(BX(m, q; n); \mathbb{F}_p) &\cong \mathbb{F}_p[u_1, \dots, u_{n-1}, v], \\ \deg u_i &= 2im, \quad \deg v = 2nq. \end{aligned}$$

Then the p -compact group $X(m, q; n) = \Omega BX(m, q; n)$ is a simple simply connected p -compact group with

$$\begin{aligned} H^*(X(m, q; n); \mathbb{F}_p) &\cong \Lambda(x_1, \dots, x_{n-1}, y) \\ \deg x_i &= 2im - 1, \quad \deg y = 2nq - 1. \end{aligned}$$

Note that

$$Spin(2n)_p^\wedge \simeq X(2, 1; n) \quad \text{and} \quad Spin(2n+1)_p^\wedge \simeq X(2, 2; n).$$

Then, we show the following¹

Theorem 1.1. *If p is an odd prime, then any simple simply connected p -torsion free p -compact group other than $X(m, q; n)$ with $q < m$ satisfies $(\mathcal{P})$.*

A complete list for the Steenrod action on the cohomology of simple simply connected p -torsion free p -compact group is given in section 4. We note that $\mathcal{P}^i$ vanishes for $i \geq p$ except for the ones of Nos. 1 and 2a for dimensional reasons.

Now it is proved by Adams and Wilkerson [1, Corollary 1.3] and Dwyer, Miller and Wilkerson [8, Theorem 1.1] that if p is an odd prime, then any simply connected p -torsion free p -compact group splits as a product of simple p -compact groups. Thus, from the above theorem we have the following

Theorem 1.2. *Let X be a simply connected p -torsion free p -compact group for an odd prime p . Then X splits as*

$$X \simeq X_1 \times \cdots \times X_m,$$

where each X_i is a p -compact group satisfying $(\mathcal{P})$ or the p -completion of an odd sphere.

For the case of $p = 2$, Thomas [21] proved a similar fact to $(\mathcal{P})$ for 2-torsion free simply connected H -spaces with primitively generated mod 2 cohomology. Moreover, his result includes a stronger consequence. In fact, he showed that if x is a primitive mod 2 cohomology class with $\deg x = 2n - 1$, then for any $t > 0$ with $\binom{n-t-1}{t} \not\equiv 0 \pmod{2}$, there exists a primitive class y with $\deg y = 2(n-t) - 1$ such that $Sq^{2t}(y) = x$, which is not generally true if p is an odd prime.

Since the above theorem determines the action of the Steenrod algebra on the cohomology of all the simple simply connected p -compact groups, we shall have many applications. For example, we have the following

Corollary 1.3. *Let X be a simply connected p -compact group. Suppose that the action of the Steenrod algebra on $H^*(X; \mathbb{F}_p)$ is trivial. If $p \geq 5$, then X is p -regular, that is, X is p -equivalent to a product of odd spheres; if $p = 3$, then X is 3-equivalent to a product of odd spheres and some copies of the exceptional Lie group G_2 .*

¹After the final submission, the authors found that Theorem 1.1. was independently proved by preprint; D. Davis; Homotopy type and v_1 -periodic homotopy groups of p -compact groups; preprint

In the above corollary we need not to assume that X is p -torsion free.

The authors would like to thank Yusuke Kawamoto for his useful suggestions.

§2. Simple p -Compact Groups

From now on, we assume that p is an odd prime. Moreover, any p -compact group is assumed to be simply connected and p -torsion free.

Let B be a p -complete space such that the mod p cohomology $H^*(B; \mathbb{F}_p)$ is a finitely generated polynomial algebra:

$$H^*(B; \mathbb{F}_p) \cong \mathbb{F}_p[u_1, \dots, u_k], \quad \deg u_i = 2n_i.$$

Then, by Dwyer, Miller and Wilkerson [8, Theorem 1.1], there exists a p -adic pseudoreflection group $W \subset GL(k, \mathbb{Z}_p^\wedge)$ and a map $f: BT \rightarrow B$ equivariant up to homotopy with respect to the natural action of W on BT and the trivial action of W on B such that f induces an isomorphism $H^*(B; \mathbb{F}_p) \cong H^*(BT; \mathbb{F}_p)^W$, where $\mathbb{Z}_p^\wedge$ is the ring of p -adic integers, and BT is the classifying space of the p -completion of the k -dimensional torus T^k . The space B is called a realization of W . We notice that the order of W equals to the product $n_1 \dots n_k$. If the order of W is divisible by p , then W and $\mathbb{F}_p[u_1, \dots, u_k]$ are called modular.

All irreducible p -adic pseudoreflection groups are classified by Clark and Ewing [7]. Their result is based on the classification of the irreducible complex reflection groups by Shephard and Todd [20]. A p -compact group X is simple if its classifying space BX realizes one of the irreducible p -adic pseudoreflection groups in the Clark-Ewing list.

Then Theorem 1.2 and Corollary 1.3 are proved from Theorem 1.1.

Proof of Theorem 1.2 from Theorem 1.1. Let X be a p -compact group. According to Adams and Wilkerson [1, Corollary 1.3] and Dwyer, Miller and Wilkerson [8, Theorem 1.1], X splits as the product of some simple p -compact groups. Moreover, it is proved by [5, Corollary 1.4] that if $q < m$, then

$$X(m, q; n) \simeq_p X(m, m; n-1) \times (S^{2nq-1})_p^\wedge.$$

Thus, X splits as the product of simple p -compact groups other than $X(m, q; n)$ with $q < m$ and the p -completion of odd spheres. Thus we have the result by Theorem 1.1. $\square$

Proof of Corollary 1.3. First we note that the integral homology of X has no p -torsion. In fact, if it does, then the action of the Bockstein operation

on $H^*(X; \mathbb{F}_p)$ is non trivial ([13]). Thus, we have only to show the statement for the case that X is simple by [1, Corollary 1.3] and [8, Theorem 1.1]. Then, by using Theorem 1.1, we see easily that X is either of No. 2b of type (2, 6) for $p = 3$ or of type $(n_1, \dots, n_k)$ with $n_k - n_1 < p - 1$. For the former case X is 3-equivalent to G_2 , while for the later case X is p -regular by [12]. $\square$

§3. Proof of the Main Theorem

Let X be a p -compact group. Then the mod p cohomology of X is an exterior algebra:

$$(3.1) \quad H^*(X; \mathbb{F}_p) \cong \Lambda(x_1, \dots, x_k)$$

for some primitive classes x_i . The sequence $(n_1, \dots, n_k)$, where $\deg x_i = 2n_i - 1$, is called the type of X . We note that the type of X is the same as the one of the polynomial algebra $H^*(BX; \mathbb{F}_p)$, where the type of a polynomial algebra $\mathbb{F}_p[u_1, \dots, u_k]$ is a sequence $(n_1, \dots, n_k)$ with $\deg u_i = 2n_i$ ($1 \leq i \leq k$).

Now we show the following

Lemma 3.1. *Let X be a p -compact group of type $(n_1, \dots, n_k)$. Let n be a positive integer with $n \not\equiv 0 \pmod{p}$ and $n > p$. Suppose that $\mathcal{P}^1: PH^{2tn-2(p-1)-1}(X; \mathbb{F}_p) \rightarrow PH^{2tn-1}(X; \mathbb{F}_p)$ is a monomorphism for $2 \leq t \leq p$, where $PH^*(X; \mathbb{F}_p)$ is the primitive module of $H^*(X; \mathbb{F}_p)$. Then $\mathcal{P}^1: PH^{2n-2(p-1)-1}(X; \mathbb{F}_p) \rightarrow PH^{2n-1}(X; \mathbb{F}_p)$ is an epimorphism.*

Proof. By the assumption, we have $H^*(BX; \mathbb{F}_p) \cong \mathbb{F}_p[u_1, \dots, u_k]$ for some u_i ($1 \leq i \leq k$) with $\deg u_i = 2n_i$. Here, by the same reason as [11, Lemma 4.2] we can choose u_i to satisfy the following condition:

If $\mathcal{P}^1(u_i)$ is not decomposable then $\mathcal{P}^1(u_i) = u_j$ for some u_j ; if $\mathcal{P}^1(u_i) = \mathcal{P}^1(u_j) = u_k$ then $u_i = u_j$.

Then, the assumption that $\mathcal{P}^1: PH^{2tn-2(p-1)-1}(X; \mathbb{F}_p) \rightarrow PH^{2tn-1}(X; \mathbb{F}_p)$ is a monomorphism is equivalent to say that if $\deg u_s = 2tn - 2(p-1)$ then $\mathcal{P}^1 u_s = u_m$ for some m . Then we show that if $\deg u_j = 2n$ then $\mathcal{P}^1(u_i) = u_j$ for some u_i , which implies the result clearly.

Let J be the ideal of $H^*(BX; \mathbb{F}_p)$ generated by all the generators u_s with $s \neq j$ so that $H^*(BX; \mathbb{F}_p)/J \cong \mathbb{F}_p[u_j]$. Since $n \not\equiv 0 \pmod{p}$, we have

$$u_j^p = \mathcal{P}^n u_j = \mathcal{P}^1(n^{-1} \mathcal{P}^{n-1} u_j).$$

Thus we have $\mathcal{P}^1(H^*(BX; \mathbb{F}_p)) \not\subset J$, and so there is a generator u_i with $\mathcal{P}^1 u_i \notin J$. For dimensional reasons, we have $\deg u_i = 2tn - 2(p-1)$ for some t with

$1 \leq t \leq p$. Then, by the assumption, t must be 1 and $\mathcal{P}^1 u_i = u_j$ since $\mathcal{P}^1 u_i$ is not decomposable. $\square$

Next we consider some special cases.

Proposition 3.2. *Let X be a p -compact group of type $(n_1, \dots, n_{p-1})$ with*

$$n_i = \begin{cases} i(p-1), & 1 \leq i \leq p-2 \\ p(p-1), & i = p-1 \end{cases}$$

Then, X satisfies $(\mathcal{P})$.

Proof. We show that we can choose primitive generators x_i ($1 \leq i \leq p-1$) of $H^*(X; \mathbb{F}_p)$ such that

$$(3.2) \quad x_i = \binom{p-2}{i-1}^{-1} \mathcal{P}^{i-1} x_1 \quad \text{for } 2 \leq i \leq p-2.$$

Then $(\mathcal{P})$ is satisfied. In fact, if $1 \leq i < j \leq p-2$, then $n_j - n_i = (j-i)(p-1)$, and we have

$$\begin{aligned} \mathcal{P}^{j-i} x_i &= \binom{p-2}{i-1}^{-1} \mathcal{P}^{j-i} \mathcal{P}^{i-1} x_1 \\ &= \binom{p-2}{i-1}^{-1} \binom{j-1}{i-1} \mathcal{P}^{j-1} x_1 \\ &= \binom{p-i-1}{j-i} \binom{p-2}{j-1}^{-1} \mathcal{P}^{j-1} x_1 \\ &= \binom{n_i-1}{j-i} x_j \end{aligned}$$

since $n_i - 1 = (i-1)p + (p-i-1)$ and $j-i < p$. On the other hand, $n_{p-1} - n_i = (p-i)(p-1)$ for $i \leq p-2$, and

$$\mathcal{P}^{p-i} x_i = (p-i)^{-1} \mathcal{P}^1 \mathcal{P}^{p-i-1} x_i = 0 = \binom{n_i-1}{p-i} x_{p-1}.$$

Now we show (3.2). By the assumption we have

$$H^*(BX; \mathbb{F}_p) \cong \mathbb{F}_p[u_1, u_2, \dots, u_{p-1}]$$

with

$$\deg u_i = \begin{cases} 2i(p-1), & 1 \leq i \leq p-2 \\ 2p(p-1), & i = p-1. \end{cases}$$

First, since $\mathcal{P}^{i-1}u_{p-i} \in D^2H^*(BX; \mathbb{F}_p)$ for dimensional reasons, we have

$$(3.3) \quad \mathcal{P}^i u_{p-i} = i^{-1} \mathcal{P}^1 \mathcal{P}^{i-1} u_{p-i} \in D^2H^*(BX; \mathbb{F}_p),$$

where D^tH^* is the module of t -fold decomposables of an algebra H^* .

Let J be an ideal of $H^*(BX; \mathbb{F}_p)$ generated by $u_1, \dots, u_{p-2}$. Then, we show that

$$(3.4) \quad \mathcal{P}^p u_{p-1} \notin J.$$

In fact, if $\mathcal{P}^p u_{p-1} \in J$, then for dimensional reasons, $\mathcal{P}^p u_i \in J$ for $1 \leq i \leq p-1$. Thus

$$\mathcal{P}^1(H^*(BX; \mathbb{F}_p)) \subset J \quad \text{and} \quad \mathcal{P}^p(H^*(BX; \mathbb{F}_p)) \subset J.$$

Then

$$u_{p-1}^p = \mathcal{P}^{p(p-1)} u_{p-1} = -\mathcal{P}^1 \mathcal{P}^{p(p-1)-2} \mathcal{P}^1 u_{p-1} - \mathcal{P}^p \mathcal{P}^{p(p-1)} u_{p-1} \in J,$$

and we have a contradiction.

Now we consider the BP cohomology of BX at the prime p (cf. [10, Lemma 8.1]):

$$BP^*(BX) \cong BP^*[[U_1, \dots, U_{p-1}]],$$

where $T(U_i) = u_i$ for the natural map $T: BP^*(BX) \rightarrow H^*(BX; \mathbb{F}_p)$. We notice that the kernel of T is the ideal $(p, v_1, v_2, \dots)$ generated by $p, v_1, v_2, \dots$.

Let r_t be the Landweber-Novikov operation so that

$$\chi(\mathcal{P}^t)(T(z)) = T(r_t(z))$$

for any $z \in BP^*(BX)$, where χ is the canonical anti-automorphism on the mod p Steenrod algebra. In particular, since $\chi(\mathcal{P}^t) = (-1)^t \mathcal{P}^t$ for $t \leq p$ we have

$$(-1)^t \mathcal{P}^t(T(z)) = T(r_t(z)) \quad \text{for } t \leq p.$$

Since $\mathcal{P}^p u_{p-1} \notin J$ by (3.4), we have

$$(3.5) \quad r_p U_{p-1} \equiv a U_{p-1}^2 \pmod{D^3 BP^*(BX) + (p, v_1, v_2, \dots)}$$

for some $a \in \mathbb{Z}_{(p)}$ with $a \not\equiv 0 \pmod{p}$, where $\mathbb{Z}_{(p)}$ is the ring of p -local integers.

Now we can put

$$\begin{aligned} r_1 U_{p-1} &\equiv b_1 U_1 U_{p-1} + \sum_{k=3}^{(p+1)/2} b_k U_k U_{p+1-k} \\ &\quad + v_1 \left(c_2 U_2 U_{p-1} + \sum_{k=4}^{(p+1)/2} c_k U_k U_{p+2-k} \right) \\ &\pmod{D^3 BP^*(BX) + (p, v_1, v_2, \dots)^2} \end{aligned}$$

with $b_k, c_k \in \mathbb{Z}_{(p)}$. Here, $r_{p-1}r_1 \equiv pr_p \pmod{(v_1, v_2, \dots)}$, and $r_i U_{p-i} \in D^2BP^*(BX) + (p, v_1, \dots)$ by (3.3). Thus

$$\begin{aligned} pr_p U_{p-1} &\equiv r_{p-1}r_1 U_{p-1} \\ &\equiv b_1(r_{p-1}U_1)U_{p-1} + \sum_{k=3}^{(p+1)/2} b_k(r_{p-k}U_k)(r_{k-1}U_{p+1-k}) \\ &\quad + p \left(c_2(r_{p-2}U_2)U_{p-1} + \sum_{k=4}^{(p+1)/2} c_k(r_{p-k}U_k)(r_{k-2}U_{p+2-k}) \right) \\ &\equiv pb_1 d U_{p-1}^2 \\ &\quad \pmod{D^3BP^*(BX) + (p^2, v_1, v_2, \dots)}, \end{aligned}$$

where $r_{p-1}U_1 \equiv pd U_{p-1} \pmod{D^2BP^*(BX) + (p^2, v_1, v_2, \dots)}$ with $d \in \mathbb{Z}_{(p)}$. Thus we have $b_1 d \equiv a \not\equiv 0 \pmod{p}$ by (3.5). In particular, we have

$$r_{p-1}U_1 \not\equiv 0 \pmod{D^2BP^*(BX) + (p^2, v_1, v_2, \dots)}.$$

Suppose that $\mathcal{P}^{p-3}u_1 \in D^2H^*(BX; \mathbb{F}_p)$. Then

$$r_{p-3}U_1 \equiv pe U_{p-2} \pmod{D^2BP^*(BX) + (p^2, v_1, v_2, \dots)}$$

for some $e \in \mathbb{Z}_{(p)}$. Thus, we have

$$\begin{aligned} r_{p-1}U_1 &\equiv \binom{p-1}{2}^{-1} r_2 r_{p-3}U_1 \equiv pe r_2 U_{p-2} \equiv 0 \\ &\quad \pmod{D^2BP^*(BX) + (p^2, v_1, v_2, \dots)}, \end{aligned}$$

which is a contradiction. Therefore, we have

$$\mathcal{P}^{p-3}u_1 \notin D^2H^*(BX; \mathbb{F}_p),$$

and we can choose generators u_i for $2 \leq i \leq p-2$ such that

$$u_i = \binom{p-2}{i-1}^{-1} \mathcal{P}^{i-1}u_1.$$

Thus by putting $x_i = s(u_i)$, where s is the cohomology suspension, we have (3.2), which completes the proof. $\square$

Next we consider the p -compact group $X(m, m; n)$.

Proposition 3.3. *For any positive integers m and n with $m|p-1$, the p -compact group $X(m, m; n)$ satisfies $(\mathcal{P})$.*

Proof. According to Castellana [6, p.2817] there is a map $Bc: BX(q, 1; n) \rightarrow BU(nq)_p^\wedge$ such that if we put

$$u_j = Bc^*(c_{jq}) \quad (1 \leq j \leq n)$$

then

$$H^*(BX(m, m; n); \mathbb{F}_p) = \mathbb{F}_p[u_1, \dots, u_n],$$

where c_j is the j -th Chern class so that

$$H^*(BU(nq)_p^\wedge; \mathbb{F}_p) = \mathbb{F}_p[c_1, \dots, c_{nq}].$$

Then by putting $x_i = s(u_i)$ for the cohomology suspension s we have $(\mathcal{P})$ for $X(m, m; n)$ from the one for $U(nq)_p^\wedge$ ([4, Corollaire 11.4], [16]). □

Now we show Theorem 1.1.

Proof of Theorem 1.1. Let X be a simple p -compact group of type $(n_1, \dots, n_k)$ other than $X(m, q; n)$ with $q < m$.

If $n_k - n_1 < p - 1$, then $(\mathcal{P})$ clearly holds since X is p -regular and there are no i and j with $n_j - n_i = t(p - 1)$ for any $t > 0$.

Suppose that $p - 1 \leq n_k - n_1 < 2(p - 1)$. Then, by checking the Clark-Ewing list, we can see that $n_i \neq n_j$ for $i \neq j$.

Let $n_j - n_i = t(p - 1)$ for some $t > 0$. Then it is clear that $t = 1$. Moreover, by checking the Clark-Ewing list, we can see that $n_j \not\equiv 0 \pmod p$ and $PH^{2sn_j - 2(p-1)-1}(X; \mathbb{F}_p) = 0$ for $2 \leq s \leq p$. Thus by Lemma 3.1, $\mathcal{P}^1: PH^{2n_i-1}(X; \mathbb{F}_p) \rightarrow PH^{2n_j-1}(X; \mathbb{F}_p)$ is an epimorphism. Since $n_i - 1 \equiv n_j \not\equiv 0 \pmod p$, we can choose generators x_i and x_j with

$$\mathcal{P}^1 x_i = \binom{n_i - 1}{1} x_j$$

Thus $(\mathcal{P})$ holds.

Now we consider the remaining case $n_k - n_1 \geq 2(p - 1)$. The Clark-Ewing list consists of thirty seven families of irreducible p -adic pseudoreflection groups, with two subfamilies 2a and 2b in No. 2. It is proved that all non-modular irreducible p -adic pseudoreflection groups have realizations ([7]). On the other hand, not all modular irreducible p -adic pseudoreflection groups have realizations. The non-realizable ones for an odd prime p are of Nos. 28, 35, 36 for $p = 3$, and 37 for $p = 3, 5$ ([22], [2], [18], [19]). The following is the table of

all realizable irreducible p -adic pseudoreflection groups with $n_k - n_1 \geq 2(p-1)$.

No.	type	prime
1	$(2, 3, \dots, n+1)$	$p \leq (n+1)/2$
2a	$(m, 2m, \dots, nm)$	$p \leq (n-1)m/2 + 1$
2b	$(2, 6)$	3
26	$(6, 12, 18)$	7
28	$(2, 6, 8, 12)$	5
29	$(4, 8, 12, 20)$	5
30	$(2, 12, 20, 30)$	11
31	$(8, 12, 20, 24)$	5
32	$(12, 18, 24, 30)$	7
33	$(4, 6, 10, 12, 18)$	7
34	$(6, 12, 18, 24, 30, 42)$	7, 13, 19
35	$(2, 5, 6, 8, 9, 12)$	5
36	$(2, 6, 8, 10, 12, 14, 18)$	5, 7
37	$(2, 8, 12, 14, 18, 20, 24, 30)$	7, 11, 13

Now, the p -adic pseudoreflection groups of Nos. 1, 2b, 28, 35, 36 and 37 are the Weyl groups of the Lie groups $SU(n+1)$, G_2 , F_4 , E_6 , E_7 and E_8 , respectively. If BX realizes the Weyl group of a Lie group G , then $H^*(X; \mathbb{F}_p)$ is isomorphic to $H^*(G; \mathbb{F}_p)$ as an algebra over the Steenrod algebra. Thus, for these cases, we have only to show $(\mathcal{P})$ for corresponding Lie groups, which are proved in [4, Corollaire 11.4] (see also [16]), [3], [12] and [15].

For No. 2a, we have already done in Propositions 3.3.

For Nos. 29 for $p = 5$ and 34 for $p = 7$, $(\mathcal{P})$ holds by Proposition 3.2. For the Nos. 30 for $p = 11$ and 31 for $p = 5$, we can apply Lemma 3.1 to show $(\mathcal{P})$. Note that for No. 31, we have

$$\mathcal{P}^2 x_2 = 2^{-1} \mathcal{P}^1 \mathcal{P}^1 x_2 = 0 = \begin{pmatrix} 12-1 \\ 2 \end{pmatrix} x_3.$$

For No. 34 for $p = 19$, $(\mathcal{P})$ holds by Nishinobu [17, Theorem 1.2].

Now we consider No. 32 for $p = 7$. Then by Lemma 3.1 generators x_i are chosen to satisfy

$$\mathcal{P}^1 x_2 = \begin{pmatrix} 18-1 \\ 1 \end{pmatrix} x_3, \quad \mathcal{P}^1 x_3 = \begin{pmatrix} 24-1 \\ 1 \end{pmatrix} x_4.$$

Thus, we have only to show that

$$\mathcal{P}^1 x_1 \neq 0.$$

Then by modifying x_1 we have

$$\mathcal{P}^1 x_1 = \begin{pmatrix} 12 & -1 \\ & 1 \end{pmatrix} x_2.$$

Suppose contrarily that $\mathcal{P}^1 x_1 = 0$. Then, in $H^*(BX; \mathbb{F}_7) = \mathbb{F}_7[u_1, u_2, u_3, u_4]$ the generators u_i can be chosen to satisfy

$$\mathcal{P}^1 u_1 = 0, \quad \mathcal{P}^1 u_2 = \begin{pmatrix} 18 & -1 \\ & 1 \end{pmatrix} u_3, \quad \mathcal{P}^1 u_3 = \begin{pmatrix} 24 & -1 \\ & 1 \end{pmatrix} u_4.$$

We show that $\mathcal{P}^1 u_4$ is not in the ideal (u_1, u_3, u_4) generated by u_1, u_3, u_4 . In fact, if $\mathcal{P}^1 u_4 \in (u_1, u_3, u_4)$, then $\mathcal{P}^1(H^*(BX; \mathbb{F}_7)) \subset (u_1, u_3, u_4)$ since $\mathcal{P}^1 u_i \in (u_1, u_3, u_4)$ for any $i = 1, 2, 3, 4$. On the other hand, we have

$$\mathcal{P}^1(\mathcal{P}^{17} u_2) = 18 \mathcal{P}^{18} u_2 = 18 u_2^7 \notin (u_1, u_3, u_4).$$

This is a contradiction, and so we have

$$\mathcal{P}^1 u_4 \notin (u_1, u_3, u_4).$$

Now, consider the quotient algebra $H^*(BX; \mathbb{F}_7)/(u_1) \cong \mathbb{F}_7[\bar{u}_2, \bar{u}_3, \bar{u}_4]$, where $\bar{u}_i$ is the coset of u_i . Note that this algebra does not necessarily admit the Steenrod action, but has the $\mathcal{P}^1$ action. Then, by the above argument we can put

$$\mathcal{P}^1 \bar{u}_4 = a \bar{u}_2^2$$

with $a \neq 0$. Then by iterating $\mathcal{P}^1$ we have

$$(\mathcal{P}^1)^7 \bar{u}_4 = 3a^3 \bar{u}_2^4 + a^2 \bar{u}_2 \bar{u}_3 \bar{u}_4 + 3a^2 \bar{u}_3^3 \neq 0.$$

This is a contradiction since $(\mathcal{P}^1)^7 = 0$. Thus $\mathcal{P}^1 u_1 \neq 0$ and we have $\mathcal{P}^1 x_1 \neq 0$.

The remaining cases are of Nos. 26 and 33 for $p = 7$, and 34 for $p = 13$. The idea of the proof for these cases are essentially the same. First we consider No. 26.

By Lemma 3.1 we can choose generators x_i such that

$$\mathcal{P}^1 x_2 = \begin{pmatrix} 12 & -1 \\ & 1 \end{pmatrix} x_3.$$

Thus, we prove

$$\mathcal{P}^1 x_1 \neq 0$$

to show

$$\mathcal{P}^1 x_1 = \begin{pmatrix} 6 & -1 \\ & 1 \end{pmatrix} x_2$$

for a suitable x_1 .

Suppose contrarily that $\mathcal{P}^1 x_1 = 0$. Then we can choose generators u_i ($i = 1, 2, 3$) of $H^*(BX; \mathbb{F}_p)$ such that the ideal (u_1) of $H^*(BX; \mathbb{F}_p)$ generated by u_1 is closed under the action of the Steenrod algebra for dimensional reasons. Thus $H^*(BX; \mathbb{F}_p)/(u_1)$ is a polynomial algebra of the type (12, 18) on which the Steenrod algebra acts unstably. According to Adams and Wilkerson [1, Theorem 1.2], the type of any non-modular polynomial algebra on which the Steenrod algebra acts unstably must be a union of types in the Clark-Ewing list, which is not the case. This is a contradiction, and $\mathcal{P}^1 u_1$ is indecomposable. Thus we have $\mathcal{P}^1 x_1 \neq 0$.

Next consider No. 33.

By Lemma 3.1 we can choose generators x_i such that

$$\mathcal{P}^1 x_1 = \begin{pmatrix} 4-1 \\ 1 \end{pmatrix} x_3, \quad \mathcal{P}^1 x_4 = \begin{pmatrix} 12-1 \\ 1 \end{pmatrix} x_5.$$

Thus we have only to show that

$$\mathcal{P}^1 x_2 \neq 0$$

to show

$$\mathcal{P}^1 x_2 = \begin{pmatrix} 6-1 \\ 1 \end{pmatrix} x_4$$

for a suitable x_2 .

Suppose contrarily that $\mathcal{P}^1 x_2 = 0$. Then we can choose generators u_i of $H^*(BX; \mathbb{F}_p)$ such that the ideal (u_1, u_2, u_3) of $H^*(BX; \mathbb{F}_p)$ generated by u_1, u_2, u_3 is closed under the action of the operation $\mathcal{P}^1$. Moreover, for dimensional reasons, we have $\mathcal{P}^7 u_3 \in (u_1, u_2, u_3)$. Thus (u_1, u_2, u_3) is closed under the action of the Steenrod algebra, and we have a polynomial algebra $H^*(BX; \mathbb{F}_p)/(u_1, u_2, u_3)$ of the type (12, 18) on which the Steenrod algebra acts unstably. Thus, by the same reason as the case of No. 26, we have a contradiction, and we have $\mathcal{P}^1 x_2 \neq 0$.

Finally, we consider No. 34.

By Lemma 3.1, we can choose generators x_i such that

$$\mathcal{P}^1 x_2 = \begin{pmatrix} 12-1 \\ 1 \end{pmatrix} x_4, \quad \mathcal{P}^1 x_3 = \begin{pmatrix} 18-1 \\ 1 \end{pmatrix} x_5, \quad \mathcal{P}^1 x_5 = \begin{pmatrix} 30-1 \\ 1 \end{pmatrix} x_6.$$

Thus, we have only to show that

$$\mathcal{P}^1 x_1 \neq 0$$

to show

$$\mathcal{P}^1 x_1 = \binom{6-1}{1} x_3$$

for a suitable x_1 .

Suppose contrarily that $\mathcal{P}^1 x_1 = 0$. Then we can choose generators u_i of $H^*(BX; \mathbb{F}_p)$ such that the ideal (u_1) of $H^*(BX; \mathbb{F}_p)$ generated by u_1 is closed under the action of the Steenrod algebra for dimensional reasons. Thus we have a polynomial algebra $H^*(BX; \mathbb{F}_p)/(u_1)$ of type $(12, 18, 24, 30, 42)$ on which the Steenrod algebra acts unstably. Thus, by the same reason as the case of No. 26, we have a contradiction, and we have $\mathcal{P}^1 x_1 \neq 0$. This completes the proof of the theorem. $\square$

§4. Steenrod Connection

In this section we list the complete Steenrod actions on the cohomology of simple simply connected p -torsion free p -compact groups for any odd prime p . In No. 2a we only consider $X(m, m; n)$. For $X(m, q; n)$ with $q < m$, the Steenrod actions can be given from the following isomorphism of algebras over the Steenrod algebra:

$$H^*(X(m, q; n); \mathbb{F}_p) \cong H^*(X(m, m; n-1); \mathbb{F}_p) \otimes H^*(S^{2nq-1}; \mathbb{F}_p).$$

In the following, we only list non trivial actions, where PH^n denotes the n -dimensional primitive module of $H^*(X; \mathbb{F}_p)$.

(No.1 of type: $(2, 3, \dots, n+1)$) For any odd prime p with $p \leq n$:

$$\mathcal{P}^k(PH^{2t-1}) = PH^{2(t+k(p-1))-1} \quad \text{for } \binom{t-1}{k} \not\equiv 0 \pmod{p},$$

where $t \geq 2$ and $t+k(p-1) \leq n+1$.

(No.2a of type: $(m, 2m, \dots, nm)$) For any odd prime p with $p \leq n$:

$$\mathcal{P}^k(PH^{2tm-1}) = PH^{2(tm+k(p-1))-1} \quad \text{for } \binom{tm-1}{k} \not\equiv 0 \pmod{p},$$

where $t \geq 1$ and $tm+k(p-1) \leq nm$.

(No.2b of type: $(2, p+1)$) For any odd prime p :

$$\mathcal{P}^1(PH^3) = PH^{2p+1}.$$

(No.5 of type: (6, 12)) For $p = 7$:

$$\mathcal{P}^1(PH^{11}) = PH^{23}.$$

(No.8 of type: (8, 12)) For $p = 5$:

$$\mathcal{P}^1(PH^{15}) = PH^{23}.$$

(No.9 of type: (8, 24)) For $p = 17$:

$$\mathcal{P}^1(PH^{15}) = PH^{47}.$$

(No.10 of type: (12, 24)) For $p = 13$:

$$\mathcal{P}^1(PH^{23}) = PH^{47}.$$

(No.12 of type: (6, 8)) For $p = 3$:

$$\mathcal{P}^1(PH^{11}) = PH^{15}.$$

(No.14 of type: (6, 24)) For $p = 19$:

$$\mathcal{P}^1(PH^{11}) = PH^{47}.$$

(No.16 of type: (20, 30)) For $p = 11$:

$$\mathcal{P}^1(PH^{39}) = PH^{59}.$$

(No.17 of type: (20, 60)) For $p = 41$:

$$\mathcal{P}^1(PH^{39}) = PH^{119}.$$

(No.18 of type: (30, 60)) For $p = 31$:

$$\mathcal{P}^1(PH^{59}) = PH^{119}.$$

(No.20 of type: (12, 30)) For $p = 19$:

$$\mathcal{P}^1(PH^{23}) = PH^{59}.$$

(No.24 of type: (4, 6, 14)) For $p = 11$:

$$\mathcal{P}^1(PH^7) = PH^{27}.$$

(No.25 of type: (6, 9, 12)) For $p = 7$:

$$\mathcal{P}^1(PH^{11}) = PH^{23}.$$

(No.26 of type: (6, 12, 18)) For $p = 7$:

$$\begin{aligned}\mathcal{P}^1(PH^{11}) &= PH^{23}, \\ \mathcal{P}^2(PH^{11}) &= \mathcal{P}^1(PH^{23}) = PH^{35}.\end{aligned}$$

For $p = 13$:

$$\mathcal{P}^1(PH^{11}) = PH^{35}.$$

(No.27 of type: (6, 12, 30)) For $p = 19$:

$$\mathcal{P}^1(PH^{23}) = PH^{59}.$$

(No.28 of type: (2, 6, 8, 12)) For $p = 5$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{11}, \\ \mathcal{P}^1(PH^{15}) &= PH^{23}.\end{aligned}$$

For $p = 7$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{15}, \\ \mathcal{P}^1(PH^{11}) &= PH^{23}.\end{aligned}$$

For $p = 11$:

$$\mathcal{P}^1(PH^3) = PH^{23}.$$

(No.29 of type: (4, 8, 12, 20)) For $p = 5$:

$$\begin{aligned}\mathcal{P}^1(PH^7) &= PH^{15}, \\ \mathcal{P}^2(PH^7) &= \mathcal{P}^1(PH^{15}) = PH^{23}.\end{aligned}$$

For $p = 13$:

$$\mathcal{P}^1(PH^{15}) = PH^{39}.$$

For $p = 17$:

$$\mathcal{P}^1(PH^7) = PH^{39}.$$

(No.30 of type: (2, 12, 20, 30)) For $p = 11$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{23}, \\ \mathcal{P}^1(PH^{39}) &= PH^{59}.\end{aligned}$$

For $p = 19$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{39}, \\ \mathcal{P}^1(PH^{23}) &= PH^{59}.\end{aligned}$$

For $p = 29$:

$$\mathcal{P}^1(PH^3) = PH^{59}.$$

(No.31 of type: $(8, 12, 20, 24)$) For $p = 5$:

$$\mathcal{P}^1(PH^{15}) = PH^{23},$$

$$\mathcal{P}^1(PH^{39}) = PH^{47}.$$

For $p = 13$:

$$\mathcal{P}^1(PH^{15}) = PH^{39},$$

$$\mathcal{P}^1(PH^{23}) = PH^{47}.$$

For $p = 17$:

$$\mathcal{P}^1(PH^{15}) = PH^{47}.$$

(No.32 of type: $(12, 18, 24, 30)$) For $p = 7$:

$$\mathcal{P}^1(PH^{23}) = PH^{35},$$

$$\mathcal{P}^2(PH^{23}) = \mathcal{P}^1(PH^{35}) = PH^{47},$$

$$\mathcal{P}^3(PH^{23}) = \mathcal{P}^2(PH^{35}) = \mathcal{P}^1(PH^{47}) = PH^{59}.$$

For $p = 13$:

$$\mathcal{P}^1(PH^{23}) = PH^{47},$$

$$\mathcal{P}^1(PH^{35}) = PH^{59}.$$

For $p = 19$:

$$\mathcal{P}^1(PH^{23}) = PH^{59}.$$

(No.33 of type: $(4, 6, 10, 12, 18)$) For $p = 7$:

$$\mathcal{P}^1(PH^7) = PH^{19},$$

$$\mathcal{P}^1(PH^{11}) = PH^{23},$$

$$\mathcal{P}^2(PH^{11}) = \mathcal{P}^1(PH^{23}) = PH^{35}.$$

For $p = 13$:

$$\mathcal{P}^1(PH^{11}) = PH^{35}.$$

(No.34 of type: $(6, 12, 18, 24, 30, 42)$) For $p = 7$:

$$\mathcal{P}^1(PH^{11}) = PH^{23},$$

$$\mathcal{P}^2(PH^{11}) = \mathcal{P}^1(PH^{23}) = PH^{35},$$

$$\mathcal{P}^3(PH^{11}) = \mathcal{P}^2(PH^{23}) = \mathcal{P}^1(PH^{35}) = PH^{47},$$

$$\mathcal{P}^4(PH^{11}) = \mathcal{P}^3(PH^{23}) = \mathcal{P}^2(PH^{35}) = \mathcal{P}^1(PH^{47}) = PH^{59}.$$

For $p = 13$:

$$\begin{aligned}\mathcal{P}^1(PH^{11}) &= PH^{35}, \\ \mathcal{P}^1(PH^{23}) &= PH^{47}, \\ \mathcal{P}^2(PH^{11}) &= \mathcal{P}^1(PH^{35}) = PH^{59}, \\ \mathcal{P}^3(PH^{11}) &= \mathcal{P}^2(PH^{35}) = \mathcal{P}^1(PH^{59}) = PH^{83}.\end{aligned}$$

For $p = 19$:

$$\begin{aligned}\mathcal{P}^1(PH^{11}) &= PH^{47}, \\ \mathcal{P}^1(PH^{23}) &= PH^{59}, \\ \mathcal{P}^2(PH^{11}) &= \mathcal{P}^1(PH^{47}) = PH^{83}.\end{aligned}$$

For $p = 31$:

$$\mathcal{P}^1(PH^{23}) = PH^{83}.$$

For $p = 37$:

$$\mathcal{P}^1(PH^{11}) = PH^{83}.$$

(No.35 of type: $(2, 5, 6, 8, 9, 12)$) For $p = 5$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{11}, \\ \mathcal{P}^1(PH^9) &= PH^{17}, \\ \mathcal{P}^1(PH^{15}) &= PH^{23}.\end{aligned}$$

For $p = 7$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{15}, \\ \mathcal{P}^1(PH^{11}) &= PH^{23}.\end{aligned}$$

For $p = 11$:

$$\mathcal{P}^1(PH^3) = PH^{23}.$$

(No.36 of type: $(2, 6, 8, 10, 12, 14, 18)$) For $p = 5$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{11}, \\ \mathcal{P}^1(PH^{15}) &= PH^{23}, \\ \mathcal{P}^1(PH^{19}) &= PH^{27}, \\ \mathcal{P}^2(PH^{19}) &= \mathcal{P}^1(PH^{27}) = PH^{35}.\end{aligned}$$

For $p = 7$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{15}, \\ \mathcal{P}^1(PH^{11}) &= PH^{23}, \\ \mathcal{P}^2(PH^{11}) &= \mathcal{P}^1(PH^{23}) = PH^{35}.\end{aligned}$$

For $p = 11$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{23}, \\ \mathcal{P}^1(PH^{15}) &= PH^{35}.\end{aligned}$$

For $p = 13$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{27}, \\ \mathcal{P}^1(PH^{11}) &= PH^{35}.\end{aligned}$$

For $p = 17$:

$$\mathcal{P}^1(PH^3) = PH^{35}.$$

(No.37 of type: (2, 8, 12, 14, 18, 20, 24, 30)) For $p = 7$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{15}, \\ \mathcal{P}^1(PH^{23}) &= PH^{35}, \\ \mathcal{P}^1(PH^{27}) &= PH^{39}, \\ \mathcal{P}^2(PH^{23}) &= \mathcal{P}^1(PH^{35}) = PH^{47}, \\ \mathcal{P}^3(PH^{23}) &= \mathcal{P}^2(PH^{35}) = \mathcal{P}^1(PH^{47}) = PH^{59}.\end{aligned}$$

For $p = 11$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{23}, \\ \mathcal{P}^1(PH^{15}) &= PH^{35}, \\ \mathcal{P}^1(PH^{27}) &= PH^{47}, \\ \mathcal{P}^1(PH^{39}) &= PH^{59}.\end{aligned}$$

For $p = 13$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{27}, \\ \mathcal{P}^1(PH^{15}) &= PH^{39}, \\ \mathcal{P}^1(PH^{23}) &= PH^{47}, \\ \mathcal{P}^1(PH^{35}) &= PH^{59}.\end{aligned}$$

For $p = 17$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{35}, \\ \mathcal{P}^1(PH^{15}) &= PH^{47}, \\ \mathcal{P}^1(PH^{27}) &= PH^{59}.\end{aligned}$$

For $p = 19$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{39}, \\ \mathcal{P}^1(PH^{23}) &= PH^{59}.\end{aligned}$$

For $p = 23$:

$$\begin{aligned}\mathcal{P}^1(PH^3) &= PH^{47}, \\ \mathcal{P}^1(PH^{15}) &= PH^{59}.\end{aligned}$$

For $p = 29$:

$$\mathcal{P}^1(PH^3) = PH^{59}.$$

References

- [1] J. F. Adams and C. W. Wilkerson, Finite H -spaces and algebras over the Steenrod algebra, *Ann. of Math. (2)* **111** (1980), no. 1, 95–143.
- [2] J. Aguadé, Constructing modular classifying spaces, *Israel J. Math.* **66** (1989), no. 1-3, 23–40.
- [3] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. of Math. (2)* **57** (1953), 115–207.
- [4] A. Borel and J.-P. Serre, Groupes de Lie et puissances réduites de Steenrod, *Amer. J. Math.* **75** (1953), 409–448.
- [5] N. Castellana, Quillen Grassmannians as non-modular homotopy fixed points, *Math. Z.* **248** (2004), no. 3, 477–493.
- [6] ———, On the p -compact groups corresponding to the p -adic reflection groups $G(q, r, n)$, *Trans. Amer. Math. Soc.* **358** (2006), no. 7, 2799–2819 (electronic).
- [7] A. Clark and J. Ewing, The realization of polynomial algebras as cohomology rings, *Pacific J. Math.* **50** (1974), 425–434.
- [8] W. G. Dwyer, H. R. Miller and C. W. Wilkerson, Homotopical uniqueness of classifying spaces, *Topology* **31** (1992), no. 1, 29–45.
- [9] W. G. Dwyer and C. W. Wilkerson, Homotopy fixed-point methods for Lie groups and finite loop spaces, *Ann. of Math. (2)* **139** (1994), no. 2, 395–442.
- [10] Y. Hemmi, On finite H -spaces given by sphere extensions of classical groups, *Hiroshima Math. J.* **14** (1984), no. 3, 451–470.
- [11] ———, Homotopy associative finite H -spaces and the mod 3 reduced power operations, *Publ. Res. Inst. Math. Sci.* **23** (1987), no. 6, 1071–1084.
- [12] P. G. Kumpel, Jr., Lie groups and products of spheres, *Proc. Amer. Math. Soc.* **16** (1965), 1350–1356.
- [13] J. P. Lin, Torsion in H -spaces. II, *Ann. Math. (2)* **107** (1978), no. 1, 41–88.

- [14] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Ann. of Math. Stud., 76, Princeton Univ. Press, Princeton, N.J., 1974.
- [15] M. Mimura and H. Toda, Cohomology operations and homotopy of compact Lie groups. I, *Topology* **9** (1970), 317–336.
- [16] S. Mukohda and S. Sawaki, On the $b_p^{k,j}$ coefficient of a certain symmetric function, *J. Fac. Sci. Niigata Univ. Ser. I* **1** (1954), no. 2, 6 pp.
- [17] H. Nishinobu, On the action of the Steenrod algebra on the cohomology of a certain p -compact group, *Kochi J. Math.* **2** (2007), 125–153.
- [18] D. Notbohm, For which pseudo-reflection groups are the p -adic polynomial invariants again a polynomial algebra?, *J. Algebra* **214** (1999), no. 2, 553–570.
- [19] ———, Erratum: “For which pseudo-reflection groups are the p -adic polynomial invariants again a polynomial algebra?” *J. Algebra* **218** (1999), no. 1, 286–287.
- [20] G. C. Shephard and J. A. Todd, Finite unitary reflection groups, *Canadian J. Math.* **6** (1954), 274–304.
- [21] E. Thomas, Steenrod squares and H -spaces. II, *Ann. of Math. (2)* **81** (1965), 473–495.
- [22] A. Zabrodsky, On the realization of invariant subgroups of $\pi_*(X)$, *Trans. Amer. Math. Soc.* **285** (1984), no. 2, 467–496.