

An Approximate Statement of the Main Theorem of Inter-universal Teichmüller Theory

Yuichiro Hoshi

RIMS, Kyoto University

December 7, 2015

Notation and Terminology

$$\begin{aligned} \mathcal{O}^\mu &\stackrel{\text{def}}{=} (\mathcal{O}^\times)_{\text{tor}} \subseteq \mathcal{O}^\times \stackrel{\text{def}}{=} \{|z| = 1\} \\ &\subseteq \mathcal{O}^\triangleright \stackrel{\text{def}}{=} \{0 < |z| \leq 1\} \subseteq \mathcal{O} \stackrel{\text{def}}{=} \{|z| \leq 1\} \end{aligned}$$

$$\mathcal{O}^{\times\mu} \stackrel{\text{def}}{=} \mathcal{O}^\times / \mathcal{O}^\mu$$

an *isomorph* of $A \stackrel{\text{def}}{\Leftrightarrow}$ an object which is isomorphic to A

R_+ : the underlying additive module of a ring R

F : a number field, i.e., $[F : \mathbb{Q}] < \infty$, s.t. $\sqrt{-1} \in F$

$\mathbb{V}(-)$: the set of primes of $(-)$

E : an elliptic curve over F which has

either good or split multiplicative reduction at $\forall v \in \mathbb{V}(F)$

$q_v \in \mathcal{O}_{F_v}^\times$: the q -parameter of E at $v \in \mathbb{V}(F)$

$q_E \stackrel{\text{def}}{=} (q_v)_{v \in \mathbb{V}(F)} \in \prod_{v \in \mathbb{V}(F)} \mathcal{O}_{F_v}^\times$

$\Rightarrow \deg q_E (= [F : \mathbb{Q}]^{-1} \log(\prod_{v \in \mathbb{V}(F)} \#(\mathcal{O}_{F_v}/q_v \mathcal{O}_{F_v}))) \quad (\approx 6 \cdot \text{ht}_E)$

The Szpiro Conjecture for Elliptic Curves over Number Fields

A certain upper bound of ht_E , i.e., $\deg q_E$

Suppose that the following $(*)$ holds:

$$(*): \exists N \geq 2, \exists C \geq 0 \quad \text{s.t.} \quad \deg q_E^N \leq \deg q_E + C$$

Then since $\deg q_E^N = N \cdot \deg q_E$, one may conclude that

$$\deg q_E \leq \frac{C}{N-1}.$$

In order to establish $(*)$, let us

- take two isomorphisms ${}^{\dagger}\mathfrak{S}, {}^{\ddagger}\mathfrak{S}$ of (a part of) scheme theory,
- consider a “link” between these two isomorphisms

$$\Theta_{\text{naive}}: {}^{\dagger}\mathfrak{S} \ni {}^{\dagger}q_E^N \mapsto {}^{\ddagger}q_E \in {}^{\ddagger}\mathfrak{S}, \quad \text{and}$$

- compare, via Θ_{naive} , the computation of $\deg$ of ${}^{\dagger}q_E^N$ (in ${}^{\dagger}\mathfrak{S}$) with the computation of $\deg$ of ${}^{\ddagger}q_E$ (in ${}^{\ddagger}\mathfrak{S}$).

$$\Theta_{\text{naive}}: {}^{\dagger}\mathfrak{S} \rightarrow {}^{\ddagger}\mathfrak{S}: {}^{\dagger}q_E^N \mapsto {}^{\ddagger}q_E$$

Very roughly speaking, the main theorem of IUT asserts that:

Relative to such a link, the computation of $\deg {}^{\dagger}q_E^N$ is, up to mild indeterminacies, *compatible* with the computation of $\deg {}^{\ddagger}q_E$.

$$(\Rightarrow \deg q_E^N \stackrel{\text{ind.}\curvearrowright}{=} \deg q_E \Rightarrow (*) \Rightarrow \text{the Szpiro Conjecture})$$

Terminology

- $a(n)$ (*arithmetic*) *holomorphic structure*

$\stackrel{\text{def}}{\Leftrightarrow}$ a (structure which determines a) ring structure

- a *mono-analytic structure*

$\stackrel{\text{def}}{\Leftrightarrow}$ an “underlying” (“non-holomorphic”) structure of a hol. str.

(e.g.: $\mathbb{Q}_p, \pi_1^{\text{ét}}(\mathbb{P}_{\mathbb{Q}_p}^1 \setminus \{0, 1, \infty\})$: hol.; $(\mathbb{Q}_p)_+, \mathbb{Q}_p^{\times}, G_{\mathbb{Q}_p}$: mono-an.)

$F_{\text{mod}} \subseteq F$: the field of moduli of E

l : a prime number

$$K \stackrel{\text{def}}{=} F(E[l](\overline{F}))$$

$\underline{V} \subseteq \mathbb{V}(K)$: the image of a splitting of $\mathbb{V}(K) \rightarrow \mathbb{V}_{\text{mod}} \stackrel{\text{def}}{=} \mathbb{V}(F_{\text{mod}})$

Suppose that F/F_{mod} and K/F_{mod} are *Galois*.

$$S \stackrel{\text{def}}{=} [\text{Spec } \mathcal{O}_K / \text{Gal}(K/F_{\text{mod}})] \quad (\text{the stack-theoretic quotient})$$

$\Rightarrow$ The arith. div. on $\mathcal{O}_F$ determined by q_E can be descended to an arith. div. on S , i.e., by considering the arith. div. on S det'd by

$$\mathfrak{q} \stackrel{\text{def}}{=} (q_{\underline{v}} \stackrel{\text{def}}{=} q_{\underline{v}|_F} \in \mathcal{O}_{F_{\underline{v}|_F}}^{\triangleright} \subseteq \mathcal{O}_{K_{\underline{v}}}^{\triangleright})_{\underline{v} \in \underline{V}} \in \prod_{\underline{v} \in \underline{V}} \mathcal{O}_{K_{\underline{v}}}^{\triangleright}.$$

Note that $\deg \mathfrak{q} \stackrel{\text{def}}{=} [F_{\text{mod}} : \mathbb{Q}]^{-1} \log(\prod_{\underline{v} \in \underline{V}} \#(\mathcal{O}_{K_{\underline{v}}}/q_{\underline{v}} \mathcal{O}_{K_{\underline{v}}})) = \deg q_E$.

Recall An arithmetic line bundle on $\mathcal{O}_K$

$$= \text{a certain pair of a l.b. } \mathcal{L} \text{ on } \mathcal{O}_K \text{ and a metric on } \mathcal{L} \times_{\mathbb{Z}} \mathbb{C} \\ 1 \rightarrow \mu(K) \rightarrow K^\times \xrightarrow{\text{ADiv}} \bigoplus_{w \in \mathbb{V}(K)} (K_w^\times / \mathcal{O}_{K_w}^\times) \rightarrow \text{APic } \mathcal{O}_K \rightarrow 1$$

Categories of Arithmetic Line Bundles on S

$\mathcal{F}_{\text{mod}}^*$: the Frobenioid of arithmetic line bundles on S

Module-theoretic Description

$\mathcal{F}_{\text{mod}}^*$: the Frobenioid of collections $\{a_{\underline{v}} \mathcal{O}_{K_{\underline{v}}}\}_{\underline{v} \in \underline{\mathbb{V}}}$ s.t.

$$a_{\underline{v}} \in K_{\underline{v}}^\times, \quad a_{\underline{v}} \in \mathcal{O}_{K_{\underline{v}}}^\times \text{ for almost } \underline{v} \in \underline{\mathbb{V}}$$

Multiplicative Description

$\mathcal{F}_{\text{MOD}}^*$: the Frobenioid of pairs $(T, \{t_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}})$ s.t.

$$T: \text{an } F_{\text{mod}}^\times\text{-torsor,} \quad t_{\underline{v}} \in T \times^{F_{\text{mod}}^\times} K_{\underline{v}}^\times / \mathcal{O}_{K_{\underline{v}}}^\times$$

$\Rightarrow$ The holomorphic structure of F_{mod} determines

$$\begin{array}{ccccc} \mathcal{F}_{\text{mod}}^* & \xrightarrow{\sim} & \mathcal{F}_{\text{mod}}^* & \xrightarrow{\sim} & \mathcal{F}_{\text{MOD}}^* \\ \{a_{\underline{v}} \mathcal{O}_{K_{\underline{v}}}\} & & \mapsto & & (F_{\text{mod}}^\times, \{\text{ord}_{\underline{v}}(a_{\underline{v}})\}) \end{array}$$

$\mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}}, \mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}}, \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}$: the resp. realifications of $\mathcal{F}_{\text{mod}}^*$, $\mathcal{F}_{\text{mod}}^*$, $\mathcal{F}_{\text{MOD}}^*$,
i.e., obtained by replacing $\bigoplus_{\underline{v}} (K_{\underline{v}}^\times / \mathcal{O}_{K_{\underline{v}}}^\times)$ by $\bigoplus_{\underline{v}} ((K_{\underline{v}}^\times / \mathcal{O}_{K_{\underline{v}}}^\times) \otimes \mathbb{R})$
($\Rightarrow$ The hol. str. of F_{mod} determines $\mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}} \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}} \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}$)

The multiplication by $1/N$ on $\bigoplus_{\underline{v}} ((K_{\underline{v}}^{\times} / \mathcal{O}_{K_{\underline{v}}}^{\times}) \otimes \mathbb{R})$ determines

$$\Theta_{\text{naive}} : {}^{\dagger}\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \xrightarrow{\sim} {}^{\ddagger}\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \quad \text{which maps} \quad {}^{\dagger}\mathfrak{q}^N \mapsto {}^{\ddagger}\mathfrak{q}.$$

Remark

- Θ_{naive} may be regarded as a “deformation of value groups”.
- The link “ Θ_{naive} ” will be eventually established by means of nonarchimedean theta functions (cf. p.21).

$\mathcal{F}_{\text{mod}}^{\otimes}$ depending on hol. str. suited to deg. estimates

$\mathcal{F}_{\text{MOD}}^{\otimes}$ only multiplicative str. not suited to deg. estimates

Goal

Relative to a link such as Θ_{naive} , the computation of $\deg^{\dagger} \mathbf{q}^N$ is, up to mild indeterminacies, *compatible* with the comp. of $\deg^{\dagger} \mathbf{q}$.

Note that $\nexists$ a ring automorphism of $K_{\underline{v}}$ s.t. $q_{\underline{v}}^N \mapsto q_{\underline{v}}$ (if $q_{\underline{v}} \neq 1$).

Thus, Θ_{naive} *cannot be compatible* with the holomorphic structures, i.e., Θ_{naive} may be compatible with only certain mono-analytic str.

(For instance, Θ_{naive} is *compatible* with the local Galois group

$G_{\underline{v}} \stackrel{\text{def}}{=} \text{Gal}(\overline{F}_{\underline{v}}/K_{\underline{v}})$ for each finite $\underline{v} \in \mathbb{V}$

— cf. Θ_{naive} “=” a deformation of value groups.)

On the other hand:

Remark

The “degree computation” is, at least a priori, performed by means of the holomorphic structure under consideration.

Thus, in order to obtain a certain compatibility of the degree computations, we have to establish a “*multiradial representation*” of the degree computations whose coric data consist of suitable mono-analytic structures.

Multiradial Algorithm

Suppose that we are given

- a mathematical object R , i.e., a *radial data*,
- an “underlying” object C of R , i.e., a *coric data*, and
- a func'l algorithm Φ whose input data is (an isomorph of) R .

Example


- R : the one-dimensional complex linear space $\mathbb{C}$
 C : the underlying two-dimensional real linear space $\mathbb{R}^{\oplus 2}$
- R : the field $\mathbb{Q}_p$ C : the underlying additive module $(\mathbb{Q}_p)_+$
- R : the étale fundamental gp $\pi_1^{\text{ét}}(V)$ of a hyperbolic curve $V/\mathbb{Q}_p$
 C : the absolute Galois group $G_{\mathbb{Q}_p}$ of $\mathbb{Q}_p$

Roughly speaking, we shall say that the algorithm Φ is:

- *coric* if Φ depends only on C
- *multiradial* if Φ (is rel'd to R but) may be described in terms of C
- *uniradial* if Φ is not multiradial, i.e., essentially depends on R

If one starts with a coric data “ C ” and applies the alg'm Φ , then:

- *uniradial* $\Rightarrow$ the output *depends* on the choice of a “spoke”
- *multiradial* $\Rightarrow$ the output is *unaffected* by alterations in a “spoke”



(Tautological) Example

$$(R, C) \cong (\mathbb{C}, \mathbb{R}^{\oplus 2})$$

- $\Phi(R) =$ the holomorphic structure on $R \Rightarrow$ uniradial
- $\Phi(R) =$ the real analytic structure on $R \Rightarrow$ coric
- $\Phi(R) =$ the $\mathrm{GL}_2(\mathbb{R})$ -orbit of the hol. str. on $R \Rightarrow$ multiradial

incompatible

H.S. on ${}^{\dagger}\mathbb{C}$

compatible

H.S. of ${}^{\dagger}\mathbb{C}$

$$\begin{array}{ccc} & & \downarrow \\ \text{H.S. of } {}^{\dagger}\mathbb{C} & \longrightarrow & \boxed{\mathbb{R}^{\oplus 2}} \end{array}$$

uniradial

$$\begin{array}{ccc} & & \downarrow \scriptstyle \mathrm{GL}_2(\mathbb{R}) \curvearrowright \\ \text{H.S. of } {}^{\dagger}\mathbb{C} & \xrightarrow[\curvearrowright]{\mathrm{GL}_2(\mathbb{R})} & \boxed{\mathbb{R}^{\oplus 2}} \end{array}$$

multiradial

$$(\text{cf. } \mathrm{GL}_2(\mathbb{R})/\mathbb{C}^{\times} = \mathbb{H}_+ \sqcup \mathbb{H}_-)$$

Summary

- We want to obtain a certain compatibility of the degree computations relative to a link such as Θ_{naive} .
- Θ_{naive} *cannot be compatible* w/ the holomorphic str., i.e., Θ_{naive} is compatible w/ only certain mono-an. str., e.g., $G_{\underline{v}}$.
- On the other hand, the degree computation is, at least a priori, performed by means of the holomorphic structure.
- Thus, we have to establish a *multiradial representation* of the degree computations whose coric data are suitable mono-an. str.

An Approximate Statement of the Main Theorem of IUT (tentative)

$\exists$ A suitable multiradial algorithm whose output data consist of the following three objects $\curvearrowright$ mild indeterminacies

- $\{(\mathcal{O}_{K_{\underline{v}}})_*\}_{\underline{v} \in \mathbb{V}}$ ($*$ = $+$ if $\underline{v}$ is finite, $*$ = $\emptyset$ if $\underline{v}$ is infinite)
- $\mathfrak{q}^N \curvearrowright \prod_{\underline{v} \in \mathbb{V}} (\mathcal{O}_{K_{\underline{v}}})_*$
- $F_{\text{mod}} \curvearrowright \prod_{\underline{v} \in \mathbb{V}} ((K_{\underline{v}})_* \text{ “via } (\mathcal{O}_{K_{\underline{v}}})_* \text{”})$

Moreover, this algorithm is *compatible* with

$$\Theta_{\text{naive}}: {}^\dagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \xrightarrow{\sim} {}^\ddagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}; {}^\dagger \mathfrak{q}^N \mapsto {}^\ddagger \mathfrak{q}.$$

Main Theorem of IUT $\Rightarrow$ Szpiro Conjecture

$$\Theta_{\text{naive}} : {}^\dagger\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \xrightarrow{\sim} {}^\ddagger\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}; \quad {}^\dagger\mathbf{q}^N \mapsto {}^\ddagger\mathbf{q}$$

$$\xRightarrow{\text{Th'm}}$$

$$\left(\begin{array}{c} \{({}^\dagger\mathcal{O}_{K_{\underline{v}}})_*\}_{\underline{v} \in \mathbb{V}} \\ {}^\dagger\mathbf{q}^N \curvearrowright \prod ({}^\dagger\mathcal{O}_{K_{\underline{v}}})_* \\ {}^\dagger F_{\text{mod}} \curvearrowright \prod ({}^\dagger K_{\underline{v}})_* \end{array} \right) \xrightarrow[\sim]{\text{ind.} \curvearrowright} \left(\begin{array}{c} \{({}^\ddagger\mathcal{O}_{K_{\underline{v}}})_*\}_{\underline{v} \in \mathbb{V}} \\ {}^\ddagger\mathbf{q}^N \curvearrowright \prod ({}^\ddagger\mathcal{O}_{K_{\underline{v}}})_* \\ {}^\ddagger F_{\text{mod}} \curvearrowright \prod ({}^\ddagger K_{\underline{v}})_* \end{array} \right)$$

$\Rightarrow$

- $c_{\text{mod}} : {}^\dagger\mathcal{F}_{\text{mod}}^{\otimes} \xrightarrow[\sim]{\text{ind.} \curvearrowright} {}^\ddagger\mathcal{F}_{\text{mod}}^{\otimes}$ which maps $\{{}^\dagger q_{\underline{v}}^N {}^\dagger\mathcal{O}_{K_{\underline{v}}}\} \mapsto \{{}^\ddagger q_{\underline{v}}^N {}^\ddagger\mathcal{O}_{K_{\underline{v}}}\}$
- $c_{\square} : {}^\square\mathcal{F}_{\text{mod}}^{\otimes} \xrightarrow{\sim} {}^\square\mathcal{F}_{\text{MOD}}^{\otimes}$ which maps $\{{}^\square q_{\underline{v}}^{(N)} {}^\square\mathcal{O}_{K_{\underline{v}}}\} \mapsto {}^\square\mathbf{q}^{(N)}$

which are *compatible* with Θ_{naive} , i.e.,

fit into a diagram that is *commutative*, up to *mild indeterminacies*

$$\begin{array}{ccc}
 {}^{\dagger}\mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}} & \xrightarrow{c_{\text{mod}}} & {}^{\ddagger}\mathcal{F}_{\text{mod}}^{\otimes \mathbb{R}} \\
 c_{\dagger} \downarrow & & \downarrow c_{\ddagger} \\
 {}^{\dagger}\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} & \xrightarrow{\Theta_{\text{naive}}} & {}^{\ddagger}\mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}
 \end{array}$$

$$\Rightarrow \left(\begin{array}{ccc}
 \{ {}^{\dagger}q_v^{N\dagger} \mathcal{O}_{K_v} \} & \Rightarrow & \{ {}^{\ddagger}q_v^{N\ddagger} \mathcal{O}_{K_v} \}, \quad \{ {}^{\ddagger}q_v^{\ddagger} \mathcal{O}_{K_v} \} \\
 \uparrow & & \downarrow \\
 {}^{\dagger}\mathfrak{q}^N & \Leftarrow & {}^{\ddagger}\mathfrak{q}
 \end{array} \right)$$

$$\Rightarrow \{ {}^{\ddagger}q_v^{\ddagger} \mathcal{O}_{K_v} \} \overset{\text{log-vol.}}{\subseteq} \bigcup_{\text{indeterminacies}} \{ {}^{\ddagger}q_v^{N\ddagger} \mathcal{O}_{K_v} \}$$

$$\Rightarrow -\deg \mathfrak{q} \leq -\deg \mathfrak{q}^N + C, \quad \text{i.e., } (*) \text{ in p.4}$$

$$\Rightarrow \text{the Szpiro Conjecture}$$

An Approximate Statement of the Main Theorem of IUT (tentative)

$\exists$ A suitable multiradial algorithm whose output data consist of the following three objects $\curvearrowright$ mild indeterminacies

- $\{(\mathcal{O}_{K_{\underline{v}}})_*\}_{\underline{v} \in \mathbb{V}}$ ($*$ = $+$ if $\underline{v}$ is finite, $*$ = $\emptyset$ if $\underline{v}$ is infinite)
- $\mathfrak{q}^N \curvearrowright \prod_{\underline{v} \in \mathbb{V}} (\mathcal{O}_{K_{\underline{v}}})_*$
- $F_{\text{mod}} \curvearrowright \prod_{\underline{v} \in \mathbb{V}} ((K_{\underline{v}})_* \text{ “via } (\mathcal{O}_{K_{\underline{v}}})_* \text{”})$

Moreover, this algorithm is *compatible* with

$$\Theta_{\text{naive}}: {}^\dagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \xrightarrow{\sim} {}^\ddagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}; {}^\dagger \mathfrak{q}^N \mapsto {}^\ddagger \mathfrak{q}.$$

Recall ($\underline{v} \in \underline{\mathbb{V}}$: fin.; $p \stackrel{\text{def}}{=} p_{\underline{v}}$) $(\mathcal{O}_{K_{\underline{v}}})_+ \in \text{output}$, $G_{\underline{v}}$: coric

Unfortunately, it is known that:

$\nexists$ a func'l (w.r.t. open injections) alg'm for rec. " $(\mathcal{O}_{K_{\underline{v}}})_+$ " from " $G_{\underline{v}}$ ".

$\mathcal{I}_{K_{\underline{v}}} \stackrel{\text{def}}{=} \frac{1}{2p} \text{Im}(\mathcal{O}_{K_{\underline{v}}}^{\times} \rightarrow \mathcal{O}_{K_{\underline{v}}}^{\times} \otimes_{\mathbb{Z}} \mathbb{Q} \xrightarrow{\log_p} (K_{\underline{v}})_+)$: the *log-shell* of $K_{\underline{v}}$

- $\mathcal{I}_{K_{\underline{v}}}$: a finitely generated free $\mathbb{Z}_p$ -module
- $(\mathcal{O}_{K_{\underline{v}}})_+, \log_p(\mathcal{O}_{K_{\underline{v}}}^{\times}) \subseteq \mathcal{I}_{K_{\underline{v}}} \subseteq \mathcal{I}_{K_{\underline{v}}} \otimes_{\mathbb{Z}} \mathbb{Q} = (K_{\underline{v}})_+$
- $[\mathcal{I}_{K_{\underline{v}}} : (\mathcal{O}_{K_{\underline{v}}})_+] (< \infty)$ can be comp'd by the top. gp str. of $G_{\underline{v}}$
- $G_{\underline{v}} \xrightarrow[\text{algorithm}]{\exists \text{ func'l}}$ an isomorph of $\mathcal{I}_{K_{\underline{v}}}$

Thus, " $\{(\mathcal{O}_{K_{\underline{v}}})_*\}_{\underline{v} \in \underline{\mathbb{V}}}$ " in Th'm should be replaced by $\{\mathcal{I}_{\underline{v}} \stackrel{\text{def}}{=} \mathcal{I}_{K_{\underline{v}}}\}_{\underline{v} \in \underline{\mathbb{V}}}$
 (where the log-shell at an infinite $\underline{v} \in \underline{\mathbb{V}} \stackrel{\text{def}}{=} \pi \cdot \mathcal{O}_{K_{\underline{v}}}$).

Recall $\mathfrak{q}^N, F_{\text{mod}} \in \text{output}$

Unfortunately, by various indeterminacies arising from the operation of “passing from holom’c str. to mono-anal’c str.”, it is difficult to obtain multiradial representations of $\mathfrak{q}^N, F_{\text{mod}}$ themselves directly. To establish a mul’l alg’m of the desired type, we rep. multiradially a *suitable function* whose special value is $q_{\underline{v}}^N$ or an $\in F_{\text{mod}}$.

- “ $q_{\underline{v}}^N$ ” will be represented as a special value of a (multiradially represented) *theta function*.
- “ F_{mod} ” will be represented as a set of special values of (multiradially represented) *κ -coric functions*.

An Approximate Statement of the Main Theorem of IUT

For a “general E/F ”,

$\exists$ a suitable multiradial algorithm whose output data consist of the following three objects $\curvearrowright$ mild indeterminacies

- the collection of log-shells $\{\mathcal{I}_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$
- the theta values $(= \{q^{j^2/2l}\}_{1 \leq j \leq l^* \stackrel{\text{def}}{=} \frac{l-1}{2}}) \curvearrowright \prod_{\underline{v} \in \underline{\mathbb{V}}} \mathcal{I}_{\underline{v}}$
- F_{mod} via κ -coric functions $\curvearrowright \prod_{\underline{v} \in \underline{\mathbb{V}}} ((K_{\underline{v}})_+ \text{ “via } \mathcal{I}_{\underline{v}} \text{”})$

Moreover, this alg'm is *compatible* w/ the Θ -link (more precisely,

$\Theta_{\text{LGP}}^{\times \mu}$ -link) “ $\dagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}} \xrightarrow{\sim} \ddagger \mathcal{F}_{\text{MOD}}^{\otimes \mathbb{R}}$ ”; “ $\dagger$ theta values $\mapsto \ddagger q^{1/2l}$ ”.