

Limits of tangents on a hypersurface  
(in collaboration with J. P. G. Henry)

by Lê Dũng Tráng (Ecole Polytechnique, Paris)

Let  $V := \{z \in \mathbb{C}^{n+1} \mid f(z) = 0\}$  be an analytic complex hypersurface. We shall suppose that  $0 \in V$  and that  $0$  is a singular point of  $V$ . We want to know what is the set of all possible limits of sequences of tangent hyperplanes to  $V$  at sequences of smooth points of  $V$  tending to  $0$ . More precisely, let us denote by  $\check{\mathbb{P}}^n$  the space of hyperplane directions of  $\mathbb{C}^{n+1}$ , let  $\Sigma$  be the singular locus of  $V$  and let  $\varphi: V - \Sigma \rightarrow \check{\mathbb{P}}^n$  be the map which makes correspond to a smooth point of  $V$  the direction of its tangent hyperplane. Then let us call  $\tilde{V}$  the closure of the graph of  $\varphi$  in  $V \times \check{\mathbb{P}}^n$ . The first projection  $V \times \check{\mathbb{P}}^n \rightarrow V$  induces a mapping  $\pi: \tilde{V} \rightarrow V$  which is called the Jacobian blowing-up of  $V$ . The set we are interested about is  $\pi^{-1}(0)$ .

Notice that  $\dim \pi^{-1}(\Sigma) = \dim V - 1$ . Thus if  $0 \in V$  is an isolated singularity  $\dim \pi^{-1}(0) = \dim V - 1$ .

First notice the following lemma:

Lemma 1 ([1]) Let  $(x_n)$  be a sequence of smooth points of  $V$  such that the lines  $0x_n$  tend to  $l$  and the hyperplanes  $T(x_n, V)$  tend to  $T$ , then  $l$  is contained in  $T$ .

Recall that all possible limits of lines  $0x_n$  for sequences of points of  $V$  tending to  $0$  define the tangent cone of  $V$  at  $0$  and the corresponding set in the projective space  $\mathbb{P}^n$  of lines through  $0$  is the Proj of this tangent cone.

Now we have the following lemma which was indicated to us by Professor O. Zariski:

Lemma 2 Let us suppose that the tangent cone of  $V$  at  $0$  is reduced. Then consider  $(x_n)$  a sequence of  $\overset{\text{smooth}}{\text{points}}$  of  $V$  which tends to  $0$  and such that  $T(x_n, V)$  tends to  $T$  and  $Ox_n$  tends to  $l$ . Moreover suppose that  $l$  gives a smooth point of the Proj of the tangent cone of  $V$  at  $0$ , then  $T$  is tangent to the tangent cone of  $V$  at  $0$  along  $l$ .

Such a lemma has led our interests to compare the limits of tangents at  $0$ , say  $\pi^{-1}(0)$  defined above, and the limits of secants at  $0$ , say the Proj of the tangent cone at  $0$ .

We obtain the following result :

Theorem 3 Suppose  $n=2$  and  $0 \in V$  is an isolated singularity, then the limits of tangents of  $V$  at  $0$  is the union of the dual curve of the curve, Proj of the reduced tangent cone at  $0$ , and a finite number of lines which corresponds to pencils of hyperplanes going through the singular lines of the  $\overset{\text{reduced}}{\text{Tangent}}$  cone and a finite number of non-singular lines of this reduced tangent cone.

Remark : The non-singular lines of the theorem are specified in the proof (cf theorem 6)

The proof of the theorem 3 is based on some results of B. Teissier from [4] and a geometrical study involving some results <sup>about</sup> equisingularity of O. Zariski [5].

First let us define:

Definition We shall say that a hyperplane  $H$  cuts  $V$  generically at  $0$  if  $V \cap H$  has an isolated singularity at  $0$  and the Milnor number of  $V \cap H$  at  $0$  is minimum among all <sup>the</sup> hyperplane sections with an isolated singularity at  $0$ .

Let us remind that such a hyperplane section has a well-defined topology <sup>at least</sup> (when  $n \neq 3$ ) because of the results of [2].

The result of B. Teissier can be stated as follows:

If  $0 \in V$  is an isolated singularity,

Theorem 4 (cf [4]) The hyperplane  $H$  cuts  $V$  generically at  $0$  if and only if  $H$  is not a limit of tangents of  $V$  at  $0$ , i.e.  $H \notin \pi^{-1}(0)$ .

Now let  $n=2$ . Call  $\tilde{Z}_1 \xrightarrow{\pi} \mathbb{C}^3$  the blowing-up of the point  $0$ . Let  $V_1$  be the strict transform of  $V$  by  $\pi$ . Let  $H$  be a hyperplane of  $\mathbb{C}^3$  and  $H_1$  be its strict transform by  $\pi$ . Call  $C = H \cap V$  and  $C_1$  the strict transform of  $C$ . Thus  $C_1 = H_1 \cap V_1$ .

Then  $C$  is reduced if and only if  $C_1$  is reduced. Thus

$C$  is reduced if and only if  $\pi^{-1}(0) \cap H_1$  cuts transversally the reduced curve  $\pi^{-1}(0) \cap V_1$  and near each point of intersection  $H_1$  cuts the smooth part of  $V_1$  transversally.

But we have the following lemma:

Lemma 5 (cf [3]) Let  $C$  be a plane curve of multiplicity  $n$  at  $O$ , let  $C_1$  be the strict transform of  $C$  after blowing-up  $O$ . Let  $O_1, \dots, O_k$  be the points of  $C_1$  over  $O$ , then:

$$\mu(C, O) = \mu(C_1, O_1) + \dots + \mu(C_1, O_k) + n(n-1) - (k-1)$$

Thus if we apply it to our above situation we find that at each point where  $p^{-1}(O) \cap H_1$  cuts  $p^{-1}(O) \cap V_1$  the local Milnor number of  $H_1 \cap V_1$  must be the minimum one to get the minimum one for  $H \cap V$  at  $O$ .

Then consider the components  $\Gamma_1, \dots, \Gamma_\ell$  of  $p^{-1}(O) \cap V_1$ . We have the following case:

- a)  $V_1$  is singular along  $\Gamma_i$ . Thus  $V_1$  is equisingular along  $\Gamma_i$  outside a finite number of exceptional points <sup>defines lines</sup> we shall call exceptional secants of  $V$  at  $O$ . Remark that <sup>the</sup> singular points of  $\Gamma_i$  and points of  $\Gamma_i \cap \Gamma_j$  ( $j \neq i$ ) are among these exceptional points;
- b)  $V_1$  is not singular along  $\Gamma_i$  except maybe at a finite number of points. The <sup>corresponding lines of these points</sup> will be exceptional secants of  $V$  at  $O$ , too.

Then using Zariski's theory of equisingularity we can prove:

Theorem 6 A hyperplane  $H$  cuts  $V$  generically at  $O$  if and only if it does not contain any exceptional secants.

## Bibliography

- [1] Lê Dũng Tráng Singularités des hypersurfaces complexes, Acta Sc. Vietn., Hanoi, 1971.
- [2] Lê Dũng Tráng - C.P. Ramanujam The invariance of Milnor's number implies the invariance of the topological type, to appear in Amer. J. Math.
- [3] J. J. Risler Sur l'Idéal Jacobien, Bull. Soc. Mat. Fr.<sup>t.99</sup>, 1971, 30
- [4] B. Teissier Cycles évanescents, Sections planes et conditions de Whitney, in Singularités à Cargèse, Astérisque 7 et 8, France
- [5] O. Zariski Studies in Equisingularities I, II Amer. J. Math., t.87 (1965), 507-536 et 972-1006