

ASYMPTOTIC THEORY IN CHANGE POINT PROBLEM

Kumamoto University 坂田年男 (Toshio Sakata)

1. Introduction.

Let  $\Pi_k$  be a normal population whose mean and variance are  $\mu_k$  and  $\sigma_k^2$ , respectively,  $k=1,2,\dots,n$ . Many authors have examined the problem of testing the null hypothesis

$$H_1: \mu_1 = \mu_2 = \dots = \mu_n$$

against the two sided alternative

$$K_1: \bigcup_{1 \leq k \leq n-1} K_{1k}, \text{ where } K_{1k}: \mu_1 = \mu_2 = \dots = \mu_k \neq \mu_{k+1} = \dots = \mu_n,$$

or against the one-sided alternative

$$K_2: \bigcup_{1 \leq k \leq n-1} K_{2k}, \text{ where } K_{2k}: \mu_1 = \mu_2 = \dots = \mu_k > \mu_{k+1} = \dots = \mu_n.$$

This problem is known as the problem of detecting a change in means, (see, e.g., Sen and Srivastava (1975), or Srivastava and Worsley (1986)). Apart from normal means, Hsu (1979) treated the problem of detecting a change of means in a sequence of gamma variables. Worsley (1986) treated the problem somewhat generally, that is, in the framework of a sequence of exponential family random variables. He derived the likelihood ratio test for the two sided alternative and obtained the null distribution of the test, conditional on the total sum of observations, in the terms of iterated integrals. More recently, Yao and Davis (1986)

has determined the asymptotic null distribution of the likelihood ratio test as the number of the populations  $n$  tends to infinity for the case of normal means. Siegmund (1986) also has treated the approximation of the asymptotic null distribution of the modified likelihood ratio test.

In this paper we will proceed as follows. In Section 2 some mathematical preliminaries are provided. Section 3 is expository and the asymptotic distribution of the likelihood ratio test for a shift in normal means given by Yao and Davis (1986) reviewed briefly. In Section 4 we treat the asymptotic null distribution of a modified likelihood ratio test for a shift in normal variances.

## 2. Mathematical preliminaries

Definition 2.1. A process  $W(t)$ ,  $t \geq 0$ , satisfying the following conditions is called the standard Brownian motion.

- (1) Gaussian
- (2)  $EW(t) = 0$ ,  $\text{Cov}(W(s), W(t)) = s \wedge t$
- (3) continuous

Proposition 2.1 (Law of the Iterated Logarithm). Let  $W(t)$ ,  $t \geq 0$ , be the standard Brownian motion. Then

$$\limsup_{t \rightarrow 0^+} |W(t)| / \sqrt{t \log |\log t|} = 1 \quad \text{a.s.}$$

$$\limsup_{t \rightarrow \infty} |W(t)| / \sqrt{t \log |\log t|} = 1 \quad \text{a.s.}$$

Definition 2.2. A Gaussian process  $W^0(t)$  defined by

$$W^0(t) = W(t) - tW(1), \quad 0 \leq t \leq 1,$$

is called the standard Brownian bridge.

Proposition 2.2. Let  $W^0(t)$  be the standard Brownian bridge. Then

$$W^0(0) = W^0(1) = 0, \quad EW^0(t) = 0, \quad \text{and} \quad \text{Cov}(W^0(s), W^0(t)) = s(1-t), \quad s \leq t.$$

Definition 2.3. A stationary Gaussian process  $X(t)$ ,  $-\infty < t < \infty$ , satisfying

$EX(t) = 0$  and  $\text{Cov}(X(s), X(t)) = e^{-(t-s)}$ ,  $s \leq t$ , is called the Ornstein-Uhlenbeck (O-U) process.

Note that we use the symbol  $W(t)$ ,  $W^0(t)$  and  $X(t)$  for these three particular process respectively.

In this paper some transformations among these three process play an essential role, and we summarize them here.

Proposition 2.4. Let  $Y((1/2)\log t) = W(t)/\sqrt{t}$ ,  $t > 0$ . Then  $Y(t)$ ,  $-\infty < t < \infty$ , is the O-U process  $X(t)$ .

Remark: Proposition 2.4 implies that a square root boundary for  $W(t)$  is transformed to a constant boundary for  $X(t)$ .

Proposition 2.5. (DeLong (1981), p. 2212). Let  $Y(t)$ ,  $t \geq 0$ , be defined by

$$Y(t) = (1+t)W^0(t/(1+t)).$$

Then  $Y(t)$  is the standard Brownian motion  $W(t)$  and

$$P\left\{\max_{t_1 < t < t_2} W^0(t)/\sqrt{t(1-t)} < c\right\} = P\left\{\max_{1 < t < f(t_1, t_2)} W(t)/\sqrt{t} < c\right\},$$

where  $f(t_1, t_2) = t_2(1-t_1)/t_1(1-t_2)$ .

Remark: Proposition 2.5 implies that a  $\sqrt{t(1-t)}$  boundary for  $W^0(t)$  is transformed to a  $\sqrt{t}$  boundary for  $W(t)$  and therefore to a constant boundary for  $X(t)$ .

Proposition 2.6 (DeLong(1981), p. 2203).

$$P\left\{\max_{1 \leq t \leq T} |W(t)|/\sqrt{t} < c\right\} = \sum_{i=1}^{\infty} \alpha_i(c, q) T^{-\beta_i(c, q)},$$

where  $q=1/2$ ,  $\beta_i$  is the  $i$ -th root of  $M(\beta_i+q, q, -c^2/2)=0$ ,  $M$  is the confluent hypergeometric function, and  $\alpha_i$  is equal to

$$\frac{e^{-c^2/2} (c^2/2)^q M(\beta_i+q, q+1, -c^2/2)}{\Gamma(q+1) \beta_i (d/dx)M(x+q, q, -c^2/2)}$$

with the derivative of  $M$  evaluated at  $x=\beta_i$ .

Proposition 2.7 (DeLong (1981), p. 2205). For  $T$  and  $c$  large,

$$P\left\{\max_{1 \leq t \leq T} |W(t)|/\sqrt{t} \geq c\right\} \sim \frac{c}{\sqrt{2\pi}} e^{-c^2/2} \{(\log T)(1-1/c^2)+2/c^2+O(c^{-4})\}$$

Next we summarize the extreme value theory briefly, (see, e.g., Resnik(1987), p. 9). Let  $X_1, X_2, \dots, X_n$  be a i.i.d. sequence of random variables with common distribution  $F(x)$ . Set  $M_n = \max_{1 \leq i \leq n} X_i$

Definition 2.4. Suppose there exist  $a_n > 0$  and  $b_n \in \mathbb{R}_+^n$  such that

$P\{(M_n - b_n)/a_n < x\} \rightarrow G(x)$  weakly as  $n \rightarrow \infty$  where  $G(x)$  is non-degenerate. Then  $F$  is said to belong to the domain of attraction of  $G$  and  $G$  is said a extreme value distribution.

Proposition 2.8 (Gnedenko(1945)). Extreme value distribution is of the type of one of the following three cases:

(1)  $\Phi_{\alpha}(x) = 0$  for  $x < 0$  and  $\exp(-x^{\alpha})$  for  $x \geq 0$  with  $\alpha > 0$

(2)  $\Psi_{\alpha}(x) = \exp(-(-x)^{\alpha})$  for  $x < 0$  with  $\alpha > 0$  and 1 for  $x \geq 0$

(3)  $\Lambda(x) = \exp(-e^{-x})$   $x \in \mathbb{R}^1$ .

An extension of an extreme value distribution to the maximum of a stationary Gaussian sequence was given by Berman (1964) and extensions to the maximum of a continuous parameter

stationary Gaussian process have appeared in various literatures, (Berman(1974,1980), Pickands (1969), etc). Recently Berman(1984) has treated a stationary non-Gaussian case. Here we refer only to a theorem by Pickands (1969).

Proposition 2.9 (Pickands(1969)). Let  $Y(t)$ ,  $-\infty < t < \infty$ , be a stationary Gaussian process such that  $EY(t)=0$  and  $\text{Cov}(Y(s), Y(s+t))=r(t)$ . Let

$$Z(T) = \max_{0 \leq t \leq T} Y(t). \text{ If } r(t) = 1 - C|t|^\alpha + o(|t|^\alpha) \text{ and } \lim_{t \rightarrow \infty} r(t) \log t = 0,$$

then for all  $x$ ,  $-\infty < x < \infty$ ,

$$\lim_{T \rightarrow \infty} P\{ a(T)^{-1}(Z(T) - b(T)) \leq x \} = \exp(-e^{-x}),$$

where  $a(T) = (2 \log T)^{-1/2}$  and  $b(T) = a(T)^{-1} +$

$$a(T) \left\{ \left( (1/\alpha - 1/2) \log \log T + \log \left( (2\pi)^{-1/2} C^{1/\alpha} H 2^{(2-\alpha)/2\alpha} \right) \right) \right\}$$

and  $0 < H = \lim_{T \rightarrow \infty} T^{-1} \int_0^\infty e^{-s} P\{ \sup_{0 \leq t \leq T} Y(t) > s \} ds < \infty$ ,

where  $Y(t)$  is a stationary Gaussian process with means and covariances:

$$EY(t) = -|t|^\alpha, \text{ Cov}(Y(s), Y(t)) = |s|^\alpha + |t|^\alpha - |t-s|^\alpha.$$

Remark: In the expression of  $b(T)$ ,  $(1/\alpha - 1/2)$  is described as  $(1/2 - 1/\alpha)$  in the original paper of Pickands, but from the book of Leadbetter et al (1983), p.237, Theorem 12.3.5 or from the process of the original proof by Pickands we know  $(1/\alpha - 1/2)$  is a correct one.

For the O-U process  $X(t)$ ,  $r(t) = e^{-t} = 1 - |t| + o(|t|)$ , and  $C = \alpha = 1$ .

For  $\alpha = 1$ ,  $H = 1$  (Pickands (1969), p. 77), and we have the next

proposition.

Proposition 2.10. Let  $Z(T) = \max_{0 \leq t \leq T} X(t)$  and

$a(T) = (2 \log T)^{-1/2}$  and  $b(T) = a(T)^{-1} + a(T) \{ (1/2) \log \log T - \log \sqrt{\pi} \}$

Then

$$\lim_{T \rightarrow \infty} P\{a(T)^{-1}(Z(T) - b(T)) \leq x\} = \exp(-e^{-x}),$$

and moreover

$$\lim_{T \rightarrow \infty} P\{a(T)^{-1}(Z(T) - b^*(T)) \leq x\} = \exp(-(1/\sqrt{\pi})e^{-x}),$$

where  $b^*(T) = a(T)^{-1} + (1/2)a(T) \log \log T$ .

Proof of Proposition 2.10. The former part is trivial from Proposition 2.9. We must show the last part of the proposition.

Since  $b(T) = b^*(T) - a(T) \log \sqrt{\pi}$ ,

$$\lim_{T \rightarrow \infty} P\{a(T)^{-1}(Z(T) - b^*(T)) \leq y - \log \sqrt{\pi}\} = \exp(-e^{-y}),$$

and setting  $x = y - \log \sqrt{\pi}$ ,  $e^{-y} = (1/\sqrt{\pi})e^{-x}$ , which show the last part of Proposition 2.10.

### 3. A shift in normal means

For a shift in normal means the likelihood ratio test rejects  $H$  if  $T$  is too large, where  $T = \max_{1 \leq k \leq n-1} |T_k|$  and

$$T_k = \{(S_k/\sqrt{n} - (k/n)S_n/\sqrt{n})\} / \sqrt{(k/n)(1-(k/n))}$$

and for a one-sided shift,  $T$  must be replaced by  $T^*$  where

$T^* = \max\{0, \max_{1 \leq k \leq n-1} T_k\}$ , and  $S_k = \sum_{i=1}^k X_i$ ,  $k=1, 2, \dots, n$ , (see, e.g., Yao

and Davis, (1986), p.343).

Hawkins(1977) tried to determine the normalizing constants

and the limiting distribution from heuristic arguments, and suggested an approximation by an stationary Gaussian sequence satisfying Berman's condition(1964):  $r(n)\log r(n)\rightarrow 0$ . Yao and Davis(1986) determined exactly the normalizing constants and the limiting distribution.

One of the motivations of this section was to understand the reason why the coefficients  $2/\sqrt{\pi}$  or  $1/\sqrt{\pi}$  appear in the limiting distribution in Proposition 3.1., which do not appear in the extreme value distribution of i.i.d. sequence,  $\Lambda(x)=\exp(-e^{-x})$ , in Proposition 2.8. Another was to know the origin of the normalizing constants  $a_n$  and  $b_n$ .

Proposition 3.1 (Yao and Davis(1986)).

$$\lim_{n \rightarrow \infty} P(T \leq a_n x + b_n) = \exp(-(2/\sqrt{\pi})e^{-x}),$$

$$\lim_{n \rightarrow \infty} P(T^* \leq a_n x + b_n) = \exp(-(1/\sqrt{\pi})e^{-x}),$$

where  $a_n = (2 \log \log n)^{-1/2}$  and  $b_n = a_n^{-1} + (a_n/2) \log \log \log n$ .

For  $T^*$  their arguments are briefly reviewed here. First note  $T_k$  is represented as

$$T_k = W^0(t_k) / \sqrt{t_k(1-t_k)}, \quad t_k = k/n, \quad k=1,2,\dots,n-1,$$

and the distribution of  $T^*$  is essentially approximable by that of the maximum of a stationary Gaussian sequence, (see, Hawkins, 1979, p.184). They proceeded as follows. First they showed from the arguments based on the law of the iterated logarithm of the Brownian motion  $W(t)$  and the symmetry of the Brownian bridge  $W^0(t)$  that

$$P\left(\max_{1 \leq k \leq n-1} T_k \leq a_n x + b_n\right) \sim P\left(\max_{1 \leq k \leq n/\log n} T_k' \leq a_n x + b_n\right)^2$$

where  $T_k' = W(t_k)/\sqrt{t_k}$ ,  $k=1,2,\dots,n/\log n$ . Secondly, noting that the sequence  $T_k' = W(t_k)/\sqrt{t_k}$ ,  $k=1,2,\dots,n/\log n$ , is equal in distribution to the sequence  $T_k'' = W(k)/\sqrt{k}$ ,  $k=1,2,\dots,n/\log n$ , they showed that

$$\begin{aligned} P\left\{\max_{1 \leq k \leq n/\log n} T_k' \leq a_n X + b_n\right\}^2 &\sim P\left\{\max_{1 \leq k \leq n/\log n} T_k'' \leq a_n X + b_n\right\}^2 \\ &\sim P\left\{\max_{1 \leq k \leq n/\log n} T_k'' \leq a_{n/\log n} X + b_{n/\log n}\right\}^2. \end{aligned}$$

Finally the last term was evaluated through the theorem of Darling and Erdos(1956).

The theorem by Darling and Erdos(1956) is now briefly reviewed below. In short, the theorem evaluates the maximum of  $T_k'' = W(k)/\sqrt{k}$ ,  $k=1,2,\dots,n$ , by making the transformation given in Proposition 2.4, that is, by embedding this into the O-U process  $X(t)$ . Further the limiting distribution of the maximum of the sequence  $X(\tau_k)$ ,  $\tau_k = (1/2)\log k$ ,  $k=1,2,\dots,n$ , is shown to be the same with that of the maximum over the interval  $(0, \tau_n)$ , and the limiting distribution of the last quantity is evaluated.

Proposition 3.2 (Darling and Erdős(1956)).

$$\lim_{n \rightarrow \infty} P\left\{\max_{1 \leq k \leq n-1} S_k/\sqrt{k} \leq a_n X + b_n\right\} = \exp\left\{(-1/2\sqrt{\pi})e^{-x}\right\},$$

where  $a_n = \sqrt{2\log\log n}$  and  $b_n = a_n^{-1} + (a_n/2)\log\log\log n$ .

The proof of Proposition 3.2 is sketched as follows. After observing that

$$P\left\{\max_{1 \leq k \leq n} S_k/\sqrt{k} \leq a_n X + b_n\right\} = P\left\{\max_{1 \leq k \leq n} X(\tau_k) \leq a_n X + b_n\right\},$$

where  $X$  is the O-U process and  $\tau_k = (1/2)\log k$ , they showed the next lemma.

Lemma (Darling and Erdős(1956)).

$$(1) \lim_{c \rightarrow \infty} P\{T(c) > \mu(c)y\} = e^{-y}.$$

$$(2) (1/2)\log N(c) - T(c) \rightarrow 0 \text{ in probability as } c \rightarrow \infty.$$

where  $T(c) = \inf\{t > 0; X(t) \geq c\}$ ,  $N(c) = \min\{k \geq 1; X(\tau_k) \geq c\}$ , and

$$\mu(c) = (\sqrt{2\pi}/c)e^{c^2/2}.$$

Note that (1) implies that

$$\lim_{c \rightarrow \infty} P\left\{\max_{0 \leq t \leq \mu(c)y} X(t) \leq c\right\} = \lim_{c \rightarrow \infty} P\{T(c) > \mu(c)y\} = e^{-y},$$

and this is rewritten that

$$\lim_{T \rightarrow \infty} P\left\{\max_{0 \leq t \leq T} X(t) \leq d(T)\right\} = e^{-y},$$

where  $d(T) = a(T)^{-1} + (1/2)a(T)\log\log T - a(T)\log\sqrt{\pi}y$ ,

and  $a(T) = (2\log T)^{-1/2}$ . Setting  $b(T) = a(T)^{-1} + (1/2)a(T)\log\log T$

and  $x = \log\sqrt{\pi}y$ , we see that (1) implies

$$\lim_{T \rightarrow \infty} P\left\{\max_{0 \leq t \leq T} X(t) \leq a(T)x + b(T)\right\} = \exp\left\{-\frac{1}{\sqrt{\pi}}e^{-x}\right\}.$$

This result is coincident with Proposition 2.9.

On the other hand (2) implies

$$\begin{aligned} \lim_{c \rightarrow \infty} P\left\{\max_{0 \leq t \leq \mu(c)y} X(t) \leq c\right\} &= \lim_{c \rightarrow \infty} P\{T(c) > \mu(c)y\} = \lim_{c \rightarrow \infty} P\left\{\tau_{N(c)} \geq \mu(c)y\right\} \\ &= \lim_{c \rightarrow \infty} P\left\{\max_{0 \leq \tau_k \leq \mu(c)y} X(\tau_k) \leq c\right\} = \lim_{c \rightarrow \infty} P\left\{\max_{0 \leq \tau_k \leq T} X(\tau_k) \leq a(T)x + b(T)\right\} \\ &= \exp\left\{-\frac{1}{\sqrt{\pi}}e^{-x}\right\}. \end{aligned}$$

That is, (2) implies the limiting distributions are identical for the maximum of  $X(t)$  over the discrete points  $\tau_k$ ,  $k=1,2,\dots,n$  and over the interval  $[0, \tau_n]$ , and (1) is applicable. By solving the equation  $\mu(c)y = (1/2)\log n$ , the normalizing constants  $a_n$  and  $b_n$  in Proposition 3.1 was obtained, and this constants are also the normalizing constants for the likelihood ratio test  $T$  or  $T^*$ , as

shown by Yao and Davis.

Finally we close this section after stating a question:

Question: Can we get Proposition 3.1. ?, by observing

$$P\left\{\max_{1 \leq k \leq n} W^0(t_k) / \sqrt{t_k(1-t_k)} \leq c\right\} = P\left\{\max_{1 \leq k \leq n} X(\tau_k) \leq c\right\},$$

where  $t_k = k/n$  and  $\tau_k = (1/2) \log(t_k/(1-t_k))$ , and then following the line of the proof of Lemma of Darling and Erdős.

#### 4. A shift in variances

In this section we treat the problem of detecting a change in a sequence of variances of univariate normal populations and study the asymptotic behavior of the modified likelihood ratio test as the number of populations,  $n$ , tends to infinity.

Let  $X_k$ ,  $k=1, \dots, n$  be distributed as  $\chi^2(\nu_k)$  multiplied by a constant  $\sigma_k^2$ , where  $\chi^2(\nu)$  denotes a chi-square random variable with  $\nu$  degrees of freedom. Our problem is to test the null hypothesis

$$H: \sigma_1^2 = \dots = \sigma_n^2$$

against the alternative

$$K = \bigcup_{k=1}^{n-1} K_k, \text{ where } K_k: \sigma_1^2 = \dots = \sigma_k^2 \neq \sigma_{k+1}^2 = \dots = \sigma_n^2$$

Let

$$S_k = \sum_{i=1}^k X_i, \text{ and } S_k^* = \sum_{i=k+1}^n X_i, \text{ (} k=1, \dots, n-1 \text{) and } S = \sum_{i=1}^n X_i$$

and let

$$N_k = \sum_{i=1}^k \nu_i, \quad N_k^* = \sum_{i=k+1}^n \nu_i, \text{ and } N = \sum_{i=1}^n \nu_i.$$

The likelihood ratio test rejects  $H$  if  $\max_{1 \leq k \leq n-1} T_k$  is too large, where each  $T_k$  is the likelihood ratio test for testing  $H$  against  $K_k$  and given by the following quantity,

$$T_k = N \log(S/N) - N_k \log(S_k/N_k) - N_k \log(S_k^*/N_k^*).$$

In the problem of detecting a change of normal means, Siegmund(1986) proposed the modified likelihood ratio test given by

$$\max_{1 < m_0 \leq k \leq m_1 < n} \Lambda_k,$$

where  $\Lambda_k$  is the likelihood ratio statistic for testing the null hypothesis against the alternative that a change has occurred between the  $k$ -th population and  $(k+1)$ -th population. He treated the asymptotic approximation of the probability of the first kind of error for the modified likelihood ratio test statistic when the number of the populations  $n$ , and  $m_0$  and  $m_1$  tend to infinity in such a way that

Condition (A): for some  $0 < t_0 < t_1 < 1$ ,  $m_0/m \rightarrow t_0$ ,  $m_1/m \rightarrow t_1$

In this study we take the same modification for the likelihood ratio test. That is,

$$\text{let } T(m_0, m_1) = \max_{1 < m_0 \leq k \leq m_1 < n} T_k$$

and we call this as the modified likelihood ratio test statistic. In the following we say simply that  $n$  tends to infinity if the condition (A) holds, and we consider all the limit theorems under the null hypothesis. Finally it should be also noted that the terminology of "asymptotic equivalence" about two statistic  $F_1$  and  $F_2$  is used for both cases  $F_1 = F_2 + o_p(1)$  and  $F_1 = (1 + o_p(1))F_2$ .

First we have the next limit theorem.

Theorem 1. Let  $T_1(m_0, m_1) = \max_{1 < m_0 \leq k \leq m_1 < n} T_{1k}$ , where

$$T_{1,k} = (N_k/2) \{1 - (S_k/N_k)/(S/N)\}^2 + (N_k^*/2) \{1 - (S_k^*/N_k^*)/(S/N)\}^2.$$

Then  $T(m_0, m_1)$  and  $T_1(m_0, m_1)$  is asymptotically equivalent when  $n$  tends to infinity.

Proof. Without loss of generality we assume  $\sigma_i^2=1$ ,  $i=1, \dots, n$ . Then from Central Limit Theorem, all of  $S/N$ ,  $S_k/N_k$ , and  $S_k^*/N_k^*$  converge to 1 with probability 1. Since  $T_k$  is rewritten as

$$T_k = -N_k \log[1 - \{1 - (S_k/N_k)/(S/N)\}] - N_k^* \log[1 - \{1 - (S_k^*/N_k^*)/(S/N)\}],$$

taking the Tailor expansion of the log function about 1, we obtain

$$T_k = T_{1k} + o_p(1), \quad k = m_0, \dots, m_1.$$

This completes the proof of Theorem 1.

Corollary 1. Let  $T_2(m_0, m_1) = \max_{1 < m_0 \leq k \leq m_1 < n} T_{2k}$ ,

where

$$T_{2k} = (N_k/2) \{(S/N) - (S_k/N_k)\}^2 + (N_k^*/2) \{(S/N) - (S_k^*/N_k^*)\}^2.$$

Then  $T_2(m_0, m_1)$  is asymptotically equivalent to  $T(m_0, m_1)$  when  $n$  tends to infinity.

Proof. This is easy to see if we note that

$$T_1(m_0, m_1) = (S/N)^{-2} T_2(m_0, m_1)$$

and  $(S/N)^{-2}$  converges to 1 with probability 1.

Next simulation results show to what extent the approximations of  $T(m_0, m_1)$  by  $T_1(m_0, m_1)$  and  $T_2(m_0, m_2)$  are good.

Note that the results are based on the 10,000 generations of the

statistics.

Table I

The quantiles of  $T(m_0, m_1)$ ,  $T_1(m_0, m_1)$ , and  $T_2(m_0, m_1)$

n = 50 m <sub>0</sub> = 5 m <sub>1</sub> = 45 v <sub>1</sub> = ... v <sub>n</sub> = 20	%	$T(m_0, m_1)$	$T_1(m_0, m_1)$	$T_2(m_0, m_1)$
	10	1.3258	1.3156	1.31383
30	2.7112	2.17388	2.1685	
50	3.02059	3.0044	3.00169	
60	3.56599	3.559	3.5462	
70	4.23738	4.1963	4.2154	
80	5.14256	5.1173	5.0125	
90	6.6528	6.624	6.6366	
95	8.2040	8.2562	8.35647	

Table II

The quantiles of  $T(m_0, m_1)$ ,  $T_1(m_0, m_1)$ , and  $T_2(m_0, m_1)$

n = 100 m <sub>0</sub> = 10 m <sub>1</sub> = 90 v <sub>1</sub> = ... v <sub>k</sub> = 20	%	$T(m_0, m_1)$	$T_1(m_0, m_1)$	$T_2(m_0, m_1)$
	10	1.459	1.4626	1.4506
30	2.3075	2.3042	2.298	
50	3.2262	3.2176	3.207	
60	3.7859	3.7395	3.7515	
70	4.423	4.4177	4.4198	
80	5.3976	5.3462	5.3417	
90	6.938	6.9399	6.9465	
95	8.4808	8.4511	8.4777	

### 5. Convergence of $T_2(m_0, m_1)$

The purpose of this section is to give the convergence theorem of  $T_2(m_0, m_1)$  to a functional of the standard

Brownian bridge. First we state a well-known lemma.

Lemma 5.1 (Donskar, see Billingsley(1968), p. 137). Let  $\xi_1, \dots, \xi_n$  be identically and independently distributed random variables with mean 0 and finite variance  $\sigma^2$ . Let  $Y_n(t) = S_{[nt]} / \sqrt{n}\sigma$ . Then when  $n$  tends to infinity  $Y_n(t)$  converges to  $W(t)$ ,  $0 \leq t \leq 1$ , weakly, where  $W(t)$  denotes the standard Brownian motion.

Theorem 2. Assume  $v_1 = \dots = v_n = v$ . When  $n$  tends to infinity

$T_2(m_0, m_1)$  converges to  $Z$  in law,

where

$$Z = \max_{t_1 < t < t_2} \{ W_0(t)^2 / t(1-t) \}$$

Proof. Let  $\tilde{S} = S - N$ ,  $\tilde{S}_k = S_k - N_k$  and  $\tilde{S}_{k^*} = S_{k^*} - N_{k^*}$ . Then  $\tilde{S}$ ,  $\tilde{S}_k$ , and  $\tilde{S}_{k^*}$  are the sums of identically and independently distributed random variables with mean 0 and finite variance  $\sqrt{2v}$ , respectively. It is easy to see  $T_{2k}$  is rewritten as

$$\begin{aligned} T_{2k} &= (1/2v) [ (n/k) \{ (k/n) (\tilde{S}/\sqrt{n}) - \tilde{S}_k/\sqrt{n} \}^2 + \\ &\quad (n/k^*) \{ (k^*/n) (\tilde{S}/\sqrt{n}) - \tilde{S}_{k^*}/\sqrt{n} \}^2 ] \\ &= (1/2v) (n/k + n/k^*) \{ (k/n) (\tilde{S}/\sqrt{n}) - \tilde{S}_k/\sqrt{n} \}^2 \end{aligned}$$

Set  $Y_n(t) = \tilde{S}_{[nt]} / \sqrt{n} \sqrt{2v}$ ,  $0 \leq t \leq 1$ . Then  $T_{2k}$  is also rewritten as

$$T_{2k} = (n/k + n/k^*) \{ (k/n) Y_n(1) - Y_n(k/n) \}^2.$$

When  $n$  tends to infinity, since  $Y_n(t)$  converges to  $W(t)$  weakly,  $T_2(m_0, m_1)$  converges to  $Z$  in law, which completes the proof of Theorem 2.

## 6. Evaluation of the limiting distribution of $T(m_0, m_1)$

We have the following evaluation about the limiting

distribution of  $T(m_0, m_1)$ .

Theorem 3.

$$\lim_{n \rightarrow \infty} P\{T(m_0, m_1) \leq c^2\} = \sum_{i=1}^{\infty} \alpha_i(c, q) T^{-\beta_i(c, q)}$$

where  $q=1/2$  and  $T = t_2(1-t_1)/t_1(1-t_2)$

Proof of Theorem 3. This is immediately deduced from Corollary of Theorem 1 and Propositions 2.5 and 2.6.

Theorem 4. For large  $c$ ,

$$\lim_{n \rightarrow \infty} P\{T(m_0, m_1) \geq c^2\} \sim (ce^{-c^2/2}/\sqrt{2\pi})\{(\log T)(1-c^{-2})+2/c^2+O(c^{-4})\}$$

Proof of Theorem 4. This is an immediate consequence of Corollary of Theorem 1 and Propositions 2.5 and 2.7.

Next table III represents the approximations of the upper tail probability by the formula in Theorem 4.

Table III

n	$m_0$	$m_1$	$c^2$	exact P	approximated P
50	5	45	8.2040	0.05	0.07753
100	10	90	8.4808	0.05	0.0688

Note that the used data are parts of Tables I and II. The approximation errors seem to be moderate.

#### References

- [1] Berman, S (1964). Limit theorem for the maximum term in stationary sequences. Ann. Math. Statist. vol. 35, pp. 502-516.
- [2] Berman, S (1974). Sojourns and extremes of Gaussian process.

- Ann. Probability, vol. 2, pp. 999-1026.
- [3] Berman, S (1980). A compound Poisson limit for stationary sums and sojourns of Gaussian processes. Ann. Probability, vol. 8, pp. 511-538.
- [4] Berman, S (1984). Sojourns and extremes of stationary process. Ann. Probability, vol. 10, pp. 1-46.
- [5] Billingsley, P (1968). Convergence of Probability Measures, John Wiley.
- [6] Darling, D.A. and Erdős, P. (1956). A limit theorem for the maximum of normalized sums of independent random variables. Duke Math. J., vol. 23, pp. 143-155.
- [7] DeLong, M (1981). Crossing probabilities for a square root boundary by a Bessel process. Commun. in Statist. Theor. and Meth., vol. 10, pp. 2197-2213.
- [8] Gnedenko, B.V. (1943). Sur la distribution limite du terme d' une serie aleatoire. Ann. Math., vol. 44, pp. 423-453.
- [9] Hawkins, D.M. (1977). Testing a sequence of observations for a shift in location. J. Amer. Statist. Assoc., Vol. 72, pp. 180-186.
- [10] Hsu, D.A. (1979). Detecting shifts of parameter in gamma sequences with applications stock price and air traffic flow analysis. J. Amer. Statist. Assoc., vol. 74, pp. 31-40.
- [11] Leadbetter, M.R., Lindgren, G., and Rootzen, H. (1983). Extremes and related Properties of Random Sequence and processes. Springer Series in Statistics.
- [12] Pickands III, J. (1969). Asymptotic properties of the maximum in a stationary Gaussian process. Trans. Amer. Math. Soc., vol. 145, pp. 75-86.
- [13] Resnick, S. (1987). Extreme values, regular valuation, and point process. Springer, New York.
- [14] Sakata, T. (1988). Detectin a change in variances. Commun. Statist. Theor. and Meth., vol. 17, No., 3, (to appear)
- [15] Sen, A. and Srivastava, M.S. (1975). On tests for detecting change in mean. Ann. Statistics, vol. 3, pp. 98-108..
- [16] Siegmund, D. (1986). Boundary crossing probabilities and its

statistical applications. Ann. Statist. vol. 14, pp. 361-404.

- [17] Srivastava, M.S. and Worsley, K.J.(1986). Likelihood ratio tests for a change in the multivariate normal mean. J. Amer Statist. Assoc., Vol. 81, pp. 199-204.
- [18] Worsley, K.J. (1986). Confidence regions and tests for a change point in a sequence of exponential family random variables. Biometrika, vol. 73, pp. 91-104.
- [19] Yao, Y.C. and Davis, R.A.(1986). The asymptotic behavior of the likelihood ratio statistic for testing a shift in mean in a sequence of independent normal variates. Sankhya, vol. 48, pp. 339-353.