

On characterization of primitive commutative association schemes by a nonsymmetric relation with small valency

Mitsugu Hirasaka* (平坂 貢)
Graduate School of Mathematics
Kyushu University†

May 18, 1998

Abstract

Let (X, G) be a primitive commutative association scheme with a nonsymmetric relation of valency 4. Then, for each nonsymmetric relation g of valency 4, the graph (X, g) is uniquely determined. In particular, the cardinality of X is the cube of an odd prime.

Let (X, G) be an association scheme (see [4]). Then for each $g \in G$, (X, g) is a regular graph satisfying certain conditions. It seems an interesting problem to determine whether a given graph could be a relation of an association scheme. This problem has been studied before and will be one of the main topics in characterizing association schemes by certain intersection numbers. Until now, quite a few results for this problem under various assumptions or interests, for instance, when (X, G) is a P -polynomial association scheme or a translation association scheme (see [4] or [6]), or when we impose conditions on valencies or cardinality of X , are obtained by many researchers. The author is especially interested in association schemes with a nonsymmetric relation, namely aims to determine the directed graph (X, g) with $g \in G$ or the structure of (X, G) under some assumptions, for example, when (X, G) is primitive, commutative or when $|X|$ is a prime.

The following is an example of such association schemes.

*e-mail address: hirasaka@math.kyushu-u.ac.jp

†Department of Science, Hakozaki 6-10-1, Fukuoka 812, Japan

Example 0.1 Let F_p be a finite field of order p where p is an odd prime. We define some permutations on F_p^n as follows;

- o Let S_n be the symmetric group of degree n . For all $(x_1, x_2, \dots, x_n) \in F_p^n$ and $\sigma \in S_n$,

$$(x_1, x_2, \dots, x_n)^\sigma := (x_{\sigma^{-1}(1)}, x_{\sigma^{-1}(2)}, \dots, x_{\sigma^{-1}(n)}).$$

- o For each $(x_1, x_2, \dots, x_n) \in F_p^n$, we define $\tau \in GL(F_p^n)$ as follows;

$$(x_1, x_2, \dots, x_n)^\tau := (-x_1, x_2 - x_1, \dots, x_n - x_1).$$

- o For all $u, v \in F_p^n$,

$$u^v = u + v.$$

Then the permutation group $\Gamma := \langle S_n, \tau, F_p^n \rangle$ acts on F_p^n transitively, and Γ acts on $F_p^n \times F_p^n$ by $(u, v)^g := (u^g, v^g)$ where $(u, v) \in F_p^n \times F_p^n$ and $g \in \Gamma$. Let G be the set of orbits of Γ on $F_p^n \times F_p^n$. Then (F_p^3, G) is an association scheme (see [4, p.52]), and it can be verified that (F_p^n, G) is a primitive commutative association scheme with a nonsymmetric relation of valency $n + 1$. Indeed, the orbit containing $((0, 0, \dots, 0), (1, 0, \dots, 0))$ is a nonsymmetric relation of valency $n + 1$.

Let (X, G) be a primitive commutative association scheme. The following results are obtained until now.

- 1) If there exists a nonsymmetric relation $g \in G$ of valency 1 then (X, g) is a directed cycle of prime length.
- 2) There exists (X, G) with a nonsymmetric relation of valency 2.
- 3) If there exists a nonsymmetric relation $g \in G$ of valency 3 then we have one of the following;
 - (a) (X, g) is isomorphic to a relation of a cyclotomic scheme where $|X|$ is a prime.
 - (b) There exists a relation $\tilde{g} \in G$ isomorphic to a relation of a cyclotomic scheme where $|X|$ is an odd prime squared.

Although it is trivial to prove 1), 2), it is rather difficult to prove 3) (see [10]).

The following is a main result of this presentation.

Theorem 0.2 *Let (X, G) be a primitive commutative association scheme with a nonsymmetric relation of valency 4, denoted by g . Then the graph (X, g) is isomorphic to (F_p^3, E) where $(x, y) \in E$ if and only if $y - x \in \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (-1, -1, -1)\}$.*

We need certain intersection numbers in order to determine the graph (X, g) . The following proposition holds.

Proposition 0.3 *For each $g \in G$ with $k_g = 4$ and $g^* \neq g$, there exist $a, b, f, h, m \in G$ such that*

- (i) $g \bullet g = 2a + b, k_a = 6, k_b = 4.$
- (ii) $g \bullet g^* = 4 \cdot 1_X + h, k_h = 12.$
- (iii) $a^* \bullet a = 6 \cdot 1_X + 2h + f, k_f = 6.$
- (iv) $a = a^*, g \bullet a = 3g^* + m, k_m = 12.$
- (v) $ga \cap gb = \{m\}.$

Definition 0.4 Let $e \in G$ be such that $k_e = 4$ and $e \neq e^*$. A sequence (x_0, x_1, \dots, x_j) of elements of X is a *chain* if $(x_i, x_{i+1}) \in g$ for each i with $0 \leq i \leq j-1$, and $(x_i, x_{i+2}) \in b$ for each i with $0 \leq i \leq j-2$.

Remark 0.5 *If (x_0, x_1, \dots, x_i) is a chain then there exists a unique element $x_{i+1} \in X$ such that $(x_{i-1}, x_{i+1}) \in b$ and $(x_i, x_{i+1}) \in g$ by $p_{bg}^g = 1$, and $(x_0, x_1, \dots, x_i, x_{i+1})$ is also a chain. Hence we obtain a chain of length i for any $i > 0$, and whence there exists a closed chain since $|X|$ is finite.*

Let $(x_0, x_1, \dots, x_n = x_0)$ be a closed chain of the shortest length. We would like to show that n is a prime and identify $(x_0, x_1, \dots, x_n = x_0)$ with F_p . Proposition 0.3 makes the above things possible. Moreover, we can construct F_p^3 by repeating to connect a closed chain $(y_0, y_1, \dots, y_n = y_0)$ to another closed chain $(y'_0, y'_1, \dots, y'_n = y'_0)$ such that $(y_i, y'_i) \in g$ for all i with $0 \leq i \leq n-1$ and $y'_0 \neq y_1$.

References

- [1] Z. Arad, E. Fisman, V. Miloslavsky, M. Muzychuk, On antisymmetric homogeneous integer table algebras, preprint.
- [2] Z. Arad, H. I. Blau, On table algebras and applications to finite group theory, *J. Algebra* 138, 137–185 (1991).

- [3] Z. Arad, E. Fisman, H. Arisha, M. Muzychuk, On integral table algebras with non-real element of degree 4, 1998, in preparation.
- [4] E. Bannai, T. Ito, Algebraic Combinatorics I: Association Schemes, Benjamin/Cummings, Menlo Park, CA, 1984.
- [5] H. I. Blau, Integral table algebras, affine diagrams, and the analysis of degree two, *J. Algebra* 178, 872–918 (1995).
- [6] A. E. Brouwer, A. M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, 1989.
- [7] C. D. Godsil, Algebraic Combinatorics, Chapman and Hall, New York, 1993.
- [8] M. Hall, Jr. Combinatorial Theory, second edition, A Wiley-Interscience Publication.
- [9] A. Hanaki, I. Miyamoto, Classification of association schemes with 18 and 19 vertices, 1998, preprint.
- [10] M. Hirasaka, Association schemes with a prime number of points, 1997, preprint.
- [11] E. Nomiyama, Classification of association schemes with at most ten vertices, *Kyushu J. of Math.* 49, 163–195 (1995).
- [12] N. Yamazaki, On symmetric association schemes with $k_1 = 3$, 1995, preprint.