

N-Fractional Calculus of Some Irrational Functions

Katsuyuki Nishimoto

Institute for Applied Mathematics, Descartes Press Co.
2-13-10 Kaguike, Koriyama, 963-8833, JAPAN.

Keywords ; Fractional Calculus, N-Fractional Calculus (NFC),
NFC of Products, NFC of Composite Functions

Abstract

In this article N-fractional calculus of the functions (which have multiple root signs)

$$f(z) = \sqrt{\sqrt{z-b} - c - d}$$

are discussed.

Theorem 1. We have

$$(i) \quad (f(z))_{\gamma} = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \\ \times \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m, \quad \left(\left| \frac{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \right| < \infty \right)$$

and

$$(ii) \quad (f(z))_n = (-1)^n (z-b)^{(1/8)-n} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k [\frac{k}{2} - \frac{1}{8} + \frac{m}{4}]_n}{m! \cdot k!} \\ \times \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m, \quad (n \in \mathbb{Z}_0^+)$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

§ 0. Introduction (Definition of Fractional Calculus)

(I) Definition. (by K. Nishimoto) ([1] Vol. 1)

Let $D = \{D_-, D_+\}$, $C = \{C_-, C_+\}$,

C_- be a curve along the cut joining two points z and $-\infty + i\operatorname{Im}(z)$,

C_+ be a curve along the cut joining two points z and $\infty + i\operatorname{Im}(z)$,

D_- be a domain surrounded by C_- , D_+ be a domain surrounded by C_+ .

(Here D contains the points over the curve C).

Moreover, let $f = f(z)$ be a regular function in $D(z \in D)$,

$$f_v = (f)_v - {}_C(f)_v = \frac{\Gamma(v+1)}{2\pi i} \int_C \frac{f(\xi)}{(\xi-z)^{v+1}} d\xi \quad (v \notin \mathbb{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{v \rightarrow -m} (f)_v \quad (m \in \mathbb{Z}^+), \quad (2)$$

where $-\pi \leq \arg(\xi - z) \leq \pi$ for C_- , $0 \leq \arg(\xi - z) \leq 2\pi$ for C_+ ,

$\xi \neq z$, $z \in C$, $v \in R$, Γ ; Gamma function,

then $(f)_v$ is the fractional differintegration of arbitrary order v (derivatives of order v for $v > 0$, and integrals of order $-v$ for $v < 0$), with respect to z , of the function f , if $|(f)_v| < \infty$.

(II) On the fractional calculus operator N^v [3]

Theorem A. Let fractional calculus operator (Nishimoto's Operator) N^v be

$$N^v = \left(\frac{\Gamma(v+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{v+1}} \right) \quad (v \notin \mathbb{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with

$$N^{-m} = \lim_{v \rightarrow -m} N^v \quad (m \in \mathbb{Z}^+), \quad (4)$$

and define the binary operation \circ as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta(N^\alpha f) \quad (\alpha, \beta \in R), \quad (5)$$

then the set

$$\{N^v\} = \{N^v | v \in R\} \quad (6)$$

is an Abelian product group (having continuous index v) which has the inverse transform operator $(N^v)^{-1} = N^{-v}$ to the fractional calculus operator N^v , for the function f such that $f \in F = \{f; 0 \neq |f_v| < \infty, v \in R\}$, where $f = f(z)$ and $z \in C$. (vis. $-\infty < v < \infty$).

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

(For our convenience, we call $N^\beta \circ N^\alpha$ as product of N^β and N^α .)

Theorem B. " F.O.G. $\{N^\nu\}$ " is an " Action product group which has continuous index ν " for the set of F . (F.O.G. ; Fractional calculus operator group) [3]

(III) **Lemma.** We have [1]

$$(i) \quad ((z-c)^b)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-b)}{\Gamma(-b)} (z-c)^{b-\alpha} \quad \left(\left| \frac{\Gamma(\alpha-b)}{\Gamma(-b)} \right| < \infty \right), \quad (7)$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty), \quad (8)$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty), \quad (9)$$

where $z-c \neq 0$ for (i) and $z-c \neq 0, 1$ for (ii), (iii) ,

§ 1. Preliminary

[I] The theorem below is reported by K. Nishimoto already (cf. J. Frac. Calc. Vol. 29, May (2006), p.37). [12]

Theorem D. We have

$$(i) \quad (((z-b)^\beta - c)^\alpha)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha\beta-\gamma} \\ \times \sum_{k=0}^{\infty} \frac{[-\alpha]_k \Gamma(\beta k - \alpha\beta + \gamma)}{k! \Gamma(\beta k - \alpha\beta)} \left(\frac{c}{(z-b)^\beta} \right)^k \quad \left(\left| \frac{\Gamma(\beta k - \alpha\beta + \gamma)}{\Gamma(\beta k - \alpha\beta)} \right| < \infty \right) \quad (1)$$

and

$$(ii) \quad (((z-b)^\beta - c)^\alpha)_n = (-1)^n (z-b)^{\alpha\beta-n} \\ \times \sum_{k=0}^{\infty} \frac{[-\alpha]_k [\beta k - \alpha\beta]_n}{k!} \left(\frac{c}{(z-b)^\beta} \right)^k \quad (n \in \mathbb{Z}_0^+) \quad (2)$$

where

$$|c/(z-b)^\beta| < 1,$$

and

$$[\lambda]_k = \lambda(\lambda+1)\cdots(\lambda+k-1) = \Gamma(\lambda+k)/\Gamma(\lambda) \quad \text{with} \quad [\lambda]_0 = 1.$$

(Notation of Pochhammer).

[I I] The theorem below is reported by K. Nishimoto already (cf. J. Frac. Calc. Vol. 31, May (2007), p.13). [13]

Theorem E. We have

$$(i) (((z-b)^\beta - c)^\alpha - d)^\delta)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha\beta\delta-\gamma}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k \Gamma(\beta k - \alpha\beta(\delta-m) + \gamma)}{m! \cdot k! \Gamma(\beta k - \alpha\beta(\delta-m))} \left(\frac{c}{(z-b)^\beta} \right)^k \left(\frac{d}{(z-b)^{\alpha\beta}} \right)^m$$

$$\left(\left| \frac{\Gamma(\beta k - \alpha\beta(\delta-m) + \gamma)}{\Gamma(\beta k - \alpha\beta(\delta-m))} \right| < \infty \right) \quad (3)$$

and

$$(ii) (((z-b)^\beta - c)^\alpha - d)^\delta)_n = (-1)^n (z-b)^{\alpha\beta\delta-n}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k [\beta k - \alpha\beta(\delta-m)]_n}{m! \cdot k!} \left(\frac{c}{(z-b)^\beta} \right)^k \left(\frac{d}{(z-b)^{\alpha\beta}} \right)^m$$

$$(n \in \mathbb{Z}_0^+) \quad (4)$$

where

$$((z-b)^\beta - c)^\alpha - d \neq 0, \quad |c/(z-b)^\beta| < 1, \quad |d/(z-b)^{\alpha\beta}| < 1.$$

§ 2. N-Fractional Calculus of Functions

$$\sqrt{\sqrt{\sqrt{z-b} - c} - d}$$

Theorem 1. We have

$$(i) \left(\sqrt{\sqrt{\sqrt{z-b} - c} - d} \right)_\gamma = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (1)$$

$$\left(\left| \frac{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{\Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \right| < \infty \right)$$

and

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b} - c} - d} \right)_n = (-1)^n (z-b)^{(1/8)-n} \\ \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k [\frac{k}{2} - \frac{1}{8} + \frac{m}{4}]_n}{m! \cdot k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2) \\ (n \in \mathbb{Z}_0^+)$$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

Proof of (i). We have

$$\sqrt{\sqrt{\sqrt{z-b} - c} - d} = \left(((z-b)^{1/2} - c)^{1/2} - d \right)^{1/2}, \quad (3)$$

hence, operating N^γ to the both sides of (3), we obtain

$$\left(\sqrt{\sqrt{\sqrt{z-b} - c} - d} \right)_\gamma = \left(\left(((z-b)^{1/2} - c)^{1/2} - d \right)^{1/2} \right)_\gamma \quad (4)$$

$$= e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \\ \times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4} + \gamma)}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (5)$$

setting $\alpha = \beta = \delta = 1/2$ in Theorem E. (i) in preliminary, under the conditions stated before.

Proof of (ii). Set $\gamma = n$ in (i).

Corollary 1. We have

$$(i) \quad \left(\sqrt{\sqrt{\sqrt{z-b} - c}} \right)_\gamma = e^{-i\pi\gamma} (z-b)^{(1/8)-\gamma} \\ \times \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{1}{8} + \gamma)}{k! \Gamma(\frac{k}{2} - \frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (6) \\ \left(\left| \frac{\Gamma(\frac{k}{2} - \frac{1}{8} + \gamma)}{\Gamma(\frac{k}{2} - \frac{1}{8})} \right| < \infty \right)$$

Katsuyuki Nishimoto

and

$$(i) \quad \left(\sqrt{\sqrt{z-b} - c} \right)_n = (-1)^n (z-b)^{(1/8)-n} \times \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k [\frac{k}{2} - \frac{1}{8}]_n}{k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (7)$$

$(n \in \mathbb{Z}_0^+)$

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1 .$$

Proof of (i). Set $d = 0$ in Theorem 1(i).

We have (6) from Theorem D, (i), setting $\beta = 1/2$ and $\alpha = 1/4$.

Proof of (ii). Set $\gamma = n$ in (i).

Corollary 2. We have

$$(i) \quad \left(\sqrt{\sqrt{z-b}} \right)_\gamma = e^{-i\pi\gamma} \frac{\Gamma(-\frac{1}{8} + \gamma)}{\Gamma(-\frac{1}{8})} (z-b)^{(1/8)-\gamma} \quad (8)$$

$(|\Gamma(-\frac{1}{8} + \gamma)| < \infty)$

and

$$(ii) \quad \left(\sqrt{\sqrt{z-b}} \right)_n = (-1)^n [-\frac{1}{8}]_n (z-b)^{(1/8)-n} \quad (9)$$

$(n \in \mathbb{Z}_0^+)$

where

$$z-b \neq 0 .$$

Proof of (i). Set $c = 0$ in Corollary 1(i).

Or set $c = d = 0$ in Theorem 1.(i).

Proof of (ii). Set $\gamma = n$ in (i).

Note. We have

$$\left(\sqrt{\sqrt{z-b}} \right)_\gamma = ((z-b)^{1/8})_\gamma = e^{-i\pi\gamma} \frac{\Gamma(-\frac{1}{8} + \gamma)}{\Gamma(-\frac{1}{8})} (z-b)^{(1/8)-\gamma} \quad (10)$$

from Lemma (i), directly.

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

§ 3. Semi Derivatives and Integrals

Theorem 2. We have

$$(i) \quad \left(\sqrt{\sqrt{z-b} - c} - d \right)_{1/2} = -i(z-b)^{-3/8}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} + \frac{3}{8} + \frac{m}{4})}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (1)$$

(semi derivatives)

and

$$(ii) \quad \left(\sqrt{\sqrt{z-b} - c} - d \right)_{-1/2} = i(z-b)^{5/8}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{5}{8} + \frac{m}{4})}{m! \cdot k! \Gamma(\frac{k}{2} - \frac{1}{8} + \frac{m}{4})} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2)$$

(semi integrals)

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1, \quad \left| \frac{d}{\sqrt[4]{z-b}} \right| < 1.$$

Proof. Set $\gamma = 1/2$ and $-1/2$ in Theorem 1(i) we have then (i) and (ii) respectively, clearly.

Corollary 3. We have

$$(i) \quad \left(\sqrt{\sqrt{z-b} - c} \right)_{1/2} = -i(z-b)^{-3/8} \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2} + \frac{3}{8})}{k! \Gamma(\frac{k}{2} - \frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (3)$$

(semi derivatives)

and

$$(ii) \quad \left(\sqrt{\sqrt{z-b} - c} \right)_{-1/2} = i(z-b)^{5/8} \sum_{k=0}^{\infty} \frac{[-\frac{1}{4}]_k \Gamma(\frac{k}{2} - \frac{5}{8})}{k! \Gamma(\frac{k}{2} - \frac{1}{8})} \left(\frac{c}{\sqrt{z-b}} \right)^k \quad (4)$$

(semi integrals)

where

$$\left| \frac{c}{\sqrt{z-b}} \right| < 1.$$

Katsuyuki Nishimoto

Proof. Set $d = 0$ in Theorem 2.

Corollary 4. We have

$$(i) \quad \left(\sqrt{\sqrt{z-b} - d} \right)_{1/2} = -i(z-b)^{-3/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m \Gamma(\frac{3}{8} + \frac{m}{4})}{m!! \Gamma(-\frac{1}{8} + \frac{m}{4})} \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (5)$$

(semi derivatives)

and

$$(ii) \quad \left(\sqrt{\sqrt{z-b} - d} \right)_{-1/2} = i(z-b)^{5/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m \Gamma(-\frac{5}{8} + \frac{m}{4})}{m! \Gamma(-\frac{1}{8} + \frac{m}{4})} \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \quad (2)$$

(semi integrals)

where

$$\left| \frac{d}{\sqrt[4]{z-b}} \right| < 1 .$$

Proof. Set $c = 0$ in Theorem 2.

§ 4. Some Special Cases

[I] When $n = 0$, we have

$$\begin{aligned} \left(\sqrt{\sqrt{z-b} - c - d} \right)_0 &= (z-b)^{1/8} \\ &\times \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k}{m! k!} \left(\frac{c}{\sqrt{z-b}} \right)^k \left(\frac{d}{\sqrt[4]{z-b}} \right)^m \end{aligned} \quad (1)$$

from § 2. (2).

Now we have

$$\begin{aligned} \text{RHS of (1)} &= (z-b)^{1/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \\ &\quad \left(S = \frac{c}{\sqrt{z-b}}, \quad T = \frac{d}{\sqrt[4]{z-b}} \right) \end{aligned} \quad (2)$$

$$= (z-b)^{1/8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m (1-S)^{(1/4)-(m/2)} \quad (3)$$

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

$$= (z - b)^{1/8} (1 - S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m T^m}{m! (1 - S)^{m/2}} \quad (4)$$

$$= (z - b)^{1/8} (1 - S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} \left(\frac{d}{\sqrt{\sqrt{z - b} - c}} \right)^m \quad (5)$$

$$= (z - b)^{1/8} \left(1 - \frac{c}{\sqrt{z - b}} \right)^{1/4} \left(1 - \frac{d}{\sqrt{\sqrt{z - b} - c}} \right)^{1/2} \quad (6)$$

$$= \sqrt{\sqrt{\sqrt{z - b} - c} - d} \quad (\text{LHS of (1)}). \quad (7)$$

[II] When $n = 1$, we have

$$\left(\sqrt{\sqrt{\sqrt{z - b} - c} - d} \right)_1 = -(z - b)^{-7/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k [\frac{k}{2} + \frac{m}{4} - \frac{1}{8}]_1}{m! \cdot k!} S^k T^m \quad (8)$$

$$= -(z - b)^{-7/8} \sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (\frac{k}{2} + \frac{m}{4} - \frac{1}{8})}{m! \cdot k!} S^k T^m \quad (9)$$

from § 2. (2).

Now we have the identities;

$$\sum_{k=0}^{\infty} \frac{[\lambda]_k}{k!} z^k = (1 - z)^{-\lambda}, \quad (10)$$

$$[\lambda]_0 = 1, \quad [\lambda]_1 = \lambda, \quad (11)$$

$$[\lambda]_{k+1} = \lambda [\lambda + 1]_k, \quad (12)$$

hence

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{[\lambda]_k k}{k!} z^k &= \sum_{k=1}^{\infty} \frac{[\lambda]_k}{(k-1)!} z^k = z \sum_{k=0}^{\infty} \frac{[\lambda]_{k+1}}{k!} z^k \\ &= \lambda z (1 - z)^{-\lambda-1}. \end{aligned} \quad (13)$$

Therefore, we have

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (\frac{k}{2})}{m! \cdot k!} S^k T^m = \frac{1}{2} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k k}{k!} S^k \quad (14)$$

Katsuyuki Nishimoto

$$= \frac{1}{2} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \left(\frac{m}{2} - \frac{1}{4} \right) S \sum_{k=0}^{\infty} \frac{[\frac{m}{2} + \frac{3}{4}]_k}{k!} S^k \quad (15)$$

$$= \frac{1}{2} S(1-S)^{-3/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m (\frac{m}{2} - \frac{1}{4})}{m!} U^m \quad \left(U = \frac{d}{\sqrt{\sqrt{z-b}-c}} \right) \quad (16)$$

$$= \frac{1}{2} S(1-S)^{-3/4} \left\{ -\frac{1}{4} U(1-U)^{-1/2} - \frac{1}{4} (1-U)^{1/2} \right\} \quad (17)$$

$$= -\frac{1}{8} S(1-S)^{-3/4} (1-U)^{-1/2}, \quad (18)$$

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (\frac{m}{4})}{m! k!} S^k T^m = \frac{1}{4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \quad (19)$$

$$= \frac{1}{4} (1-S)^{1/4} \sum_{m=1}^{\infty} \frac{[-\frac{1}{2}]_m}{(m-1)!} U^m \quad (20)$$

$$= \frac{1}{4} (1-S)^{1/4} U \left(-\frac{1}{2} \right) \sum_{m=0}^{\infty} \frac{[\frac{1}{2}]_m}{m!} U^m \quad (21)$$

$$= -\frac{1}{8} (1-S)^{1/4} U (1-U)^{-1/2}. \quad (22)$$

and

$$\sum_{m,k=0}^{\infty} \frac{[-\frac{1}{2}]_m [\frac{m}{2} - \frac{1}{4}]_k (-\frac{1}{8})}{m! k!} S^k T^m = -\frac{1}{8} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} T^m \sum_{k=0}^{\infty} \frac{[\frac{m}{2} - \frac{1}{4}]_k}{k!} S^k \quad (23)$$

$$= -\frac{1}{8} (1-S)^{1/4} \sum_{m=0}^{\infty} \frac{[-\frac{1}{2}]_m}{m!} U^m \quad (24)$$

$$= -\frac{1}{8} (1-S)^{1/4} (1-U)^{1/2}. \quad (25)$$

Then applying (18), (22) and (25) to (9) we obtain

$$\left(\sqrt{\sqrt{z-b}-c-d} \right)_1 = \frac{1}{8} (z-b)^{-7/8} (1-S)^{1/4} (1-U)^{-1/2} \{ S(1-S)^{-1} + 1 \} \quad (26)$$

N- Fractional Calculus of Some Functions Which Have Multiple Root Signs

$$= \frac{1}{8} (z - b)^{-7/8} (1 - S)^{1/4} (1 - U)^{-1/2} (1 - S)^{-1} \quad (27)$$

$$= \frac{1}{8} (z - b)^{-7/8} (1 - S)^{-3/4} (1 - U)^{-1/2} \quad (28)$$

$$= \frac{1}{8} (z - b)^{-1/2} (\sqrt{z - b} - c)^{-1/2} (\sqrt{\sqrt{z - b} - c} - d)^{-1/2} . \quad (29)$$

This result (29) coincides with the one obtained by classical calculus.

References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator N^ν (On an action group), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives (A serendipity in fractional calculus), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N-Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto ; On the fractional calculus $(a - z)^\beta$ and $\log(a - z)$, J. Frac. Calc. Vol.3, May (1993), 19 - 27.
- [7] K. Nishimoto and S.- T. Tu ; Fractional calculus of Psi functions (Generalized Polygamma unctions), J. Frac. Calc. Vol.5 May (1994), 27 - 34.
- [8] S.- T. Tu and K. Nishimoto ; On the fractional calculus of functions $(cz - a)^\beta$ $\log(cz - a)$, J. Frac.Calc.Vol.5, May (1994), 35 - 43.
- [9] K. Nishimoto ; N-Fractional Calculus of the Power and Logarithmic Functions and Some Identities, J. Frac. Calc. Vol.21, May (2002), 1 - 6.
- [10] K. Nishimoto ; Some Theorems for N-Fractional Calculus of Logerithmic Functions I , J. Frac Calc.Vol.21, May (2002), 7 - 12.
- [11] K. Nishimoto ; N-Fractional Calculus of Products of Some Power Functions, J. Frac.Calc. Vol.27, May (2005), 83 - 88.
- [12] K. Nishimoto ; N-Fractional Calculus of Some Composite Functions, J. Frac. Calc. Vol. 29, May (2006), 35 - 44.
- [13] K. Nishimoto ; N-Fractional Calculus of Some Composite Algebraic Functions,J. Frac.Calc. Vol. 31, May (2006), 11 - 23.
- [14] K. Nishimoto ; N-Fractional Calculus of Some Elementary Functions and Their Semi Differintegrations, J. Frac. Caic. Vol. 31, May (2007), 1 - 10.
- [15] K. Nishimoto and T. Miyakoda ; N-Fractional Calculus and n -th Derivatives of Some Algebraic Functions, J. Frac. Calc. Vol. 31, May (2007), 53 - 62.

Katsuyuki Nishimoto

- [16] T. Miyakoda ; N- Fractional Calculus of Certain Algebraic Functions, J. Frac. Calc. Vol. 31, May (2007), 63 - 76.
- [17] David Dunmmit and Richard M. Foote ; Abstract Algebra, Prentice Hall (1991).
- [18] K. B. Oldham and J. Spanier ; The Fractional Calculus, Academic Press (1974).
- [19] A.C. McBride ; Fractional Calculus and Integral Transforms of Generalized Functions, Research Notes, Vol. 31, (1979), Pitman.
- [20] S.G. Samko, A.A. Kilbas and O.I. Marichev ; Fractional Integrals and Derivatives, and Some Their Applications (1987), Naoka, USSR.
- [21] K.S. Miller and B. Ross ; An Introduction to The Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, (1993).
- [22] V. Kiryakova ; Generalized fractional calculus and applications, Pitman Research Notes, No.301, (1994), Longman.
- [23] A.Carpinteri and F. Mainardi (Ed.) ; Fractals and Fractional Calculus in Continuum Mechanics (1997), Springer, Wien, New York.
- [24] Igor Podlubny ; Fractional Differential Equations (1999), Academic Press.
- [25] R. Hilfer (Ed.) ; Applications of Fractional Calculus in Physics, (2000), World Scientific, Singapor, New Jersey, London, Hong Kong.
- [26] A.P. Prudnikov, Yu. A. Bryckov and O.I. Marichev ; Integrals and Series, Vol. I, Gordon and Breach, New York, (1986).
- [27] S. Moriguchi, K.Udagawa and S. Hitotsumatsu ; Mathematical Formulae, Vol.2, Iwanami Zensho, (1957), Iwanami, Japan.

Katsuyuki Nishimoto
Institute for Applied Mathematics
Descartes Press Co.
2 - 13 - 10 Kaguike, Koriyama
963 - 8833 Japan