

On the Interaction of a Pair of Coaxial Vortex Rings

Masashi AIKI

Department of Mathematics,
Faculty of Science and Technology, Tokyo University of Science

1 Introduction

We are interested in the mathematical analysis of the interaction of two vortex rings sharing a common axis of symmetry (coaxial vortex rings) in an incompressible and inviscid fluid. A vortex ring is a thin torus-shaped region in the fluid in which the vorticity of the fluid is concentrated. The study of such interaction dates back to 1858, where in his seminal paper Helmholtz [1] observed that a pair of vortex rings may exhibit what is now known as “leapfrogging”. Leapfrogging is a motion pattern where two vortex rings pass through each other repeatedly due to the induced flow of the rings acting on each other. Under the classical definition of leapfrogging motion, the pair as a whole also moves in one direction along the common axis of symmetry. This is the case we focus on in this paper. Dyson [2, 3] also considered the motion of coaxial vortex rings and proposed a system of ordinary differential equations describing such motion. Based on this model system, Dyson also observed that leapfrogging may occur. The complex, yet tangible nature of the leapfrogging phenomenon fascinated many researchers and since the observation by Helmholtz, leapfrogging of a pair of coaxial vortex rings as well as the interaction of coaxial vortex rings in general are well studied theoretically, numerically, and experimentally. Notably, although the leapfrogging phenomenon was theoretically observed for a long time, the first experiment which successfully provided photographic proof of the leapfrogging phenomenon in a laboratory setting was the one conducted by Yamada and Matsui [16] in 1978. They used vortex rings made of air and used smoke for visualization and successfully created a leapfrogging pair of rings.

In more recent years, Borisov, Kilin, and Mamaev [20] gave a thorough description of the possible motion patterns of two interacting vortex rings moving under Dyson’s model. Hence, much is already known for Dyson’s model, but the model has one drawback in that it is derived as a system of ordinary differential equations for the radius and the

displacement along the common axis of the rings. It is observed by Maxworthy [31], Widnall and Tsai [32], Widnall and Sullivan [33], and Fukumoto and Hattori [34], that even a small perturbation which destroys the axisymmetry of a vortex ring can grow and eventually cause instability (this kind of instability is called the curvature instability by Fukumoto and Hattori). This suggests that when considering the motion of vortex rings, it is important to model the motion within a framework which can incorporate the effects of these kind of perturbations in order to further understand the behavior of a pair of coaxial vortex rings, but this is not possible under Dyson's model.

Given these situations, the author proposed a new model describing the interaction of coaxial vortex rings. In particular, the new model is a system of partial differential equations which can incorporate the effects of non-symmetric perturbations.

The rest of the paper is organized as follows. In Section 2, we introduce the model system of equations which will be considered in this paper. The system describes the interaction of two vortex filaments with general shape. In Section 3, we consider the case when the two filaments are circular with a common axis of symmetry and the vorticity strengths have the same sign. We show that the problem can be reduced to a two-dimensional Hamiltonian system. From here, we give a condition for leapfrogging to occur, and prove that the condition is necessary and sufficient. The precise statement will be given in the beginning of Section 3. In Section 4, we focus on another type of motion pattern that is observed in the real world. Namely, we prove that the model system is also capable of describing the head-on collision of vortex rings.

2 Interaction of Two Vortex Filaments

We consider the interaction of two vortex filaments. The author proposed the following system of partial differential equations.

$$\begin{cases} \mathbf{X}_t = \Gamma_1 \frac{\mathbf{X}_\xi \times \mathbf{X}_{\xi\xi}}{|\mathbf{X}_\xi|^3} - \alpha \Gamma_2 \frac{\mathbf{Y}_\xi \times (\mathbf{X} - \mathbf{Y})}{|\mathbf{X} - \mathbf{Y}|^3}, \\ \mathbf{Y}_t = \Gamma_2 \frac{\mathbf{Y}_\xi \times \mathbf{Y}_{\xi\xi}}{|\mathbf{Y}_\xi|^3} - \alpha \Gamma_1 \frac{\mathbf{X}_\xi \times (\mathbf{Y} - \mathbf{X})}{|\mathbf{X} - \mathbf{Y}|^3}, \end{cases} \quad (2.1)$$

where $\mathbf{X}(\xi, t) = {}^t(X_1(\xi, t), X_2(\xi, t), X_3(\xi, t))$ and $\mathbf{Y}(\xi, t) = {}^t(Y_1(\xi, t), Y_2(\xi, t), Y_3(\xi, t))$ are the position of the two filaments, parametrized by ξ at time t , \times is the exterior product in the three-dimensional Euclidean space, $\Gamma_1, \Gamma_2 \in \mathbf{R} \setminus \{0\}$ are the vorticity strengthes of the filaments \mathbf{X} and \mathbf{Y} respectively, and $\alpha > 0$ is a real parameter introduced in the derivation of the model.

The purpose of this paper is to prove that system (2.1) is capable of describing the leapfrogging phenomenon. More precisely, we give a necessary and sufficient condition for a pair of coaxial vortex ring to exhibit leapfrogging.

3 Leapfrogging for a Pair of Filaments with Vorticity Strengths of the Same Sign

We consider the case when the two filaments are arranged as coaxial circles and $\Gamma_1, \Gamma_2 > 0$. Rescaling the time variable by a factor of Γ_2 in (2.1) yields

$$\begin{cases} \mathbf{X}_t = \beta \frac{\mathbf{X}_\xi \times \mathbf{X}_{\xi\xi}}{|\mathbf{X}_\xi|^3} - \alpha \frac{\mathbf{Y}_\xi \times (\mathbf{X} - \mathbf{Y})}{|\mathbf{X} - \mathbf{Y}|^3}, \\ \mathbf{Y}_t = \frac{\mathbf{Y}_\xi \times \mathbf{Y}_{\xi\xi}}{|\mathbf{Y}_\xi|^3} - \alpha\beta \frac{\mathbf{X}_\xi \times (\mathbf{Y} - \mathbf{X})}{|\mathbf{X} - \mathbf{Y}|^3}, \end{cases} \quad (3.1)$$

where $\beta = \Gamma_1/\Gamma_2$. We assume without loss of generality that $\beta \geq 1$, since the case $\beta < 1$ is reduced to the case $\beta > 1$ by renaming the filaments.

Suppose that for some $R_{1,0}, R_{2,0} > 0$ and $z_{1,0}, z_{2,0} \in \mathbf{R}$, the initial filaments \mathbf{X}_0 and \mathbf{Y}_0 are parametrized by $\xi \in [0, 2\pi)$ as follows.

$$\mathbf{X}_0(\xi) = {}^t(R_{1,0} \cos(\xi), R_{1,0} \sin(\xi), z_{1,0}), \quad \mathbf{Y}_0(\xi) = {}^t(R_{2,0} \cos(\xi), R_{2,0} \sin(\xi), z_{2,0}),$$

where we assume that $(R_{1,0} - R_{2,0})^2 + (z_{1,0} - z_{2,0})^2 > 0$, which means that the two circles are not overlapping. Now, we make the ansatz

$$\mathbf{X}(\xi, t) = {}^t(R_1(t) \cos(\xi), R_1(t) \sin(\xi), z_1(t)), \quad \mathbf{Y}(\xi, t) = {}^t(R_2(t) \cos(\xi), R_2(t) \sin(\xi), z_2(t)),$$

and substitute it into (3.1). From the equation for \mathbf{X} we have

$$\begin{aligned} \dot{R}_1 \cos(\xi) &= -\frac{\alpha R_2(z_1 - z_2) \cos(\xi)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{R}_1 \sin(\xi) &= -\frac{\alpha R_2(z_1 - z_2) \sin(\xi)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{z}_1 &= \frac{\beta}{R_1} + \frac{\alpha R_2(R_1 - R_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}. \end{aligned}$$

The dependence of the system on ξ is eliminated by multiplying the first two equations by $\cos(\xi)$ and $\sin(\xi)$, respectively, and adding. The equations for \mathbf{Y} are calculated in the

same way and we arrive at

$$\left\{ \begin{array}{l} \dot{R}_1 = -\frac{\alpha R_2(z_1 - z_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{z}_1 = \frac{\beta}{R_1} + \frac{\alpha R_2(R_1 - R_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{R}_2 = \frac{\alpha\beta R_1(z_1 - z_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{z}_2 = \frac{1}{R_2} - \frac{\alpha\beta R_1(R_1 - R_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ (R_1(0), z_1(0), R_2(0), z_2(0)) = (R_{1,0}, z_{1,0}, R_{2,0}, z_{2,0}). \end{array} \right. \quad (3.2)$$

We note here that a system similar to (3.2) was derived independently by Munakata [40] by directly approximating the induced velocities of vortex rings. First, we observe that z_1 and z_2 can be reduced to one variable, namely $W = z_1 - z_2$. Furthermore, we see by direct calculation that $\beta R_1^2 + R_2^2$ is a conserved quantity. Hence, setting $d^2 = \beta R_{1,0}^2 + R_{2,0}^2$ with $d > 0$, we make the change of variables

$$R_1(t) = \frac{d}{\beta^{1/2}} \cos(\theta(t)), \quad R_2(t) = d \sin(\theta(t))$$

to further reduce the system. We then arrive at

$$\left\{ \begin{array}{l} \dot{\theta} = \frac{\alpha\beta^{1/2}W}{\left(\frac{d^2}{\beta}(\beta^{1/2}\sin\theta - \cos\theta)^2 + W^2\right)^{3/2}} =: F_1(\theta, W), \\ \dot{W} = \frac{\beta^{3/2}\sin\theta - \cos\theta}{d\sin\theta\cos\theta} - \frac{\alpha d^2(\sin\theta + \beta^{1/2}\cos\theta)(\beta^{1/2}\sin\theta - \cos\theta)}{\beta^{1/2}\left(\frac{d^2}{\beta}(\beta^{1/2}\sin\theta - \cos\theta)^2 + W^2\right)^{3/2}} =: F_2(\theta, W), \end{array} \right. \quad (3.3)$$

with initial data (θ_0, W_0) . Here, $W_0 = z_{1,0} - z_{2,0}$ and θ_0 is determined uniquely from the relation

$$R_{1,0} = \frac{d}{\beta^{1/2}} \cos\theta_0, \quad R_{2,0} = d \sin\theta_0.$$

Note that from our problem setting, (θ_0, W_0) is contained in the open set $\Omega_\beta \subset \mathbf{R}^2$ given

by

$$\Omega_\beta = \{(\theta, W) \in \mathbf{R}^2 \mid 0 < \theta < \frac{\pi}{2}, W \in \mathbf{R}, (\theta, W) \neq (\theta_\beta, 0)\},$$

where θ_β is the unique solution of

$$\beta^{1/2} \sin \theta_\beta - \cos \theta_\beta = 0,$$

which is given explicitly by $\theta_\beta = \arctan(1/\beta^{1/2})$. The excluded point in the above definition corresponds to the two filaments overlapping. Since we can reconstruct the solution of (3.2) from the solution $(\theta(t), W(t))$ of (3.3) by

$$\begin{aligned} R_1(t) &= \frac{d}{\beta^{1/2}} \cos(\theta(t)), \quad R_2(t) = d \sin(\theta(t)), \\ z_1(t) &= \int_0^t \frac{\beta}{R_1(\tau)} + \frac{\alpha R_2(\tau)(R_1(\tau) - R_2(\tau))}{((R_1(\tau) - R_2(\tau))^2 + W(\tau)^2)^{3/2}} d\tau, \\ z_2(t) &= \int_0^t \frac{1}{R_2(\tau)} - \frac{\alpha \beta R_2(\tau)(R_1(\tau) - R_2(\tau))}{((R_1(\tau) - R_2(\tau))^2 + W(\tau)^2)^{3/2}} d\tau, \end{aligned}$$

we focus on the solvability and behavior of the solution to system (3.3). It can be checked by direct calculation that the system (3.3) is a Hamiltonian system and the Hamiltonian \mathcal{H} is given by

$$\mathcal{H}(\theta, W) = \frac{1}{2d} \log \left(\frac{(1 - \sin \theta)^{\beta^{3/2}} (1 - \cos \theta)}{(1 + \sin \theta)^{\beta^{3/2}} (1 + \cos \theta)} \right) - \frac{\alpha \beta^{1/2}}{\left(\frac{d^2}{\beta} (\beta^{1/2} \sin \theta - \cos \theta)^2 + W^2 \right)^{1/2}}. \quad (3.4)$$

In other words, $F_1 = \frac{\partial \mathcal{H}}{\partial W}$ and $F_2 = -\frac{\partial \mathcal{H}}{\partial \theta}$. Of course, the Hamiltonian is a conserved quantity of motion. In this formulation, closed orbits revolving around the point $(\theta_\beta, 0)$ correspond to leapfrogging. From here, we treat (3.3) as a two-dimensional dynamical system in Ω_β with parameters d, β , and α , and make use of many tools known for two-dimensional dynamical systems and Hamiltonian systems, for example in Hirsch and Smale [41], to determine the dynamics of the filaments.

We state our main theorems.

Theorem 3.1 *For any $\alpha, d > 0$, $\beta \geq 1$, and $(\theta_0, W_0) \in \Omega_\beta$, there exists a unique time-global solution $(\theta, W) \in C^1(\mathbf{R}) \times C^1(\mathbf{R})$ of (3.3).*

Theorem 3.2 *In addition to the assumptions of Theorem 3.1, if we assume $0 < \alpha < 1/3$,*

then system (3.3) has two equilibrium points $(\theta_*, 0)$ and $(\theta_{**}, 0)$ with $\theta_* \in (0, \theta_\beta)$ and $\theta_{**} \in (\theta_\beta, \pi/2)$, and the following two statements are equivalent.

- (i) The solution with initial data (θ_0, W_0) is a leapfrogging solution. In other words, the solution curve is a closed orbit revolving around the point $(\theta_\beta, 0)$.
- (ii) $\theta_0 \in (\theta_*, \theta_{**})$ and $\mathcal{H}(\theta_0, W_0) < \min\{\mathcal{H}(\theta_*, 0), \mathcal{H}(\theta_{**}, 0)\}$.

Remark 3.3 (Note on the assumption for α in Theorem 3.2) Recall that $\alpha > 0$ was given by $\alpha = 2\delta/\log(\frac{L}{\varepsilon})$, where $\delta, \varepsilon > 0$ were small parameters with $L > 0$ fixed. These parameters were introduced in the course of the derivation of the model system (2.1). Hence, it is natural to assume that α is small and also important that the smallness assumption for α in Theorem 3.2 is independent of the parameters d and β .

The rest of the section is devoted to the proof of the above two theorems.

Proof of Theorem 3.1. Since F_1 and F_2 are smooth in Ω_β , the time-local unique solvability is known. Suppose the maximum existence time $T > 0$ is finite. From the standard theory of dynamical systems, for any compact set $K \subset \Omega_\beta$, there exists $t' \in [0, T)$ such that $(\theta(t'), W(t')) \notin K$. On the other hand, since the Hamiltonian is conserved, there exists $\eta > 0$ and $r > 0$ such that for all $t \in [0, T)$,

$$(\theta(t), W(t)) \in ([\eta, \frac{\pi}{2} - \eta] \times \mathbf{R}) \setminus B_r(\theta_\beta, 0),$$

where $B_r(\theta_\beta, 0)$ is the open ball in \mathbf{R}^2 with center $(\theta_\beta, 0)$ and radius r . This follows from the fact that the Hamiltonian diverges to $-\infty$ at $\theta = 0, \pi/2$ uniformly with respect to W and at the point $(\theta_\beta, 0)$. In particular, since the solution curve is uniformly separated from the point $(\theta_\beta, 0)$, there exists $c_0 > 0$ such that

$$\frac{d^2}{\beta}(\beta^{1/2} \sin \theta(t) - \cos \theta(t))^2 + W(t)^2 \geq c_0$$

for all $t \in [0, T)$. Hence from the second equation in (3.3), we have

$$|\dot{W}| \leq \frac{\beta^{3/2} + 1}{d \sin \eta \cos(\pi/2 - \eta)} + \frac{\alpha d^2 (\beta^{1/2} + 1)^2}{\beta^{1/2} c_0^{3/2}} =: M,$$

which yields

$$|W(t)| \leq |W(0)| + Mt \leq |W_0| + MT$$

for all $t \in [0, T)$. Finally, this shows that for all $t \in [0, T)$, $(\theta(t), W(t))$ is contained in the compact set K' given by

$$K' = \left([\eta, \frac{\pi}{2} - \eta] \times [-|W_0| - MT, |W_0| + MT]\right) \setminus B_r(\theta_\beta, 0),$$

which is a contradiction. The same argument holds for $t < 0$ and hence, the solution exists globally in time and is defined for all $t \in \mathbf{R}$. \square

Proof of Theorem 3.2. We divide the proof of Theorem 3.2 into subsections. First we prove that system (3.3) has exactly two equilibriums as stated in Theorem 3.2.

3.1 Equilibriums of System (3.3)

From the form of F_1 , we see that an equilibrium can only exist on the line segment $(0, \pi/2) \times \{0\}$, and thus, we set $f(\theta) := F_2(\theta, 0)$ and investigate the zeroes of f . First we consider the zeroes in the interval $(0, \theta_\beta)$. Keeping in mind that $\beta^{1/2} \sin \theta - \cos \theta < 0$ in $(0, \theta_\beta)$, by a change of variable $\theta = \arctan x$ we have

$$f(\arctan x) = \frac{(1 + x^2)^{1/2} g_\alpha(x)}{dx(\beta^{1/2}x - 1)^2},$$

where g_α is given by

$$g_\alpha(x) = \beta^{5/3}x^3 - \beta(2\beta + 1)x^2 + \beta^{1/2}(\beta + 2)x - 1 + \alpha\beta(x^2 + \beta^{1/2}x)$$

for $x \in (0, 1/\beta^{1/2})$. We further make the change of variable $y = \beta^{1/2}x$ for simplification and investigate the zeroes of the function h_α given by

$$h_\alpha(y) = \beta y^3 - (2\beta + 1)y^2 + (\beta + 2)y - 1 + \alpha(y^2 + \beta y)$$

in the interval $I_1 = (0, 1)$. We treat h_α as a perturbation of h_0 given by

$$h_0(y) = \beta y^3 - (2\beta + 1)y^2 + (\beta + 2)y - 1,$$

which is h_α with $\alpha = 0$ and prove that h_α has exactly one zero in I_1 . We see from direct calculation that h_0 has one local maximum and one local minimum at

$$y_1 = \frac{\beta + 2}{3\beta}, \quad y_2 = 1,$$

respectively, and

$$h_0(y_1) = \frac{4}{27\beta^2}(\beta - 1)^3 > 0, \quad h_0(y_2) = 0.$$

Since the zero at y_2 is singular, we cannot directly apply the method of perturbation to h_α . Instead, we analyze the positions of the local extrema for $0 < \alpha < 1/3$ to determine the number of zeroes of h_α . First, we observe that the discriminant Δ of the quadratic equation $h'_\alpha(y) = 0$ is given by

$$\Delta = 4[(1 - 3\alpha)\beta^2 - 2(1 + 2\alpha)\beta + (\alpha - 1)] =: 4\phi(\beta).$$

$\phi(\beta) = 0$ has two roots β_\pm given by

$$\beta_- = \frac{1 + 2\alpha - \sqrt{3\alpha(3 - \alpha - \alpha^2)}}{1 - 3\alpha}, \quad \beta_+ = \frac{1 + 2\alpha + \sqrt{3\alpha(3 - \alpha - \alpha^2)}}{1 - 3\alpha}$$

and under the assumption $0 < \alpha < 1/3$, we see that

$$\phi(\beta) < 0 \text{ for } 1 \leq \beta < \beta_+, \quad \phi(\beta) \geq 0 \text{ for } \beta_+ \leq \beta,$$

where we also used the fact that $\phi(1) = -\alpha(9 - \alpha) < 0$. This shows that when $1 \leq \beta < \beta_+$, $\Delta < 0$ which implies $h'_\alpha > 0$ for $y \in (0, 1)$. Since, $h_\alpha(0) = -1$ and $h_\alpha(1) = \alpha(1 + \beta) > 0$, there is exactly one zero in I_1 .

When $\beta_+ \leq \beta$, the roots y_\pm of $h'_\alpha(y) = 0$ are given by

$$y_\pm = \frac{2\beta + 1 - \alpha \pm \sqrt{\phi(\beta)}}{3\beta},$$

where y_- is the local maximum and y_+ is the local minimum. Since h_α is a third order polynomial, it is sufficient to prove that $h_\alpha(y_+) > 0$ to prove that h_α has exactly one root. We have

$$\begin{aligned} y_+ &\geq \frac{1}{3\beta}(2\beta + 1 - \alpha) \geq \frac{1}{3\beta}(\beta + 2 + (\beta_+ - 1) - \alpha) \\ &= \frac{1}{3\beta}\left\{\beta + 2 + \frac{\alpha^{1/2}}{1 - 3\alpha}[(3(3 - \alpha - \alpha^2))^{1/2} + 5\alpha^{1/2} - (1 - 3\alpha)\alpha^{1/2}]\right\} \\ &\geq \frac{\beta + 2}{3\beta}, \end{aligned}$$

which implies $h_0(y_+) \geq 0$. Finally, we have

$$h_\alpha(y_+) = h_0(y_+) + \alpha(y_+^2 + \beta y_+) > 0$$

which shows that h_α also has exactly one root when $\beta_+ \leq \beta$. Hence we have proven that for any $\beta \geq 1$ and $0 < \alpha < 1/3$, h_α has exactly one zero y_* in I_1 and $h'_\alpha(y_*) > 0$. Hence, $\theta_* = \arctan(y_*/\beta^{1/2})$ is the desired zero of $f(\theta)$ in the interval $(0, \theta_\beta)$ and we see

that $f'(\theta_*) > 0$. By a similar argument, we see that there exists a unique $\theta_{**} \in (\theta_\beta, \pi/2)$ such that $f(\theta_{**}) = 0$ and $f'(\theta_{**}) > 0$. We note here that because θ_* and θ_{**} are the only zeroes in the interval $(0, \theta_\beta)$ and $(\theta_\beta, \pi/2)$ respectively, and $f'(\theta_*), f'(\theta_{**}) > 0$, we have the following property for $f(\theta)$.

$$\begin{aligned} f(\theta) &< 0, \quad \text{for } \theta \in (0, \theta_*) \cup (\theta_{**}, \pi/2), \\ f(\theta) &> 0, \quad \text{for } \theta \in (\theta_*, \theta_\beta) \cup (\theta_\beta, \theta_{**}). \end{aligned} \tag{3.5}$$

3.2 Analysis for Solutions with Initial Data of the Form $(\theta_0, 0)$

Since a leapfrogging solution corresponds to a closed orbit revolving around the point $(\theta_\beta, 0)$ in Ω_β , a leapfrogging solution always crosses the lines $(0, \theta_\beta) \times \{0\}$ and $(\theta_\beta, \pi/2) \times \{0\}$ in Ω_β . To this end, we first characterize the solutions with initial data of the form $(\theta_0, 0)$, and prove that the condition given in Theorem 3.2 is necessary and sufficient for leapfrogging to occur.

First we prove that (ii) implies (i). Set $H_* := \min\{\mathcal{H}(\theta_*, 0), \mathcal{H}(\theta_{**}, 0)\}$. Let $\theta_0 \in (\theta_*, \theta_{**})$ satisfy $\mathcal{H}(\theta_0, 0) < H_*$. To make the situation more concrete, we further assume that $\mathcal{H}(\theta_*, 0) > \mathcal{H}(\theta_{**}, 0)$ and make a remark on the case $\mathcal{H}(\theta_*, 0) \leq \mathcal{H}(\theta_{**}, 0)$ at the end. From (3.5) and the fact that $\frac{\partial \mathcal{H}}{\partial \theta}(\theta, 0) = -f(\theta)$, we have

$$\begin{aligned} \frac{\partial \mathcal{H}}{\partial \theta}(\theta, 0) &> 0, \quad \text{for } \theta \in (0, \theta_*) \cup (\theta_{**}, \pi/2), \\ \frac{\partial \mathcal{H}}{\partial \theta}(\theta, 0) &< 0, \quad \text{for } \theta \in (\theta_*, \theta_\beta) \cup (\theta_\beta, \theta_{**}). \end{aligned} \tag{3.6}$$

Moreover, since $\mathcal{H}(\theta_*, 0) > \mathcal{H}(\theta_{**}, 0)$, and $\mathcal{H}(\theta, 0) \rightarrow -\infty$ monotonically as $\theta \rightarrow \theta_\beta -$, there exists a unique $\tilde{\theta} \in (\theta_*, \theta_\beta)$ such that $\mathcal{H}(\tilde{\theta}, 0) = H_*$. This implies that $\theta_0 \in (\tilde{\theta}, \theta_{**}) \setminus \{\theta_\beta\}$.

We assume that $\theta_0 \in (\tilde{\theta}, \theta_\beta)$ since the arguments for the case $\theta_0 \in (\theta_\beta, \theta_{**})$ is the same. We prove that the unique time-global solution $(\theta(t), W(t))$ starting from $(\theta_0, 0)$ obtained in Theorem 3.1, which is defined for $t \in \mathbf{R}$, is a closed orbit revolving around $(\theta_\beta, 0)$. First, we show that the solution is bounded. We observe that as a function of W , the Hamiltonian achieves a minimum at $W = 0$ for each fixed θ . Hence for all $W \in \mathbf{R}$, we have

$$\begin{aligned} \mathcal{H}(\tilde{\theta}, W) &\geq \mathcal{H}(\tilde{\theta}, 0) = H_* > \mathcal{H}(\theta_0, 0), \\ \mathcal{H}(\theta_{**}, W) &\geq \mathcal{H}(\theta_{**}, 0) = H_* > \mathcal{H}(\theta_0, 0). \end{aligned}$$

The above and from the conservation and continuity of the Hamiltonian, there exists

$\eta > 0$ and $r > 0$ such that

$$(\theta(t), W(t)) \in ([\tilde{\theta} + \eta, \theta_{**} - \eta] \times \mathbf{R}) \setminus B_r(\theta_\beta, 0),$$

for all $t \in \mathbf{R}$. Furthermore, if we set

$$\phi(\theta) := \frac{1}{2d} \log \left(\frac{(1 - \sin \theta)^{\beta^{3/2}} (1 - \cos \theta)}{(1 + \sin \theta)^{\beta^{3/2}} (1 + \cos \theta)} \right),$$

we see that as a function of θ , $\mathcal{H}(\theta, W)$ converges to ϕ uniformly as $W \rightarrow \infty$. Since we have

$$\phi'(\theta) = -\frac{(\beta^{3/2} \sin \theta - \cos \theta)}{d \cos \theta \sin \theta},$$

we see that ϕ achieves a maximum at $\theta = \arctan(1/\beta^{3/2}) =: \theta_c$ with $0 < \theta_c < \theta_\beta$ and ϕ is monotone in the intervals $(0, \theta_c)$ and $(\theta_c, \pi/2)$. If $0 < \theta_c \leq \tilde{\theta}$, for $\varepsilon_1 > 0$ given by

$$\varepsilon_1 = \frac{\alpha \beta^{1/2}}{2 \left\{ \frac{d^2}{\beta} (\beta^{1/2} \sin \theta_{**} - \cos \theta_{**})^2 \right\}^{1/2}},$$

there exists $W_1 > 0$ such that for all $\theta \in (\tilde{\theta}, \theta_{**})$, and $W > W_1$ we have

$$\mathcal{H}(\theta, W) > \phi(\theta) - \varepsilon_1 > \phi(\theta_{**}) - 2\varepsilon_1 = \mathcal{H}(\theta_{**}, 0) = H_* > \mathcal{H}(\theta_0, 0).$$

If $\tilde{\theta} < \theta_c < \theta_\beta$, choose $\theta' \in \{\tilde{\theta}, \theta_{**}\}$ so that $\phi(\theta') = \min\{\phi(\tilde{\theta}), \phi(\theta_{**})\}$. Then for $\varepsilon_2 > 0$ given by

$$\varepsilon_2 = \frac{\alpha \beta^{1/2}}{2 \left\{ \frac{d^2}{\beta} (\beta^{1/2} \sin \theta' - \cos \theta')^2 \right\}^{1/2}},$$

there exists $W_2 > 0$ such that for all $\theta \in (\tilde{\theta}, \theta_{**})$ and $W > W_2$, we have

$$\mathcal{H}(\theta, W) > \phi(\theta) - \varepsilon_2 > \phi(\theta') - 2\varepsilon_2 = \mathcal{H}(\theta', 0) = H_* > \mathcal{H}(\theta_0, 0).$$

In either case, we see that the value of the Hamiltonian on the segment $[\tilde{\theta}, \theta_{**}] \times \{W_*\}$, where $W_* = \max\{W_1, W_2\}$, is strictly greater than $\mathcal{H}(\theta_0, 0)$ and hence the solution curve cannot cross this segment. Since the Hamiltonian is symmetric with respect to $W = 0$, we finally see that

$$(\theta(t), W(t)) \in ([\tilde{\theta} + \eta, \theta_{**} - \eta] \times [-W_*, W_*]) \setminus B_r(\theta_\beta, 0) =: K_*,$$

for all $t \in \mathbf{R}$, and in particular, the solution is bounded.

Next we set

$$L_0 := \{(\theta, W) \in \Omega_\beta \mid \mathcal{H}(\theta, W) = \mathcal{H}(\theta_0, 0)\} \cap K_*.$$

As a closed subset of the compact set K_* , L_0 is a compact subset of Ω_β . From the conservation of the Hamiltonian and the way we chose η , r , and W_* , we see that L_0 is also an invariant set and hence we have

$$L_\omega(\theta_0, 0) \subset L_0,$$

where $L_\omega(\theta_0, 0)$ is the ω -limit set of $(\theta_0, 0)$. Since $(\theta(t), W(t))$ is bounded for $t > 0$, it converges along some series $\{t_n\}_{n=1}^\infty$ with $t_n \rightarrow \infty$ as $n \rightarrow \infty$, and in particular, $L_\omega(\theta_0, 0)$ is not empty. Since $L_\omega(\theta_0, 0)$ is a non-empty compact set and contains no equilibriums (recall that the equilibriums $(\theta_*, 0)$ and $(\theta_{**}, 0)$ are outside the set L_0), it is a closed orbit by the Poincaré–Bendixson Theorem. Moreover, the point $(\theta_\beta, 0)$ is in the interior of this closed orbit, because if it is not, then the closed orbit would enclose an open subset of Ω_β in which an equilibrium must exist, which leads to a contradiction. This proves that $L_\omega(\theta_0, 0)$ is a closed orbit revolving around $(\theta_\beta, 0)$. Since $L_\omega(\theta_0, 0) \subset L_0$, there exists $\theta_1 \in (\tilde{\theta} + \eta, \theta_\beta)$ and $\theta_2 \in (\theta_\beta, \theta_{**} - \eta)$ such that $(\theta_1, 0), (\theta_2, 0) \in L_\omega(\theta_0, 0)$. The values θ_1 and θ_2 satisfying this property are unique in their respective intervals because the Hamiltonian is monotone along the line segments $[\tilde{\theta} + \eta, \theta_\beta] \times \{0\}$ and $[\theta_\beta, \theta_{**} - \eta] \times \{0\}$. This uniqueness implies that $\theta_1 = \theta_0$, which proves that $L_\omega(\theta_0, 0)$ coincides with the orbit starting from $(\theta_0, 0)$.

In summary, we have proven that the orbit starting from $(\theta_0, 0)$ is a closed orbit revolving around $(\theta_\beta, 0)$ corresponding to a leapfrogging solution. We further have the characterization

$$L_\omega(\theta_0, 0) = L_0,$$

which we prove by contradiction. Suppose there exists $(\bar{\theta}, \bar{W}) \in L_0$ such that $(\bar{\theta}, \bar{W}) \notin L_\omega(\theta_0, 0)$. We first see that $\bar{W} \neq 0$, since $(\bar{\theta}, 0) \in L_0$ implies $\bar{\theta} = \theta_1$ or θ_2 , which contradicts $(\bar{\theta}, 0) \notin L_\omega(\theta_0, 0)$. Henceforth, we assume $\bar{W} > 0$ since the proof for the other case is the same. Now, if $\bar{\theta} \in [\tilde{\theta} + \eta, \theta_1]$, we have

$$\mathcal{H}(\bar{\theta}, \bar{W}) > \mathcal{H}(\bar{\theta}, 0) \geq \mathcal{H}(\theta_1, 0) = \mathcal{H}(\theta_0, 0)$$

from the monotonicity of \mathcal{H} along the line $\{\bar{\theta}\} \times \mathbf{R}$ and the monotonicity along the line segment $[\bar{\theta}, \theta_1] \times \{0\}$, and this contradicts $(\bar{\theta}, \bar{W}) \in L_0$. The case $\bar{\theta} \in [\theta_2, \theta_{**} - \eta]$ leads to a contradiction by the same argument. If $\bar{\theta} \in (\theta_1, \theta_2)$ and $(\bar{\theta}, \bar{W})$ is in the interior of the

closed orbit $L_\omega(\theta_0, 0)$, there exists $\tilde{W} > \overline{W}$ such that $(\bar{\theta}, \tilde{W}) \in L_\omega(\theta_0, 0)$. Then we have

$$\mathcal{H}(\bar{\theta}, \overline{W}) < \mathcal{H}(\bar{\theta}, \tilde{W}) = \mathcal{H}(\theta_0, 0),$$

which contradicts $(\bar{\theta}, \overline{W}) \in L_0$. Similarly, if $(\bar{\theta}, \overline{W})$ is outside of the closed orbit, there exists $\tilde{W} < \overline{W}$ such that $(\bar{\theta}, \tilde{W}) \in L_\omega(\theta_0, 0)$. Again, this implies the estimate

$$\mathcal{H}(\bar{\theta}, \overline{W}) > \mathcal{H}(\bar{\theta}, \tilde{W}) = \mathcal{H}(\theta_0, 0),$$

which contradicts $(\bar{\theta}, \overline{W}) \in L_0$. Hence we have $L_\omega(\theta_0, 0) = L_0$. We can express L_0 as

$$L_0 = \{(\theta, W) \in \Omega_\beta \mid \mathcal{H}(\theta, W) = \mathcal{H}(\theta_0, 0)\} \cap M,$$

with $M = [\theta_*, \theta_{**}] \times \mathbf{R}$, because the value of the Hamiltonian on $M \setminus K_*$ is different from $\mathcal{H}(\theta_0, 0)$, and thus, replacing K_* with M does not add any points. This expression will be utilized to derive the necessary and sufficient condition for leapfrogging to occur for solutions with general initial data.

Finally, we make some remarks on the case $\mathcal{H}(\theta_*, 0) \not\asymp \mathcal{H}(\theta_{**}, 0)$. When $\mathcal{H}(\theta_*, 0) = \mathcal{H}(\theta_{**}, 0)$, the same proof holds with $\hat{\theta} = \theta_*$. When $\mathcal{H}(\theta_*, 0) < \mathcal{H}(\theta_{**}, 0)$, there is a unique $\hat{\theta} \in (\theta_\beta, \theta_{**})$ such that $\mathcal{H}(\hat{\theta}, 0) = H_*$. This $\hat{\theta}$ plays the same role as $\tilde{\theta}$, and the same arguments for the case $\mathcal{H}(\theta_*, 0) < \mathcal{H}(\theta_{**}, 0)$ holds.

Next we prove that (i) implies (ii). Suppose that a solution starting from $(\theta_0, 0)$ is a leapfrogging solution. Since $\mathcal{H}(\theta_*, 0)$ and $\mathcal{H}(\theta_{**}, 0)$ are the maximum value of $\mathcal{H}(\theta, 0)$ in their respective intervals $(0, \theta_\beta)$ and $(\theta_\beta, \pi/2)$, in order for a solution curve to cross over the segments $(0, \theta_\beta) \times \{0\}$ and $(\theta_\beta, \pi/2) \times \{0\}$, the value of the Hamiltonian on this solution curve must be less than or equal to the smaller of the two. In other words, $\mathcal{H}(\theta_0, 0) \leq H_*$ holds. If $\mathcal{H}(\theta_0, 0) = H_*$ holds, the only possible points at which the solution curve can cross the segments $(0, \theta_\beta) \times \{0\}$ and $(\theta_\beta, \pi/2) \times \{0\}$ are at the equilibrium points. This would result in the solution converging to one of the equilibrium points, and is not a leapfrogging solution. Hence, for a leapfrogging solution, $\mathcal{H}(\theta_0, 0) < H_*$ holds.

Furthermore, $(\theta_0, 0)$ is not on the lines $\{\theta_*\} \times \mathbf{R}$ or $\{\theta_{**}\} \times \mathbf{R}$ since the value of the Hamiltonian is greater than or equal to H_* along these lines. Consequently, if $\theta_0 \in (0, \theta_*) \cup (\theta_{**}, \pi/2)$, the solution curve cannot cross over from one side of these lines to the other, which means that the solution is not a leapfrogging solution. This implies that $\theta_0 \in (\theta_*, \theta_{**})$, and condition (ii) holds.

We summarize the conclusions of this subsection in the following lemma.

Lemma 3.4 *For initial data of the form $(\theta_0, 0) \in \Omega_\beta$, we have the following.*

- (i) *If $\theta_0 \in (\theta_*, \theta_{**})$ and $\mathcal{H}(\theta_0, 0) < H_*$, then the solution starting from $(\theta_0, 0)$ is a leapfrogging solution. Moreover, the closed orbit $L_\omega(\theta_0, 0)$ can be expressed as*

$$L_\omega(\theta_0, 0) = \{(\theta, W) \in \Omega_\beta \mid \mathcal{H}(\theta, W) = \mathcal{H}(\theta_0, 0)\} \cap M,$$

where $M = [\theta_*, \theta_{**}] \times \mathbf{R}$.

(ii) Otherwise, the solution is not a leapfrogging solution.

3.3 Remarks on Solutions with General Initial Data

Let $(\theta_0, W_0) \in \Omega_\beta$ satisfy $\theta_0 \in (\theta_*, \theta_{**})$ and $\mathcal{H}(\theta_0, W_0) < H_*$. Since $\mathcal{H}(\theta, 0)$ takes all values between $-\infty$ and H_* on the set $(\theta_*, \theta_\beta) \cup (\theta_\beta, \theta_{**})$, there exists $\theta_{LF} \in (\theta_*, \theta_\beta) \cup (\theta_\beta, \theta_{**})$ such that $\mathcal{H}(\theta_{LF}, 0) = \mathcal{H}(\theta_0, W_0)$. Moreover, from Lemma 3.4, the orbit containing $(\theta_{LF}, 0)$ is a closed orbit corresponding to a leapfrogging solution. Since

$$(\theta_0, W_0) \in \{(\theta, W) \in \Omega_\beta \mid \mathcal{H}(\theta, W) = \mathcal{H}(\theta_{LF}, 0)\} \cap M,$$

Lemma 3.4 implies that (θ_0, W_0) is on the closed orbit containing $(\theta_{LF}, 0)$ and hence, the solution starting from (θ_0, W_0) is a leapfrogging solution.

On the other hand, suppose either $\mathcal{H}(\theta_0, W_0) \geq H_*$ or $\theta_0 \notin (\theta_*, \theta_{**})$ holds. We prove that solution curves starting from these initial data are not leapfrogging solutions. If $\mathcal{H}(\theta_0, W_0) \geq H_*$, then the solution starting from (θ_0, W_0) is not a leapfrogging solution since the value of the Hamiltonian of a leapfrogging solution is strictly less than H_* from Lemma 3.4. If $\theta_0 \notin (\theta_*, \theta_{**})$ holds, we only need to consider the case when $\mathcal{H}(\theta_0, W_0) < H_*$ also holds. Since $\mathcal{H}(\theta_0, W_0) < H_*$, $\theta_0 \in (0, \theta_*) \cup (\theta_{**}, \pi/2)$ because the value of the Hamiltonian on the lines $\{\theta_*\} \times \mathbf{R}$ and $\{\theta_{**}\} \times \mathbf{R}$ are greater than or equal to H_* . Furthermore, since the Hamiltonian is conserved, the solution curve starting from (θ_0, W_0) cannot cross over from one side of these lines to the other and hence, the solution is not a leapfrogging solution. This finishes the proof of Theorem 3.2. \square

4 Head-on Collision

So far we have focused on leapfrogging motion, but the model (2.1) has the potential to mathematically describe other patterns of motion that are observed in the real world. One such motion pattern is head-on collision. When two coaxial vortex rings move towards each other, there is a chance that the two rings collide, which is called head-on collision. Here we give a sufficient condition for head-on collision to occur. We consider the system

of ordinary differential equations considered earlier.

$$\left\{ \begin{array}{l} \dot{R}_1 = -\frac{\alpha R_2(z_1 - z_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{z}_1 = \frac{\beta}{R_1} + \frac{\alpha R_2(R_1 - R_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{R}_2 = \frac{\alpha \beta R_1(z_1 - z_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ \dot{z}_2 = \frac{1}{R_2} - \frac{\alpha \beta R_1(R_1 - R_2)}{((R_1 - R_2)^2 + (z_1 - z_2)^2)^{3/2}}, \\ (R_1(0), z_1(0), R_2(0), z_2(0)) = (R_{1,0}, z_{1,0}, R_{2,0}, z_{2,0}), \end{array} \right.$$

and set $\beta = -1$. From direct calculation, we see that $R_1^2 - R_2^2$ is conserved. We further assume $R_{1,0} = R_{2,0}$, which implies $R_1 = R_2$ throughout the motion. Further setting

$$\theta := \log(R_1) \quad \text{and} \quad W = z_1 - z_2,$$

we have

$$\left\{ \begin{array}{l} \dot{\theta} = -\frac{\alpha W}{|W|^3}, \\ \dot{W} = -2e^{-\theta}, \\ (\theta, W)(0) = (\theta_0, W_0), \end{array} \right. \quad (4.1)$$

which has a Hamiltonian $\mathcal{H}(\theta, W)$ of the form

$$\mathcal{H}(\theta, W) = -2e^{-\theta} + \frac{\alpha}{|W|}.$$

We prove the following.

Theorem 4.1 *For any $\alpha > 0$ and $(\theta_0, W_0) \in \mathbf{R} \times (\mathbf{R} \setminus \{0\})$, the following holds.*

- (i) *When $W_0 > 0$ and $\mathcal{H}(\theta_0, W_0) > 0$, there exists a $T_* > 0$ such that there exists a unique finite-time solution to (4.1) satisfying $(\theta, W) \in C^1([0, T_*)) \times C^1([0, T_*))$, which corresponds to rings colliding at $t = T_*$.*
- (ii) *When $W_0 < 0$, there exists a unique time-global solution to (4.1) satisfying $(\theta, W) \in C^1([0, \infty)) \times C^1([0, \infty))$.*

Proof of Theorem 4.1. Most of the claims in the theorem follow from standard theory of ordinary differential equations. We focus on the case (i) and prove that the solution exists only for a finite time and is in fact a solution corresponding to the two rings colliding.

First we set $H_0 := \mathcal{H}(\theta_0, W_0)$ and observe that

$$\dot{\theta} = -\frac{1}{\alpha}(H_0 + 2e^{-\theta})^2 \leq -\frac{4}{\alpha}e^{-2\theta}.$$

Hence, if we set ψ as the solution of

$$\begin{cases} \dot{\psi} = -\frac{4}{\alpha}e^{-2\psi} \\ \psi(0) = \theta_0 \end{cases}$$

we have $\theta(t) \leq \psi(t)$. Direct calculation shows that

$$\psi(t) = \frac{1}{2} \log \left(e^{\theta_0} - \frac{8}{\alpha} t \right).$$

This implies that $\theta(t)$ is a finite-time solution defined on $[0, T_*)$ for some T_* satisfying

$$T_* \leq \frac{\alpha}{8} e^{\theta_0}$$

and that $\theta(t)$ diverges to $-\infty$ as $t \rightarrow T_*-$. Hence the solution (θ, W) is a finite-time solution.

From the equation for W , we have $\dot{W}(t) \leq 0$ for all $t \in [0, T_*)$. Also from the conservation of the Hamiltonian, if $H_0 > 0$ and $W_0 > 0$, we have $W(t) > 0$ for all $t \in [0, T_*)$. This implies that there exists some $W_* \geq 0$ such that $W(t) \rightarrow W_*$ as $t \rightarrow T_*-$. Finally, if $W_* > 0$, $W(t)$ is bounded from below by some positive constant near $t = T_*$. This would lead to an a priori estimate for θ , which contradicts the fact that θ only exists for finite time. Hence, $W_* = 0$, which means that the two vortex rings collide at $t = T_*$. \square .

References

- [1] H. HELMHOLTZ, Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen, *J. Reine Angew. Math.* **55**:25–55 (1858).
- [2] F. DYSON, The Potential of an Anchor Ring, *Philos. Trans. Roy. Soc. London Ser. A* **184**:43–95 (1893).
- [3] F. DYSON, The Potential of an Anchor Ring - Part II, *Philos. Trans. Roy. Soc. London Ser. A* **184**:1107–1169 (1893).

- [4] W. M. HICKS, On the mutual threading of vortex rings, *Proc. R. Soc. Long. A* **102**:111–131 (1922).
- [5] H. LAMB, Hydrodynamics, 6th Edition, *Cambridge University Press*, 1932.
- [6] A.A. GURZHII, M.Y. KONSTANTINOV, and V.V. MELESHKO, Interaction of Coaxial Vortex Rings in an Ideal Fluid, *Izv. Akad. Nauk SSSR Mekh. Zhidk. Gaza* **2**:78–84 (1988). [*Fluid Dyn.* **23**:224–229 (1988) (English translation)].
- [7] V.I. BOYARINTSEV, E.S. LEVCHENKO, and A.S. SAVIN, Motion of Two Vortex Rings, *Izv. Akad. Nauk SSSR Mekh. Zhidk. Gaza* **5**:176–177 (1985). [*Fluid Dyn.* **20**:818–819 (1985) (English translation)].
- [8] K. SHARIFF, A. LEONARD, and J.H. FERZIGER, Dynamical systems analysis of fluid transport in time-periodic vortex ring flows, *Phys. Fluid* **18**:11 pages (page number unavailable) (2006).
- [9] N. RILEY and D.P. STEVENS, A note on leapfrogging vortex rings, *Fluid Dynam. Res.* **11**:235–244 (1993).
- [10] Y. NAKANISHI, K. KAMEMOTO, and M. NISHIO, Modification of Vortex Model for Consideration of Viscous Effect: Examination of Vortex Model on Interaction of Vortex Rings, *JSME International Journal Series B* **37**:815–820 (1994).
- [11] T.T. LIM, A note on the leapfrogging between two coaxial vortex rings at low Reynolds numbers, *PHYS. FLUID* **9**:239–241 (1997).
- [12] M. CHENG, J. LOU, and T.T. LIM, Leapfrogging of multiple coaxial viscous vortex rings, *Phys. Fluid* **27**:8 pages (page number unavailable) (2015).
- [13] T. MAXWORTHY, Some experimental studies of vortex rings, *J. Fluid Mech.* **81**:465–495 (1977).
- [14] Y. OSHIMA, The Game of Passing-Through of a Pair of Vortex Rings, *J. Phys. Soc. Jpn.* **45**:660–664 (1978).
- [15] Y. OSHIMA, T. KAMBE, and S. ASAKA, Interaction of Two Vortex Rings Moving along a Common Axis of Symmetry, *J. Phys. Soc. Jpn.* **38**:1159–1166 (1975).
- [16] H. YAMADA and T. MATSUI, Preliminary study of mutual slip-through of a pair of vortices, *Phys. Fluids* **21**:292–294 (1978).
- [17] P.G. SAFFMAN, The Velocity of Viscous Vortex Rings, *Studies in Appl. Math.* **54**:371–380 (1975).
- [18] P.G. SAFFMAN, Vortex Dynamics, *Cambridge University Press*, 1993.

- [19] Y. FUKUMOTO, Higher-order asymptotic theory for the velocity field induced by an inviscid vortex ring, *Fluid Dynam. Res.* **30**:65–92 (2002).
- [20] A.V. BORISOV, A.A. KILIN, and I.S. MAMAEV, The Dynamics of Vortex Rings: Leapfrogging, Choreographies and the Stability Problem, *Regul. Chaotic Dyn.* **18**:33–62 (2013).
- [21] E.A. NOVIKOV, Generalized dynamics of three-dimensional vortical singularities(vortons), *Zh. Eksp. Teor. Fiz.* **84**:975–981 (1983). [*J. Exp. Theor. Phys.* **57**:566–569 (1983) (English translation)].
- [22] V.V. MELESHKO and G.J.F. VAN HEIJST, Two approaches to the analysis of the coaxial interaction of vortex rings, *Int. J. Fluid Mech. Res.* **30**:166–183 (2003).
- [23] K. SHARIFF, A. LEONARD, Vortex Rings, *Annu. Rev. Fluid Mech.* **24**:235–279 (1992).
- [24] V.V. MELESHKO, Coaxial axisymmetric vortex rings: 150 years after Helmholtz, *Theor. Comput. Fluid Dyn.* **24**:403–431 (2010).
- [25] D.G. CRIGHTON, Jet noise and the effects of jet forcing, *In: The Role of Coherent Structures in Modelling Turbulence and Mixing. Lecture Notes in Physics* **136**, Springer, Berlin, Heidelberg, 1981.
- [26] A.K.M.F. HUSSAIN and K.B.M.Q. ZAMAN, Vortex pairing in a circular jet under controlled excitation. Part 2. Coherent structure dynamics, *J. Fluid Mech.* **101**:493–544 (1980).
- [27] K.B.M.Q. ZAMAN, Far-field noise of a subsonic jet under controlled excitation, *J. Fluid Mech.* **152**:83–111 (1985).
- [28] L.S. DA RIOS, Sul moto d’un liquido indefinito con un filetto vorticoso di forma qualunque (in Italian), *Rend. Circ. Mat. Palermo* **22**:117–135 (1906).
- [29] Y. MURAKAMI, H. TAKAHASHI, Y. UKITA, and S. FUJIWARA, On the vibration of a vortex filament (in Japanese), *Applied Physics Colloquium*, **6**:1–5 (1937).
- [30] R.J. ARMS and F.R. HAMA, Localized-induction concept on a curved vortex and motion of an elliptic vortex ring, *Phys. Fluids* **8**:553–559 (1965).
- [31] T. MAXWORTHY, The structure and stability of vortex rings, *J. Fluid Mech.* **51**:15–32 (1972).
- [32] S.E. WIDNALL and C.Y. TSAI, The instability of the thin vortex ring of constant vorticity, *Phil. Trans. R. Soc. Lond. A* **287**:273–305 (1977).

- [33] S.E. WIDNALL and J.P. SULLIVAN, On the stability of vortex rings, *Phil. Trans. R. Soc. Lond. A* **332**:335–353 (1973).
- [34] Y. FUKUMOTO and Y. HATTORI, Curvature instability of a vortex ring, *J. Fluid Mech.* **526**:77–115 (2005).
- [35] R. KLEIN, A.J. MAJDA, and K. DAMODARAN, Simplified equations for the interaction of nearly parallel vortex filaments, *J. Fluid Mech.* **288**:201–248 (1995).
- [36] R.L. JERRARD and C. SEIS, On the Vortex Filament Conjecture for Euler Flows, *Arch. Ration. Mech. Anal.* **224**:135–172 (2017).
- [37] Y. GIGA and T. MIYAKAWA, Navier–stokes flow in R^3 with measures as initial vorticity and morrey spaces, *Comm. Partial Differential Equations* **14**:577–618 (1989).
- [38] H. FENG and V. ŠVERÁK, On the Cauchy Problem for Axi-Symmetric Vortex Rings, *Arch. Ration. Mech. Anal.* **215**:89–123 (2015).
- [39] P. K. NEWTON, The N-Vortex Problem, *Applied Mathematical Sciences* **145** Springer-Verlag, New York 2001.
- [40] T. MUNAKATA, The Mathematics of Vortex Motion (in Japanese), *Master Thesis* (2016), Unpublished.
- [41] M. W. HIRSCH and S. SMALE, Differential Equations, Dynamical Systems, and Linear Algebra, *Academic Press*, 1974.

Department of Mathematics,
Faculty of Science and Technology, Tokyo University of Science
2641 Yamazaki, Noda, Chiba 278-8510, Japan
E-mail: aiki_masashi@ma.noda.tus.ac.jp