

LARGE DEVIATION PRINCIPLE FOR ARITHMETIC MEAN OF CONTINUED FRACTION DIGITS

Hiroki Takahasi
Keio Institute of Pure and Applied Sciences
Keio University

¹ Each irrational number $x \in (0, 1)$ has the continued fraction expansion

$$x = \frac{1}{a_1 + \frac{1}{a_2 + \cdots}},$$

where each a_i is positive integers. Khinchin [4] noted that for λ -almost every $x \in (0, 1)$ the inequality $a_n(x) > n \log n$ holds for infinitely many $n \in \mathbb{N}$ as a consequence of Borel-Bernstein's theorem [1, 2, 3], and thus

$$\limsup_{n \rightarrow \infty} \frac{a_n}{n} = \infty \quad \lambda\text{-a.e.},$$

and S_n/n does not converge where $S_n = a_1 + a_2 + \cdots + a_n$. In fact,

$$(0.1) \quad \lim_{n \rightarrow \infty} \frac{S_n}{n} = \infty \quad \lambda\text{-a.e.}$$

Hence, none of the classical limit theorems in probability, such as the weak and strong laws of large numbers and central limit theorems hold for the sum. In this paper we establish the level-1 Large Deviation Principle (LDP), i.e., show the existence of a rate function which estimates exponential probabilities with which the time average S_n/n stays away from ∞ .

Main Theorem. *Let $\psi: \mathbb{N} \setminus \{0\} \rightarrow \mathbb{R}$ satisfy $\lim_{n \rightarrow \infty} \psi(n) = \infty$ and $\int \psi(a_1(x)) dx = \infty$. There exists $\alpha^- \in \mathbb{R}$ such that the following holds:*

- for every $\alpha \in \mathbb{R}$ the limit

$$J(\alpha) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \lambda \left\{ x \in (0, 1) \setminus \mathbb{Q} : \frac{1}{n} \sum_{i=1}^n \psi(a_i(x)) \leq \alpha \right\}$$

exists and is finite if and only if $\alpha \geq \alpha^-$. The function $\alpha \in [\alpha^-, \infty) \mapsto J(\alpha)$ is lower semi-continuous, strictly positive, convex, strictly monotone decreasing and $J(\alpha) \rightarrow 0$ as $\alpha \rightarrow \infty$;

¹Part of the contents of this paper has been taken from the introduction of [5].

- for every $\alpha \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \lambda \left\{ x \in (0, 1) \setminus \mathbb{Q} : \frac{1}{n} \sum_{i=1}^n \psi(a_i(x)) \geq \alpha \right\} = 0.$$

To obtain the result for the arithmetic mean of digits, take $\psi(n) = n$. For a proof of the main theorem, see [5]. The arithmetic mean of digits of the backward continued fraction expansion has a totally different behavior, see [6].

REFERENCES

- [1] Bernstein, F.: Über eine Anwendung der Mengenlehre auf ein aus der Theorie der säkularen Störungen herrührendes Problem, *Math. Ann.* **71**, 417–439 (1912)
- [2] Borel, E.: Les probabilités dénombrables et leurs applications arithmétiques, *Rend. Circ. Mat. Palermo* **27**, 247–271 (1909)
- [3] Borel, E.: Sur un problème de probabilités relatif aux fractions continues. *Math. Ann.* **72**, 578–584 (1912)
- [4] Khinchin, A.Y.: *Continued Fractions*. University of Chicago Press, Chicago, London, 1964.
- [5] Takahasi, H.: Large deviation principle for arithmetic functions in continued fraction expansion, *Monatshefte für Mathematik*, **190** 137–152 (2019)
- [6] Takahasi, H.: Large deviation principle for harmonic/geometric/arithmetic mean of digits in backward continued fraction expansion. arXiv:1912.11815

KEIO INSTITUTE OF PURE AND APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS,
KEIO UNIVERSITY, YOKOHAMA, 223-8522, JAPAN

Email address: hiroki@math.keio.ac.jp

URL: <http://www.math.keio.ac.jp/~hiroki/>