

## A LOGIC OF INJECTIVITY

J. ADÁMEK, M. HÉBERT AND L. SOUSA

(communicated by George Janelidze)

### *Abstract*

Injectivity of objects with respect to a set  $\mathcal{H}$  of morphisms is an important concept of algebra, model theory and homotopy theory. Here we study the logic of injectivity consequences of  $\mathcal{H}$ , by which we understand morphisms  $h$  such that injectivity with respect to  $\mathcal{H}$  implies injectivity with respect to  $h$ . We formulate three simple deduction rules for the injectivity logic and for its finitary version where morphisms between finitely ranked objects are considered only, and prove that they are sound in all categories, and complete in all “reasonable” categories.

## 1. Introduction

Recall that an object  $A$  is injective w.r.t. a morphism  $h : P \rightarrow P'$  provided that every morphism from  $P$  to  $A$  factors through  $h$ . We address the following problem: given a set  $\mathcal{H}$  of morphisms, which morphisms  $h$  are *injectivity consequences* of  $\mathcal{H}$  in the sense that every object injective w.r.t. all members of  $\mathcal{H}$  is also injective w.r.t.  $h$ ? We denote the injectivity consequence relationship by  $\mathcal{H} \models h$ .

This is a classical topic in general algebra: the *equational logic* of Garrett Birkhoff [10] is a special case. In fact, an equation  $s = t$  is a pair of elements of a free algebra  $F$ , and that pair generates a congruence  $\sim$  on  $F$ . An algebra  $A$  satisfies  $s = t$  iff it is injective w.r.t. the canonical epimorphism

$$h : F \rightarrow F / \sim .$$

Thus, if we restrict our sets  $\mathcal{H}$  to regular epimorphisms with free domains, then the logic of injectivity becomes precisely the equational logic. However, there are other important cases in algebra: recall for example the concept of injective module, where  $\mathcal{H}$  is the set of all monomorphisms (in the category of modules).

To mention an example from homotopy theory, recall that a *Kan complex* [14] is a simplicial set injective w.r.t. all the monomorphisms  $\Delta_n^k \hookrightarrow \Delta_n$  (for  $n, k \in \mathbb{N}$ ,  $k \leq n$ ) where  $\Delta_n$  is the complex generated by a single  $n$ -simplex and  $\Delta_n^k$  is the subcomplex obtained by deleting the  $k$ -th 1-simplex and all adjacent faces. We can

---

The third author acknowledges financial support by the Center of Mathematics of the University of Coimbra and the School of Technology of Viseu

Received July 27, 2006, revised May 16, 2007; published on December 27, 2007.

2000 Mathematics Subject Classification: 18A15, 18C10, 18C35, 03B99, 03C75, 55U35.

Key words and phrases: Injectivity, logic of injectivity, deduction rules, Small Object Argument, locally presentable categories, strongly locally ranked categories

© 2007, J. Adámek, M. Hébert and L. Sousa. Permission to copy for private use granted.

ask for example whether Kan complexes can be specified by a simpler collection of monomorphisms, as a special case of our injectivity logic.

Injectivity establishes a Galois correspondence between objects and morphisms of a category. The closed families on the side of objects are called *injectivity classes*: for every set  $\mathcal{H}$  of morphisms we obtain the injectivity class  $\text{Inj}\mathcal{H}$ , i.e., the class of all objects injective w.r.t.  $\mathcal{H}$ . In [5] small-injectivity classes in locally presentable categories were characterized as precisely the full accessible subcategories closed under products, and in [18] this was sharpened in the following sense. Let us call a morphism  $\lambda$ -ary if its domain and codomain are  $\lambda$ -presentable objects. Injectivity classes with respect to  $\lambda$ -ary morphisms are precisely the full subcategories closed under products,  $\lambda$ -filtered colimits, and  $\lambda$ -pure subobjects. For injectivity w.r.t. cones or trees of morphisms similar results are in [7] and [15].

In the present paper we study closed sets on the side of morphisms, i.e., we develop a deduction system for the above injectivity consequence relationship  $\models$ . It has altogether three deduction rules, which are quite intuitive. Firstly, observe that every object injective w.r.t. a composite  $h = h_2 \cdot h_1$  is injective w.r.t. the first morphism  $h_1$ . This gives us the first deduction rule

$$\text{CANCELLATION} \quad \frac{h_2 \cdot h_1}{h_1}$$

It is also easy to see that injectivity w.r.t.  $h$  implies injectivity w.r.t. any morphism  $h'$  opposite to  $h$  in a pushout (along an arbitrary morphism), which yields the rule

$$\text{PUSHOUT} \quad \frac{h}{h'} \quad \text{for every pushout} \quad \begin{array}{ccc} & \xrightarrow{h} & \\ \downarrow & & \downarrow \\ & \xrightarrow{h'} & \end{array}$$

Finally, an object injective w.r.t. two composable morphisms is also injective w.r.t. their composite. The same holds for three, four, ... morphisms – but also for a *transfinite composite* as used in homotopy theory. For example, given an  $\omega$ -chain of morphisms

$$A_0 \xrightarrow{h_0} A_1 \xrightarrow{h_1} A_2 \xrightarrow{h_2} \dots$$

then their  $\omega$ -composite is the first morphism  $c_0 : A_0 \rightarrow C$  of (any) colimit cocone  $c_n : A_n \rightarrow C$  ( $n \in \mathbb{N}$ ) of the chain. Observe that  $c_0$  is indeed an injectivity consequence of  $\{h_i; i < \omega\}$ . For every ordinal  $\lambda$  we have the concept of a  $\lambda$ -composite of morphisms (see 2.10 below) and the following deduction rule, expressing the fact that an object injective w.r.t. each  $h_i$  is injective w.r.t. the transfinite composite:

$$\text{TRANSFINITE COMPOSITION} \quad \frac{h_i (i < \lambda)}{h} \quad \text{for every } \lambda\text{-composite } h \text{ of } (h_i)_{i < \lambda}$$

We are going to prove that the Injectivity Logic based on the above three rules is sound and complete. That is, given a set  $\mathcal{H}$  of morphisms, then  $\mathcal{H} \models h$  holds for precisely those morphisms  $h$  which can be proved from assumptions in  $\mathcal{H}$  using the three deduction rules above. This holds in a number of categories, e.g., in

- (a) every variety of algebras,
- (b) the category of topological spaces and many nice subcategories (e.g. Hausdorff spaces), and
- (c) every locally presentable category of Gabriel and Ulmer.

We introduce the concept of a strongly locally ranked category encompassing (a)-(c) above, and prove the soundness and completeness of our Injectivity Logic in all such categories.

Observe that the above logic is infinitary, in fact, it has a proper class of deduction rules: one for every ordinal  $\lambda$  in the instance of TRANSFINITE COMPOSITION. We also study, following the footsteps of Grigore Roşu, the completeness of the corresponding Finitary Injectivity Logic: it is the restriction of the above logic to  $\lambda$  finite. Well, all we need to consider are the cases  $\lambda = 2$ , called COMPOSITION, and  $\lambda = 0$ , called IDENTITY:

$$\begin{array}{l} \text{COMPOSITION} \quad \frac{h_1 \quad h_0}{h} \quad \text{for } h = h_1 \cdot h_0 \\ \\ \text{IDENTITY} \quad \frac{}{\text{id}_A} \end{array}$$

The resulting finitary deductive system (introduced in [6] as a slight modification of the deduction system of Grigore Roşu [19]) has four deduction rules; it is clearly sound, and the main result of our paper (Theorem 6.2) says that it is also complete with respect to finitary morphisms, i.e., morphisms with domain and codomain of finite rank. This implies the expected compactness theorem: every finitary injectivity consequence of a set  $\mathcal{H}$  of finitary morphisms is an injectivity consequence of some finite subset of  $\mathcal{H}$ .

The completeness theorem for Finitary Injectivity Logic will then be extended to the  $k$ -ary Injectivity Logic, defined in the expected way. Then the full completeness theorem easily follows.

The fact that the full Injectivity Logic above is complete in strongly locally ranked categories can also be derived from Quillen’s Small Object Argument [17], see Remark 3.9 below. However our sharpening to the  $k$ -ary logic for every cardinal  $k$  cannot be derived from that paper, and we consider this to be a major step.

**Related work** Bernhard Banaschewski and Horst Herrlich showed thirty years ago that implications in general algebra can be expressed categorically via injectivity w.r.t. regular epimorphisms, see [9]. A generalization to injectivity w.r.t. cones or even trees of morphisms was studied by Hajnal Andréka, István Németi and Ildikó Sain, see e.g. [7, 8, 15].

To see more precisely how that work relates to ours and to classical logic, consider injectivity in the category of all  $\Sigma$ -structures (and  $\Sigma$ -homomorphisms), where  $\Sigma$  is any signature. Then recall from [4], 5.33 that there is a natural way to associate to a (finitary) morphism  $f : A \rightarrow B$  a (finitary) sentence

$$f' := \forall X (\wedge A'(X) \rightarrow \exists Y (\wedge B'(X, Y)))$$

(where  $A'(X)$  and  $B'(X, Y)$  are sets of atomic formulas) such that an object  $C$

satisfies  $f'$  if and only if it is injective with respect to  $f$  (see 2.22 below for more on this). Such sentences are called *regular sentences*. In this paper we concentrate on the proof theory for the (finite and infinite) regular logics. As mentioned above, the restriction to epimorphisms correspond to considering only the quasi-equations (i.e., no existential quantifiers), and just equations if we impose they have projective domains.

Recently, Grigore Roşu introduced a deduction system for injectivity, see [19], and he proved that the resulting logic is sound and complete for epimorphisms which are finitely presentable, see 3.5, and have projective domains. A slight modification of Roşu's system was introduced in [6]: this is the deduction system 2.4 below. It differs from [19] by formulating PUSHOUT more generally and using COMPOSITION in place of Roşu's UNION. In [6] completeness is proved for sets of epimorphisms with finitely presentable domains and codomains. (This is slightly stronger than requiring the epimorphisms to be finitely presentable, however, without the too restrictive assumption of projectivity of the domains the logic fails to be complete for finitely presentable epimorphisms in general, see [6].)

In the present paper completeness of the finitary logic is proved for arbitrary morphisms (not necessarily epimorphisms) with finitely presentable domains and codomains. The fact that the assumption of epimorphism is dropped makes the proof substantially more difficult. We present a short proof in locally presentable categories first, and then a proof of a more general result for strongly locally ranked categories. We also formulate the appropriate infinitary logic dealing with arbitrary morphisms.

There are other generalizations of Birkhoff's equational logic which are, except for the common motivation, not related to our approach. For example the categorical approach to logic of (ordered) many-sorted algebras of Razvan Diaconescu [11], and the logic of implications in general algebra of Robert Quackenbush [16].

In our joint paper [1] we are taking another route to generalize the equational logic: we consider orthogonality of objects to a morphism instead of injectivity. The deduction system is similar: the rule CANCELLATION has to be weakened, and an additional rule concerning coequalizers is added. We prove the completeness of the resulting logic of orthogonality in locally presentable categories. The corresponding sentences are the so called limit sentences,  $\forall X(\wedge A'(X) \rightarrow \exists!Y(\wedge B'(X, Y)))$ , where  $\exists!Y$  means "there exists exactly one  $Y$  such that".

## 2. Logic of injectivity

**2.0. Assumption** Throughout the paper we assume that we are working in a cocomplete category.


**2.1. Definition** A morphism  $h$  is called an *injectivity consequence* of a set of morphisms  $\mathcal{H}$ , notation

$$\mathcal{H} \models h$$


provided that every object injective w.r.t. all morphisms in  $\mathcal{H}$  is also injective w.r.t.  $h$ .

**2.2. Examples** (1) A composite  $h = h_2 \cdot h_1$  is an injectivity consequence of  $\{h_1, h_2\}$ .


(2) Conversely, in every composite  $h = h_2 \cdot h_1$  the morphism  $h_1$  is an injectivity consequence of  $h$ :



(3) In every pushout



$h'$  is an injectivity consequence of  $h$ :



**2.3. Remark** The above examples are exhaustive. More precisely, the following deduction system, introduced in [6], see also [19], (where, however, it was only applied to epimorphisms) will be proved complete below:

**2.4. Definition** The *Finitary Injectivity Deduction System* consists of one axiom

$$\text{IDENTITY} \quad \frac{}{\text{id}_A}$$

and three deduction rules

$$\text{COMPOSITION} \quad \frac{h \quad h'}{h' \cdot h} \quad \text{if } h' \cdot h \text{ is defined}$$

$$\text{CANCELLATION} \quad \frac{h' \cdot h}{h}$$

and

$$\text{PUSHOUT} \quad \frac{h}{h'}$$

We say that a morphism  $h$  is a *formal consequence* of a set  $\mathcal{H}$  of morphisms (notation  $\mathcal{H} \vdash h$ ) in the Finitary Injectivity Logic if there exists a proof of  $h$  from  $\mathcal{H}$  (which means a finite sequence  $h_1, \dots, h_n = h$  of morphisms such that for every  $i = 1, \dots, n$  the morphism  $h_i$  lies in  $\mathcal{H}$  or is a conclusion of one of the deduction rules whose premises lie in  $\{h_1, \dots, h_{i-1}\}$ ).

**2.5. Lemma** *The Finitary Injectivity Logic is sound, i.e., if a morphism  $h$  is a formal consequence of a set of morphisms  $\mathcal{H}$ , then  $h$  is an injectivity consequence of  $\mathcal{H}$ . Briefly:  $\mathcal{H} \vdash h$  implies  $\mathcal{H} \models h$ .*

The proof follows from 2.2.

**2.6. Remark** Later we define finitary morphisms (as morphisms whose domains and codomains are finitely presentable (Section 3) or of finite rank (Section 5)), and in Section 6 we prove that the resulting Finitary Injectivity Logic is complete, i.e., that

$$\mathcal{H} \models h \text{ implies } \mathcal{H} \vdash h$$

for every set  $\mathcal{H}$  of finitary morphisms and every  $h$  finitary.

**2.7. Example** The following rule

$$\text{FINITE COPRODUCT} \quad \frac{h_1 \quad h_2}{h_1 + h_2}$$

(where for  $h_i : A_i \rightarrow B_i$  the morphism  $h_1 + h_2 : A_1 + A_2 \rightarrow B_1 + B_2$  is the canonical coproduct morphism) is obviously sound. Here is a proof in the Finitary Injectivity Logic:

Using the pushouts

we can write


$$\frac{\frac{h_1}{h_1 + \text{id}_{A_2}} \quad h_2}{\text{id}_{B_1} + h_2}}{h_1 + h_2} \quad \begin{array}{l} \text{via PUSHOUT} \\ \text{via COMPOSITION} \end{array}$$

since  $h_1 + h_2 = (\text{id}_{B_1} + h_2) \cdot (h_1 + \text{id}_{A_2})$ .

**2.8. Example** The following rule

$$\text{FINITE WIDE PUSHOUT} \quad \frac{h_1 \dots h_n}{h}$$

for every wide pushout




is sound. Here is a proof in the Finitary Injectivity Logic:

If  $n = 2$  we have

$$\frac{h_1 \quad h_2}{k_2} \quad \text{via PUSHOUT}$$

$$\frac{\quad}{h = k_2 \cdot h_2} \quad \text{via COMPOSITION}$$

If  $n = 3$  denote by  $r$  a pushout of  $h_1, h_2$ , then a pushout,  $h'_3$ ,



of  $h_3$  along  $r$  forms a wide pushout of  $h_1, h_2$  and  $h_3$ :

$$\frac{h_1 \quad h_2 \quad h_3}{k_2} \quad \text{via PUSHOUT}$$

$$\frac{\quad}{r} \quad \text{via COMPOSITION}$$

$$\frac{\quad}{k_3} \quad \text{via PUSHOUT}$$

$$\frac{\quad}{h = k_3 \cdot h_3} \quad \text{via COMPOSITION}$$

Etc.

**2.9. Remark** We want to define a composition of a chain of  $\lambda$  morphisms for every ordinal  $\lambda$  (see the case  $\lambda = \omega$  in the Introduction). Recall that a  $\lambda$ -chain is a functor  $A$  from  $\lambda$ , the well-ordered category of all ordinals  $i < \lambda$ .

Recall further that  $\lambda^+$  denotes the successor ordinal, i.e., the set of all  $i \leq \lambda$ .

**2.10. Definition** (i) We call a  $\lambda$ -chain  $A$  *smooth* if for every limit ordinal  $i < \lambda$  we have

$$A_i = \text{colim}_{j < i} A_j$$

with the colimit cocone of all  $a_{ji} = A(j \rightarrow i)$ .

(ii) A morphism  $h$  is called a  $\lambda$ -composite of morphisms  $(h_i)_{i < \lambda}$ , where  $\lambda$  is an ordinal, if there exists a smooth  $\lambda^+$ -chain  $A$  with connecting morphisms  $a_{ij} : A_i \rightarrow A_j$  for  $i \leq j \leq \lambda$  such that

$$h_i = a_{i,i+1} \quad \text{for all } i < \lambda$$

and

$$h = a_{0,\lambda}.$$

**2.11. Examples**  $\lambda = 0$ : No morphism  $h_i$  is given, just an object  $A_0$ ; and  $h = a_{0,0}$  is the identity morphism of  $A_0$ .

$\lambda = 1$ : A morphism  $h_0$  is given, and we have  $h = a_{0,1} = h_0$ . Thus, a 1-composite of  $h_0$  is  $h_0$ .

$\lambda = 2$ : This is the usual concept of composition: given morphisms  $h_0, h_1$ , their 2-composite exists iff they are composable. Then  $h_1 \cdot h_0$  is the 2-composite.


$\lambda = \omega$ : This is the case mentioned in the Introduction. Observe that, unlike the previous cases, an  $\omega$ -composite is only unique up to isomorphism.

**2.12. Lemma** *A  $\lambda$ -composite of morphisms  $(h_i)_{i < \lambda}$  is an injectivity consequence of these morphisms.*

**Proof** This is a trivial transfinite induction on  $\lambda$ . In case  $\lambda = 0$  this states that  $\text{id}_A$  is an injectivity consequence of  $\emptyset$ , etc.

**2.13. Definition** The *Injectivity Deduction System* consists of the deduction rules

CANCELLATION  $\frac{h' \cdot h}{h}$

PUSHOUT  $\frac{h}{h'}$  for every pushout 

and the rule scheme (one rule for every ordinal  $\lambda$ )

TRANSFINITE COMPOSITION  $\frac{h_i (i < \lambda)}{h}$  for every  $\lambda$ -composite  $h$  of  $(h_i)_{i < \lambda}$

We say that a morphism  $h$  is a *formal consequence* of a set  $\mathcal{H}$  of morphisms (notation  $\mathcal{H} \vdash h$ ) in the Injectivity Logic if there exists a proof of  $h$  from  $\mathcal{H}$  (which means a chain  $(h_i)_{i \leq n}$  of morphisms, where  $n$  is an ordinal, such that  $h = h_n$ , and each  $h_i$  either lies in  $\mathcal{H}$ , or is a conclusion of one of the deduction rules whose premises lie in  $\{h_j\}_{j < i}$ ).

**2.14. Lemma** *The Injectivity Logic is sound, i.e., if a morphism  $h$  is a formal consequence of a set  $\mathcal{H}$  of morphisms, then  $h$  is an injectivity consequence of  $\mathcal{H}$ . Briefly:  $\mathcal{H} \vdash h$  implies  $\mathcal{H} \models h$ .*

The proof (using 2.12) is elementary.




**2.15. Remark** In 2.13 we can replace TRANSFINITE COMPOSITION by the deduction rule WIDE PUSHOUT, see below, which makes use of the (obvious) fact that an object  $A$  injective w.r.t. a set  $\{h_i\}_{i<\lambda}$  of morphisms having a common domain is also injective w.r.t. their wide pushout. Let us note here that this rule does not replace PUSHOUT of 2.13 (because in the latter a pushout of  $h$  along an *arbitrary* morphism is considered).

**2.16. Definition** The deduction rule

$$\text{WIDE PUSHOUT} \quad \frac{h_i \ (i < \lambda)}{h} \quad \text{for } h \text{ a wide pushout of } \{h_i\}_{i<\lambda}$$

applies, for every cardinal  $\lambda$ , to an arbitrary object  $P$  and an arbitrary set  $\{h_i\}$  of  $\lambda$  morphisms with the common domain  $P$  and the following wide pushout



**Remark** Again, this is a scheme of deduction rules: for every cardinal  $\lambda$  we have one rule  $\lambda$ -WIDE PUSHOUT. Observe that  $\lambda = 0$  yields the rule IDENTITY.

**2.17. Lemma** *The Injectivity Deduction System 2.13 is equivalent to the deduction system*

COMPOSITION, CANCELLATION, PUSHOUT and WIDE PUSHOUT.

**Proof** (1) We can derive WIDE PUSHOUT from 2.13. For every ordinal number  $\lambda$  we derive the rule

$$\frac{h_i \ (i < \lambda)}{h} \quad \text{for } h \text{ a wide pushout of } \{h_i\}_{i<\lambda}$$

by transfinite induction on the ordinal  $\lambda$ . We are given an object  $P$  and morphisms  $h_i : P \rightarrow P_i \ (i < \lambda)$ . The case  $\lambda = 0$  is trivial, from  $\lambda$  derive  $\lambda + 1$  by using PUSHOUT, and for limit ordinals  $\lambda$  form the restricted multiple pushouts  $Q_j$  of morphisms  $h_i$  for  $i < j$ , and observe that they form a smooth chain whose composite is a multiple pushout of all  $h_i$ 's.

(2) From the system in 2.17 we can derive the rule  $\lambda$ -COMPOSITION, where  $\lambda$  is an arbitrary ordinal: the case  $\lambda = 0$  follows from 0-WIDE PUSHOUT. The isolated step uses COMPOSITION: the  $(\lambda + 1)$ -composite of  $(h_i)_{i \leq \lambda}$  is simply  $h_\lambda \cdot k$  where  $k$  is the  $\lambda$ -composite of  $(h_i)_{i < \lambda}$ . In the limit case, use the fact that a composite  $h$  of  $(h_i)_{i < \lambda}$  is a wide pushout of  $\{k_i\}_{i < \lambda}$ , where  $k_i$  is a composite of  $(h_j)_{j < i}$ .

**2.18. Remark** For every infinite cardinal  $k$  the  $k$ -ary Injectivity Deduction System is the system 2.13 where  $\lambda$  ranges through ordinals smaller than  $k$ . A proof of a

morphism  $h$  from a set  $\mathcal{H}$  in the  $k$ -ary Injectivity Logic is, then, a proof of length  $n < k$  using only the deduction rules with  $\lambda$  restricted as above. The last lemma can, obviously, be formulated under this restriction in case we use the scheme  $\lambda$ -WIDE PUSHOUT for all cardinals  $\lambda < k$ .

**2.19. Definition** The deduction rule

$$\text{COPRODUCT} \quad \frac{h_i \ (i < \lambda)}{\coprod_{i < \lambda} h_i}$$

applies, for every cardinal  $\lambda$ , to an arbitrary collection of  $\lambda$  morphisms  $h_i : A_i \rightarrow B_i$ .

**2.20. Lemma** *The Injectivity Deduction System 2.13 is equivalent to the deduction system of 2.17 with WIDE PUSHOUT replaced by*

$$\text{IDENTITY} + \text{COPRODUCT}$$

**Proof** (1) COPRODUCT follows from 2.17. In fact,  $\coprod_{i < \lambda} h_i : \coprod_{i < \lambda} A_i \rightarrow \coprod_{i < \lambda} B_i$  is a wide pushout of the morphisms  $k_j : \coprod_{i < \lambda} A_i \rightarrow \coprod_{i < j} A_i + B_j + \coprod_{j < i < \lambda} A_i$ , where  $j$  ranges through  $\lambda$ , with components  $\text{id}_{A_i}$  ( $i \neq j$ ) and  $h_j$ , and  $k_j$  is a pushout of  $h_j$  along the  $j$ -th coproduct injection of  $\coprod_{i < \lambda} A_i$ .

(2) Conversely, WIDE PUSHOUT follows from IDENTITY+COPRODUCT. We obviously need to consider only  $\lambda > 1$  and then we use the fact that given morphisms  $h_i : A \rightarrow B_i$  ( $i < \lambda$ ), their wide pushout  $h : A \rightarrow C$  can be obtained from  $\coprod_{i < \lambda} h_i$  by pushing out along the codiagonal  $\nabla : \coprod_{i < \lambda} A \rightarrow A$ :

$$\begin{array}{ccc} \coprod_{i < \lambda} A & \xrightarrow{\coprod h_i} & \coprod_{i < \lambda} B_i \\ \nabla \downarrow & & \downarrow \\ A & \xrightarrow{h} & C \end{array}$$

**2.21. Remark** The deduction system of the last lemma has five rules, but the advantage against the system 2.13 is that they are particularly simple to formulate:

$$\begin{array}{ll} \text{IDENTITY} & \frac{}{\text{id}_A} \\ \text{CANCELLATION} & \frac{h_2 \cdot h_1}{h_1} \\ \text{COMPOSITION} & \frac{h_2 \ h_1}{h_2 \cdot h_1} \quad \text{if } h_2 \cdot h_1 \text{ is defined} \end{array}$$

$$\begin{array}{l}
 \text{PUSHOUT} \quad \frac{h}{h'} \quad \text{given} \quad \begin{array}{ccc} & \xrightarrow{h} & \\ \downarrow & & \downarrow \\ & \xrightarrow{h'} & \end{array} \\
 \\
 \text{COPRODUCT} \quad \frac{h_i \ (i \in I)}{\coprod_{i \in I} h_i}
 \end{array}$$

We prove below that 2.13, and therefore the above equivalent deduction system, is not only sound but (in a number of categories) also complete.

**2.22. Remark** To relate our deduction rules to the usual ones (of classical logic), let us consider, as in the Introduction, the category of all  $\Sigma$ -structures. Then any object  $A$  can be presented by a set  $A'(X)$  of atomic formulas with parameters  $X$  in  $A$ : for the familiar algebraic structures, this is just the usual concept of generators and relations. Given a morphism  $f : A \rightarrow B$ , and such presentations  $A'(X)$  and  $B'_o(Y)$  of  $A$  and  $B$ , we can also present  $B$  by  $B'(X, Y)$ , which is the union of  $B'_o(Y)$  and the set of all the equations  $x = t(Y)$  for which  $f(x) = t(Y)$  ( $t$  a  $\Sigma$ -term). Then for the sentence

$$f' := \forall X (\wedge A'(X) \rightarrow \exists Y (\wedge B'(X, Y)))$$

we have that an object  $C$  is  $f$ -injective iff  $C \models f'$ . Note that if  $f$  is finitary (see the Introduction or 3.4 below), the presentations, and hence  $f'$ , can be chosen to be finitary (more details in [4], 5.33). Now, we can associate Gentzen-style rules to sets of atomic formulas, generalizing the idea of what was done (with more accuracy) in [6] for sets of equations: associating

$$A'(X) \Rightarrow B'(X, Y)$$

to  $\forall X (\wedge A'(X) \rightarrow \exists Y (\wedge B'(X, Y)))$ , the IDENTITY axiom is of course

$$\frac{}{A'(X) \Rightarrow A'(X)} ;$$

CANCELLATION is a categorical version of the “restriction” rule

$$\frac{A'(X) \Rightarrow (B'(X, Y) \cup C'(X, Y, Z))}{A'(X) \Rightarrow B'(X, Y)} ;$$

PUSHOUT is essentially the “weakening” rule

$$\frac{A'(X) \Rightarrow B'(X, Y)}{(A'(X) \cup C'(X, Z)) \Rightarrow B'(X, Y)} ;$$

and COMPOSITION is a “cut” rule

$$\frac{A'(X) \Rightarrow B'(X, Y), B'(X, Y) \Rightarrow C'(X, Y, Z)}{A'(X) \Rightarrow C'(X, Y, Z)} .$$

The usual stronger “cut” rule

$$\frac{A'(X) \Rightarrow B'(X, Y), ((B'(X, Y) \cup C'(X, Y, Z)) \Rightarrow D'(X, Y, Z, U))}{(A'(X) \cup C'(X, Y, Z)) \Rightarrow D'(X, Y, Z, U)}$$

corresponds to

$$\frac{A \xrightarrow{f} B, \quad B + C \xrightarrow{g} D}{A + C \xrightarrow{g \cdot (f + 1_C)} C},$$

which is proved via

$$\frac{\frac{f \quad g}{f + \text{id}_C \quad g}}{g \cdot (f + 1_C)} \quad \begin{array}{l} \text{PUSHOUT} \\ \text{COMPOSITION} \end{array}$$

### 3. Completeness in locally presentable categories

**3.1. Assumption** In the present section we study injectivity in a *locally presentable* category  $\mathcal{A}$  of Gabriel and Ulmer, see [12] or [4]. This means that:

- (a)  $\mathcal{A}$  is cocomplete,

and

- (b) there exists a regular cardinal  $\lambda$  such that  $\mathcal{A}$  has a set of  $\lambda$ -presentable objects whose closure under  $\lambda$ -filtered colimits is all of  $\mathcal{A}$ .

Recall that an object  $A$  is  *$\lambda$ -presentable* if its hom-functor  $\text{hom}(A, -) : \mathcal{A} \rightarrow \mathbf{Set}$  preserves  $\lambda$ -filtered colimits. That is, given a  $\lambda$ -filtered diagram  $D$  with a colimit  $c_i : D_i \rightarrow C$  ( $i \in I$ ) in  $\mathcal{A}$ , then for every morphism  $f : A \rightarrow C$

- (i) a factorization of  $f$  through  $c_i$  exists for some  $i \in I$ ,

and

- (ii) factorizations are essentially unique, i.e., given  $i \in I$  and  $c_i \cdot g' = c_i \cdot g''$  for some  $g', g'' : A \rightarrow D_i$ , there exists a connecting morphism  $d_{ij} : D_i \rightarrow D_j$  of the diagram with  $d_{ij} \cdot g' = d_{ij} \cdot g''$ .

**3.2. Examples** (see [4]) Sets, presheaves, varieties of algebras and simplicial sets are examples of locally presentable categories. Categories such as **Top** (topological spaces) or **Haus** (Hausdorff spaces) are not locally presentable.

**3.3. Remark** (a) In the present section we prove that the Injectivity Logic is complete in every locally presentable category. The reader may decide to skip this section since we prove a more general result in Section 6. Both of our proofs are based on the fact that for every set  $\mathcal{H}$  of morphisms the full subcategory  $\text{Inj}\mathcal{H}$  (of all objects injective w.r.t. morphisms of  $\mathcal{H}$ ) is *weakly reflective*. That is: every object  $A \in \mathcal{A}$  has a morphism  $r : A \rightarrow \bar{A}$ , called a weak reflection, such that

- (i)  $\bar{A}$  lies in  $\text{Inj}\mathcal{H}$

and

- (ii) every morphism from  $A$  to an object of  $\text{Inj}\mathcal{H}$  factors through  $r$  (not necessarily uniquely).

In the present section we will utilize the classical *Small Object Argument* of D. Quillen [17]: this tells us that every object  $A$  has a weak reflection  $r : A \rightarrow \bar{A}$  in  $\text{Inj}\mathcal{H}$  such that  $r$  is a transfinite composite of morphisms of the class

$$\widehat{\mathcal{H}} = \{k; k \text{ is a pushout of a member of } \mathcal{H} \text{ along some morphism}\}.$$

(b) The reason for proving the completeness based on the Small Object Argument in the present section is that the proof is short and elegant. However, by using a more refined construction of weak reflection in  $\text{Inj}\mathcal{H}$ , which we present in Section 5, we will be able to prove the completeness in the so-called strongly locally ranked categories, which include **Top** and **Haus**.

The spirits of the two proofs are quite different. Given an injectivity consequence  $h$  of a set of morphisms, in this section we will show how to derive a formal proof of  $h$  from Quillen’s construction of the weak reflection; this construction is “linear”, forming a transfinite composite. In the next section, a weak reflection will be constructed as a colimit of a filtered diagram which somehow presents simultaneously all the possible formal proofs.

**3.4. Definition** A morphism is called  $\lambda$ -ary provided that its domain and codomain are  $\lambda$ -presentable objects. For  $\lambda = \aleph_0$  we say *finitary*.

**3.5. Remark** (a) The  $\lambda$ -ary morphisms are precisely the  $\lambda$ -presentable objects of the arrow category  $\mathcal{A}^\rightarrow$ . In contrast, M. Hébert introduced in [13]  $\lambda$ -presentable morphisms; these are the morphisms  $f : A \rightarrow B$  which are  $\lambda$ -presentable objects of the slice category  $A \downarrow \mathcal{A}$ . In the present paper we will not use the latter concept.

(b) We work now with the *Finitary Injectivity Logic*, i.e., the deduction system 2.4 applied to finitary morphisms. We generalize this to the  $k$ -ary logic below.

**3.6. Theorem** *The Finitary Injectivity Logic is complete in every locally presentable category  $\mathcal{A}$ . That is, given a set  $\mathcal{H}$  of finitary morphisms in  $\mathcal{A}$ , then every finitary morphism  $h$  which is an injectivity consequence of  $\mathcal{H}$  is a formal consequence in the deduction system 2.4. Briefly:*

$$\mathcal{H} \models h \text{ implies } \mathcal{H} \vdash h.$$


**Proof** Given a finitary morphism  $h : A \rightarrow B$  which is an injectivity consequence of  $\mathcal{H}$ , we prove that

$$\mathcal{H} \vdash h.$$

(a) The above object  $A$  has a weak reflection

$$r : A \rightarrow \bar{A}$$


such that  $r$  is a transfinite composition of morphisms in  $\widehat{\mathcal{H}}$ , see 3.3(a). Since  $\mathcal{H} \models h$ , it follows that  $\bar{A}$  is injective w.r.t.  $h$ , which yields a morphism  $u$  forming a commutative triangle



(b) Consider all commutative triangles as above where  $r : A \rightarrow \bar{A}$  is any  $\alpha$ -composite of morphisms in  $\widehat{\mathcal{H}}$  for some ordinal  $\alpha$  and  $u$  is arbitrary. We prove that the least possible  $\alpha$  is finite. This finishes the proof of  $\mathcal{H} \vdash h$ : In case  $\alpha = 0$ , we have that  $\text{id} = u \cdot h$ , and we derive  $h$  via IDENTITY and CANCELLATION. In case  $\alpha$  is a finite ordinal greater than 0, we have that  $r$  is provable from  $\mathcal{H}$  using PUSHOUT and COMPOSITION. Consequently, via CANCELLATION, we get  $h$ .

Let  $\mathcal{C}$  be the class of all ordinals  $\alpha$  such that there are an  $\alpha$ -composite  $r$  of morphisms of  $\widehat{\mathcal{H}}$  and a morphism  $u$  with  $r = u \cdot h$ . To show that the least member  $\gamma$  of  $\mathcal{C}$  is finite, we prove that for each ordinal  $\gamma \geq \omega$  in  $\mathcal{C}$  we can find another ordinal in  $\mathcal{C}$  which is smaller than  $\gamma$ .

A. Case  $\gamma = \beta + m$ , with  $\beta$  a limit ordinal and  $m > 0$  finite. Let  $a_{i,i+1}$  ( $i < \beta + m$ ) be the corresponding chain with  $r = a_{0,\beta+m}$ . Since  $a_{\beta,\beta+1}$  lies in  $\widehat{\mathcal{H}}$ , we can express it as a pushout of some morphism  $k : D \rightarrow D'$  in  $\mathcal{H}$ :



We have a colimit  $A_\beta = \text{colim}_{i < \beta} A_i$  of a chain of morphisms. Hence, because  $D$  is finitely presentable,  $p$  factorizes as  $p = a_{i\beta} \cdot q$  for some  $i < \beta$  and some morphism  $q : D \rightarrow A_i$ . Let  $v_i$  be a pushout of  $k$  along  $q$ , and form a sequence  $v_j$  of pushouts of  $k$  along  $a_{ij} \cdot q$  ( $j < \beta$ ) as illustrated in the diagram above (taking colimits at the limit ordinals). Then it is easily seen, due to  $p = a_{i\beta} \cdot q$ , that  $v_\beta = \text{colim}_{j < \beta} v_j$  is a pushout of  $k$  along  $p$ . Thus, without loss of generality,

$$P_\beta = A_{\beta+1} \quad \text{and} \quad v_\beta = a_{\beta,\beta+1}.$$

Observe that, since  $a_{j,j+1}$  lies in  $\widehat{\mathcal{H}}$ , PUSHOUT implies that

$$p_{j,j+1} \in \widehat{\mathcal{H}} \quad \text{for all } i \leq j < \beta.$$

Also  $v_i \in \widehat{\mathcal{H}}$  since it is a pushout of  $k$  along  $q$ . Consequently,  $a_{0,\beta+1}$  is a  $\beta$ -composite of morphisms  $b_{j,j+1}$  ( $j < \beta$ ) of  $\widehat{\mathcal{H}}$  as follows (where  $l$  is the first limit ordinal after  $i$ ):

$$\begin{aligned} b_{j,j+1} &= a_{j,j+1} && \text{for all } j < i, \\ b_{i,i+1} &= v_i, \\ b_{j,j+1} &= p_{j-1,j} && \text{for all } i < j < l, \end{aligned}$$

and

$$b_{j,j+1} = p_{j,j+1} \quad \text{for all } l \leq j < \beta.$$

Thus  $r = a_{0,\beta+m}$  is a  $(\beta + (m - 1))$ -composite of morphisms of  $\widehat{\mathcal{H}}$ .

B. Case  $\gamma$  is a limit ordinal. The morphism

$$u : B \rightarrow \overline{A} = \operatorname{colim}_{i < \gamma} A_i$$

factors, since  $B$  is finitely presentable, through some  $a_{i\gamma}$ ,  $i < \gamma$ :

$$u = a_{i\gamma} \cdot \overline{u} \text{ for some } \overline{u} : B \rightarrow A_i.$$

The parallel pair

$$A = A_0 \begin{array}{c} \xrightarrow{\overline{u} \cdot h} \\ \xrightarrow{a_{0i}} \end{array} A_i$$

is clearly merged by the colimit morphism  $a_{i\gamma}$  of  $A_\gamma = \operatorname{colim}_{i < \gamma} A_i$ . Since  $A$  is finitely presentable,  $\operatorname{hom}(A, -)$  preserves that colimit, consequently (see (ii) in 3.1.b), the parallel pair is also merged by a connecting morphism  $a_{ij} : A_i \rightarrow A_j$  for some  $i < j < \gamma$ :

$$a_{ij} \cdot \overline{u} \cdot h = a_{0j}.$$

This gives us a commutative triangle

$$\begin{array}{ccccc} A_0 & \xrightarrow{a_{01}} & A_1 & \xrightarrow{a_{12}} & \cdots & \rightarrow & A_j \\ & \searrow h & & & & \nearrow a_{ij} \cdot \overline{u} & \\ & & B & & & & \end{array}$$

thus  $a_{0j}$  is a  $j$ -composite of morphisms of  $\widehat{\mathcal{H}}$  with  $j < \gamma$ .

**3.7. Remark** The above theorem immediately generalizes to the  $k$ -ary Injectivity Logic, i.e., to the deduction system of 2.18 applied to  $k$ -ary morphisms. Recall that for every set of objects in a locally presentable category there exists a cardinal  $k$  such that all these objects are  $k$ -presentable. Consequently, for every set  $\mathcal{H} \cup \{h\}$  of morphisms there exists  $k$  such that all members are  $k$ -ary. The proof that  $\mathcal{H} \models h$  implies  $\mathcal{H} \vdash h$  is completely analogously to 3.6: We show that the least possible  $\alpha$  is smaller than  $k$ , thus in Cases A. and B. we work with  $\gamma \geq k$ .

**3.8. Corollary** *The Injectivity Logic is sound and complete in every locally presentable category.*

In fact, given

$$\mathcal{H} \models h$$

find a cardinal  $k$  such that all members of  $\mathcal{H} \cup \{h\}$  are  $k$ -ary morphisms. Then  $h$  is a formal consequence of  $\mathcal{H}$  by 3.7.

**3.9. Remark** The above corollary also follows from the Small Object Argument (see 3.3(a)): if  $h : A \rightarrow B$  is an injectivity consequence of  $\mathcal{H}$  and if  $r : A \rightarrow \overline{A}$  is the corresponding weak reflection, then  $r$  is clearly a formal consequence of  $\mathcal{H}$ . Since  $\overline{A}$  is injective w.r.t.  $h$ , it follows that  $r$  factors through  $h$ , thus,  $h$  is a formal consequence of  $r$  (via CANCELLATION).

## 4. Strongly locally ranked categories

**4.1. Remark** Recall that a *factorization system* in a category is a pair  $(\mathcal{E}, \mathcal{M})$  of classes of morphisms containing all isomorphisms and closed under composition such that

- (a) every morphism  $f : A \rightarrow B$  has a factorization  $f = m \cdot e$  with  $e : A \rightarrow C$  in  $\mathcal{E}$  and  $m : C \rightarrow B$  in  $\mathcal{M}$

and

- (b) given another such factorization  $f = m' \cdot e'$  there exists a unique “diagonal fill-in” morphism  $d$  making the diagram

$$\begin{array}{ccc}
 A & \xrightarrow{e} & C \\
 e' \downarrow & \swarrow d & \downarrow m \\
 C' & \xrightarrow{m'} & B
 \end{array}$$

commutative.

The factorization system is called *left-proper* if every morphism of  $\mathcal{E}$  is an epimorphism. In that case the  $\mathcal{E}$ -quotients of an object  $A$  are the quotient objects of  $A$  represented by morphisms of  $\mathcal{E}$  with domain  $A$ .

**4.2. Definition** Let  $(\mathcal{E}, \mathcal{M})$  be a factorization system. We say that an object  $A$  has  $\mathcal{M}$ -rank  $\lambda$ , where  $\lambda$  is a regular cardinal, provided, that

- (a)  $\text{hom}(A, -)$  preserves  $\lambda$ -filtered colimits of diagrams of  $\mathcal{M}$ -morphisms (i.e., given a  $\lambda$ -filtered diagram  $D$  whose connecting morphisms lie in  $\mathcal{M}$ , then every morphism  $f : A \rightarrow \text{colim} D$  factors, essentially uniquely, through a colimit map of  $D$ )

and

- (b)  $A$  has less than  $\lambda$   $\mathcal{E}$ -quotients.

If  $\lambda = \aleph_0$  we say that the object  $A$  has *finite  $\mathcal{M}$ -rank*.

**4.3. Examples** (1) For the factorization system  $(\text{Iso}, \text{All})$ , rank  $\lambda$  is equivalent to  $\lambda$ -presentability.

(2) In the category **Top** of topological spaces, choose  $(\mathcal{E}, \mathcal{M}) = (\text{Epi}, \text{Strong Mono})$ . Here the  $\mathcal{M}$ -subobjects are precisely the embeddings of subspaces. Every topological space  $A$  of cardinality  $\alpha$  has  $\mathcal{M}$ -rank  $\lambda$  whenever  $\lambda > 2^{2^\alpha}$ . In fact,  $\text{hom}(A, -)$  preserves  $\lambda$ -directed unions of subspaces since  $\alpha < \lambda$ . And the amount of quotient objects of  $A$  (carried by epimorphisms) is at most  $\sum_{\beta \leq \alpha} E_\beta T_\beta$  where  $E_\beta$  is the number of equivalence relations on  $A$  of order  $\beta$  and  $T_\beta$  is the number of topologies on a set of cardinality  $\beta$ . Since  $E_\beta$  and  $T_\beta$  are both  $\leq 2^{2^\beta}$ , we have  $\sum_{\beta \leq \alpha} E_\beta T_\beta \leq \alpha \cdot 2^{2^\alpha} \cdot 2^{2^\alpha} < \lambda$ , thus we conclude that  $A$  has less than  $\lambda$  quotients.

**4.4. Remark** Every  $\mathcal{E}$ -quotient of an object of  $\mathcal{M}$ -rank  $\lambda$  also has  $\mathcal{M}$ -rank  $\lambda$ . In fact (a) in 4.2 follows easily by diagonal fill-in, and (b) is obvious.



**4.5. Definition** A category  $\mathcal{A}$  is called *strongly locally ranked* provided that it has a left-proper factorization system  $(\mathcal{E}, \mathcal{M})$  such that

- (i)  $\mathcal{A}$  is cocomplete;
- (ii) every object has an  $\mathcal{M}$ -rank, and all objects of the same  $\mathcal{M}$ -rank form a set up to isomorphism;
- (iii) for every cardinal  $\mu$  the collection of all objects of  $\mathcal{M}$ -rank  $\mu$  is closed under  $\mathcal{E}$ -quotients and under  $\mu$ -small colimits, i.e., colimits of diagrams with less than  $\mu$  morphisms;

and

- (iv) the subcategory of all objects of  $\mathcal{A}$  and all morphisms of  $\mathcal{M}$  is closed under filtered colimits in  $\mathcal{A}$ .

**Remark** The statement (iv) means that, given a filtered colimit with connecting morphisms in  $\mathcal{M}$ , then

- (a) the colimit cocone is formed by morphisms of  $\mathcal{M}$

and

- (b) every other cocone of  $\mathcal{M}$ -morphisms has the unique factorizing morphism in  $\mathcal{M}$ .

**4.6. Examples** (1) Every locally presentable category is strongly locally ranked: choose

$$\mathcal{E} \equiv \text{isomorphisms, } \mathcal{M} \equiv \text{all morphisms.}$$

In fact, see [4], 1.9 for the proof of (ii), whereas (iii) and (iv) hold trivially.

(2) Choose

$$\mathcal{E} \equiv \text{epimorphisms, } \mathcal{M} \equiv \text{strong monomorphisms.}$$

Here categories such as **Top** (which are not locally presentable) are included. In fact, for a space  $A$  of cardinality  $\alpha$  we have that  $\text{hom}(A, -)$  preserves  $\lambda$ -filtered colimits (=unions) of subspaces whenever  $\alpha < \lambda$ . Thus, by choosing a cardinal  $\lambda > \alpha$  bigger than the number of quotients of  $A$  we get an  $\mathcal{M}$ -rank of  $A$ . It is easy to verify (iii) and (iv) in **Top**.

(3) Let  $\mathcal{B}$  be a full, isomorphism closed,  $\mathcal{E}$ -reflective subcategory of a strongly locally ranked category  $\mathcal{A}$ . If  $\mathcal{B}$  is closed under filtered colimits of  $\mathcal{M}$ -morphisms in  $\mathcal{A}$ , then  $\mathcal{B}$  is strongly locally ranked. In fact,  $\mathcal{B}$  is closed under  $\mathcal{M}$  in the sense that given  $m : A \rightarrow B$  in  $\mathcal{M}$  with  $B \in \mathcal{B}$ , then  $A \in \mathcal{B}$ . (Indeed, we have a reflection  $r_A : A \rightarrow A'$  in  $\mathcal{E}$  and  $m = m' \cdot r_A$  for a unique  $m'$ ; this implies that  $r_A \in \mathcal{E}$  is an isomorphism, thus,  $A \in \mathcal{B}$ .) Therefore the restriction of  $(\mathcal{E}, \mathcal{M})$  to  $\mathcal{B}$  yields a factorization system. It fulfils (ii)-(iv) of 4.5 because  $\mathcal{B}$  is closed under filtered colimits of  $\mathcal{M}$ -morphisms.

(4) The category **Haus** of Hausdorff spaces is strongly locally ranked: it is an epireflective subcategory of **Top** closed under filtered unions of subspaces.

**4.7. Observation** In a strongly locally ranked category the class  $\mathcal{M}$  is closed under transfinite composition. This follows from (iv).

**4.8. Definition** A morphism is called  $k$ -ary if its domain and codomain have  $\mathcal{M}$ -rank  $k$ . In case  $k = \aleph_0$  we speak of *finitary morphisms*.

**4.9. Remark** The name “strongly locally ranked” was chosen since our requirements are somewhat stronger than those of [2]: there a category is called locally ranked in case it is cocomplete, has an  $(\mathcal{E}, \mathcal{M})$ -factorization, is  $\mathcal{E}$ -cowellpowered and for every object  $A$  there exists an infinite cardinal  $\lambda$  such that  $\text{hom}(A, -)$  preserves colimits of  $\lambda$ -chains of  $\mathcal{M}$ -monomorphisms. Our definition of rank and the condition 4.5(ii) imply that the given category is  $\mathcal{E}$ -cowellpowered. Thus, every strongly locally ranked category is locally ranked.

An example of a locally ranked category that is not strongly locally ranked is the category of  $\sigma$ -semilattices (posets with countable joins and functions preserving them): condition 4.5(iv) fails here. Consider e.g. the  $\omega$ -chain of the posets  $\text{exp}(n)$  (where  $n = \{0, 1, \dots, n - 1\}$ ),  $n \in \omega$ , with inclusion as order. The colimit of this chain is  $\text{exp}(\mathbb{N})$  ordered by inclusion. If  $M$  is the poset of all finite subsets of  $\mathbb{N}$  with an added top element, then the embeddings  $\text{exp}(n) \hookrightarrow M$  form a cocone of the chain, but the factorization morphism  $\text{exp}(\mathbb{N}) \rightarrow M$  is not a monomorphism.

## 5. A construction of weak reflections

**5.1. Assumption** In the present section  $\mathcal{A}$  denotes a strongly locally ranked category. For every infinite cardinal  $k$ ,  $\mathcal{A}_k$  denotes a chosen set of objects of  $\mathcal{M}$ -rank  $k$  closed under  $\mathcal{E}$ -quotients and  $k$ -small colimits. In particular, one may of course choose  $\mathcal{A}_k$  to be a set of representatives of all the objects of  $\mathcal{M}$ -rank  $k$  up to isomorphism.

Given a set  $\mathcal{H} \subseteq \mathcal{M}$  of  $k$ -ary morphisms of  $\mathcal{A}_k$  (considered as a full subcategory of  $\mathcal{A}$ ), [2] provides a construction of a weak reflection in  $\text{Inj } \mathcal{H}$ , which generalizes the Small Object Argument (see 3.3). However, this does not appear to be sufficient to prove our Completeness Theorem for the finitary case. The aim of this section is to present a different, more appropriate construction.

We begin with the case  $k = \omega$  and come back to the general case at the end of this section.

**5.2. Convention** (a) Morphisms with domain and codomain in  $\mathcal{A}_\omega$  are called *petty*.  
 (b) Given a set  $\mathcal{H}$  of petty morphisms, let

$$\overline{\mathcal{H}}$$

denote the closure of  $\mathcal{H}$  under finite composition and pushout in  $\mathcal{A}_\omega$ . (That is,  $\overline{\mathcal{H}}$  is the closure of  $\mathcal{H} \cup \{\text{id}_A; A \in \mathcal{A}_\omega\}$  under binary composition and pushout along petty morphisms.)

(c) Since  $\overline{\mathcal{H}} \subseteq \text{mor } \mathcal{A}_\omega$  is a set, we can, for every object  $B$  of  $\mathcal{A}_\omega$ , index all morphisms of  $\overline{\mathcal{H}}$  with domain  $B$  by a set – and that indexing set can be chosen to be independent of  $B$ . That is, we assume that a set  $T$  is given and that for every object  $B \in \mathcal{A}_\omega$ ,

$$\{h_B(t) : B \rightarrow B(t) ; t \in T\} \tag{5.1}$$

is the set of all morphisms of  $\overline{\mathcal{H}}$  with domain  $B$ .

**5.3. Diagram  $D_A$**  For every object  $A \in \mathcal{A}_\omega$  we define a diagram  $D_A$  in  $\mathcal{A}$  and later prove that a weak reflection of  $A$  in  $\text{Inj } \mathcal{H}$  is obtained as a colimit of  $D_A$ . The domain  $\mathcal{D}$  of  $D_A$ , independent of  $A$ , is the poset of all finite words

$$\varepsilon, M_1, M_1M_2, \dots, M_1 \dots M_k \quad (k < \omega)$$

where  $\varepsilon$  denotes the empty word and each  $M_i$  is a finite subset of  $T$ . The ordering is as follows:

$$M_1 \dots M_k \leq N_1 \dots N_l \quad \text{iff} \quad k \leq l \quad \text{and} \quad M_1 \subseteq N_1, \dots, M_k \subseteq N_k.$$

Observe that  $\varepsilon$  is the least element.

We denote the objects  $D_A(M_1 \dots M_k)$  of the diagram  $D_A$  by

$$A_M \quad \text{where} \quad M = M_1 \dots M_k,$$

and if  $M_1 \dots M_k \leq N_1 \dots N_l = N$ , we denote by

$$a_{M,N} : A_M \rightarrow A_N$$


the corresponding connecting morphism of  $D_A$ . We define these objects and connecting morphisms by induction on the length  $k$  of the word  $M = M_1 \dots M_k$  considered.

*Case  $k = 0$ :*  $A_\varepsilon = A$ .

*Induction step:* Assume that all objects  $A_M$  with  $M$  of length less than or equal to  $k$  and all connecting morphisms between them are defined. For every word  $M$  of length  $k + 1$  denote by

$$M^* \leq M$$

the prefix of  $M$  of length  $k$ , and define the object  $A_M$  as a colimit of the following finite diagram



where  $K$  ranges over all words  $K \in \mathcal{D}$  with  $K \leq M^*$  and  $t$  ranges over the set  $M_{k+1}$ . Thus,  $A_M$  is equipped with (the universal cone of) morphisms

$$a_{M^*,M} : A_{M^*} \rightarrow A_M \quad (\text{connecting morphism of } D_A)$$

and

$$d_M^K(t) : A_K(t) \rightarrow A_M \quad \text{for all } K \leq M^*, t \in M_{k+1},$$

forming commutative squares

$$\begin{array}{ccc}
 & A_K & \\
 a_{K,M^*} \swarrow & & \searrow h_{A_K}(t) \\
 A_{M^*} & & A_K(t) \\
 a_{M^*,M} \searrow & & \swarrow d_M^K(t) \\
 & A_M &
 \end{array} \tag{5.2}$$

This defines the objects  $A_M$  for all words of length  $k + 1$ . Next we define connecting morphisms

$$a_{N,M} : A_N \rightarrow A_M$$

for all words  $N \leq M$ . If the length of  $N$  is at most  $k$ , then  $N \leq M^*$  and we define  $a_{N,M}$  through the (already defined) connecting morphism  $a_{N,M^*}$  by composing it with the above  $a_{M^*,M}$ . If  $N$  has length  $k + 1$ , we define  $a_{N,M}$  as the unique morphism for which the diagrams

$$\begin{array}{ccc}
 & A_K & \\
 a_{K,N^*} \swarrow & & \searrow h_{A_K}(t) \\
 A_{N^*} & & A_K(t) \\
 a_{N^*,N} \searrow & & \swarrow d_N^K(t) \\
 & A_N & \\
 a_{N^*,M} \swarrow & & \searrow d_M^K(t) \\
 & A_M &
 \end{array} \tag{5.3}$$

$(K \leq N^*, t \in N_{k+1})$

commute.

It is easy to verify that the morphisms  $a_{N,M}$  are well-defined and that  $D_A : \mathcal{D} \rightarrow \mathcal{A}$  preserves composition and identity morphisms.

**5.4. Lemma** *All connecting morphisms of the diagram  $D_A$  lie in  $\overline{\mathcal{H}}$ .*

**Proof** We first observe that, given a finite diagram

$$\begin{array}{ccc}
 A_i & \xrightarrow{h_i} & B_i \\
 f_i \downarrow & & \\
 C & &
 \end{array} \tag{i \in I}$$

with all  $h_i$  in  $\overline{\mathcal{H}}$ , a colimit

$$\begin{array}{ccc}
 A_i & \xrightarrow{h_i} & B_i \\
 f_i \downarrow & & \downarrow d_i \\
 C & \xrightarrow{h} & D
 \end{array} \quad (i \in I) \tag{5.4}$$

is obtained by first considering pushouts  $h'_i$  of  $h_i$  along  $f_i$  and then forming a wide pushout  $h$  of all  $h'_i$  ( $i \in I$ ). Consequently, the connecting morphisms of  $D_A$  are formed by repeating one of the following steps: a finite wide pushout of morphisms in  $\overline{\mathcal{H}}$ , a composition of morphisms in  $\overline{\mathcal{H}}$ , and a pushout of a morphism in  $\overline{\mathcal{H}}$  along a petty morphism. Since  $\overline{\mathcal{H}}$  is closed, by 5.2, under the latter, it is closed under the first one in the obvious sense, see the construction of a finite wide pushout described in Example 2.8.

**5.5. Lemma** *For every object  $A_M$  of the diagram  $D_A$  and every morphism  $h : A_M \rightarrow B$  of  $\overline{\mathcal{H}}$  there exists a connecting morphism  $a_{M,N} : A_M \rightarrow A_N$  of  $D_A$  which factors through  $h$ .*

**Proof** We have  $M = M_1 \dots M_k$  and  $h = h_{A_M}(t)$  for some  $t \in T$ . Put

$$N = M_1 \dots M_k \{t\}.$$

Then for  $K = M$  the definition of  $d_N^K(t)$  (see (5.2)) gives the following commutative diagram:

$$\begin{array}{ccc}
 A_M & \xrightarrow{h_{A_M}(t)} & A_M(t) \\
 id \downarrow & & \downarrow d_N^K(t) \\
 A_M & \xrightarrow{a_{M,N}} & A_N
 \end{array}$$

Consequently,

$$a_{M,N} = d_N^K(t) \cdot h_{A_M}(t)$$

as required.


**5.6. Proposition** *Let  $\mathcal{H}$  be a set of petty morphisms with  $\overline{\mathcal{H}} \subseteq \mathcal{M}$ . Then for every object  $A \in \mathcal{A}_w$  a colimit  $\gamma_M : A_M \rightarrow \hat{A}$  ( $M \in \mathcal{D}$ ) of the diagram  $D_A$  yields a weak reflection of  $A$  in  $Inj\mathcal{H}$  via*

$$r_A = \gamma_\varepsilon : A \rightarrow \hat{A}.$$

**Proof** (1)  $\hat{A}$  is injective w.r.t.  $\mathcal{H}$ : We want to prove that given  $h \in \mathcal{H}$  and  $f$  as follows

$$\begin{array}{ccc}
 B & \xrightarrow{h} & C \\
 f \downarrow & & \\
 \hat{A} & & 
 \end{array}$$

then  $f$  factors through  $h$ . Firstly, since  $\hat{A} = \text{colim} D_A$  is a directed colimit of  $\overline{\mathcal{H}}$ -morphisms (see 5.4) with  $\overline{\mathcal{H}} \subseteq \mathcal{M}$ , and  $B$  has finite  $\mathcal{M}$ -rank (because  $B \in \mathcal{A}_\omega$ ), it follows that  $\text{hom}(B, -)$  preserves the colimit of  $D_A$ . Thus, there exists a colimit morphism  $\gamma_N : A_N \rightarrow \hat{A}$  through which  $f$  factors,  $f = \gamma_N \cdot f'$ .




By pushing  $h \in \mathcal{H}$  out along  $f'$  we obtain a morphism  $h' \in \overline{\mathcal{H}}$ . Then by 5.5 there exists  $M \geq N$  such that  $a_{N,M} = h'' \cdot h'$  for some  $h'' : A_N(t) \rightarrow A_M$ . The above commutative diagram proves that  $f$  factors through  $h$ .

(2) Let  $B$  be injective w.r.t.  $\mathcal{H}$ . For every morphism  $f : A \rightarrow B$  we define a compatible cocone  $f_M : A_M \rightarrow B$  of the diagram  $D_A$  by induction on


$$k = \text{the length of the word } M$$

such that  $f_\varepsilon = f$ . Then the desired factorization of  $f$  is obtained via the (unique) factorization  $g : \hat{A} \rightarrow B$  with  $g \cdot \gamma_M = f_M$ : in fact,  $g \cdot r_A = f$ .

For  $k \mapsto k + 1$ , choose for every word  $N$  of length  $k$  and every  $t \in T$  a morphism  $f_N(t)$  forming a commutative triangle



(recalling that  $B$  is  $\overline{\mathcal{H}}$ -injective because it is  $\mathcal{H}$ -injective). Then for every word  $M$  of length  $k + 1$  we have a unique factorization  $f_M : A_M \rightarrow B$  making the following diagrams



commutative for all  $K \leq M^*$  and  $t \in M_{k+1}$ .

Let us verify the compatibility

$$f_M = f_N \cdot a_{M,N} \quad \text{for all } M \leq N \text{ in } \mathcal{D}. \tag{5.6}$$

The last diagram yields  $f_{M^*} = f_M \cdot a_{M^*, M}$ . Therefore, it is sufficient to prove (5.6) for words  $M$  and  $N$  of the same length  $k + 1$ . In order to do that, we will show that

$$f_M \cdot d_M^K(t) = f_N \cdot a_{M,N} \cdot d_M^K(t), \text{ for all } K \leq M^* \text{ and } t \in M_{k+1}, \quad (5.7)$$

and

$$f_M \cdot a_{M^*, M} = f_N \cdot a_{M,N} \cdot a_{M^*, M}. \quad (5.8)$$

Concerning (5.7), we have

$$\begin{aligned} f_M \cdot d_M^K(t) &= f_K(t) \\ &= f_N \cdot d_N^K(t), \text{ by replacing } M \text{ by } N \text{ in (5.5)} \\ &= f_N \cdot a_{M,N} \cdot d_M^K(t), \text{ by (5.3)}. \end{aligned}$$

As for (5.8), we have

$$\begin{aligned} f_M \cdot a_{M^*, M} &= f_{M^*} \\ &= f_{N^*} \cdot a_{M^*, N^*} \\ &= f_N \cdot a_{N^*, N} \cdot a_{M^*, N^*}, \text{ by replacing } M \text{ by } N \text{ in (5.5)} \\ &= f_N \cdot a_{M,N} \cdot a_{M^*, M}. \end{aligned}$$

**5.7. Convention** Generalizing the above construction from  $\omega$  to any infinite cardinal  $k$ , we call the morphisms of  $\mathcal{A}_k$  *k-petty*. Let us now denote by

$$\overline{\mathcal{H}}_k$$

the closure of  $\mathcal{H}$  under  $k$ -composition (2.10) and pushout in  $A_k$ . Following 2.18,  $\overline{\mathcal{H}}_k$  is closed under  $k$ -wide pushout. We again assume that a set  $T$  is given such that, for every object  $B \in \mathcal{A}_k$  we have an indexing  $h_B(t) : B \rightarrow B(t)$ ,  $t \in T$  of all morphisms of  $\overline{\mathcal{H}}_k$  with domain  $B$ .

**5.8. Diagram  $\mathbf{D}_A$**  The poset  $\mathcal{D}$  of 5.3 is generalized to a poset  $\mathcal{D}_k$ : Let  $\mathcal{P}_k T$  be the poset of all subsets of  $T$  of cardinality  $< k$ . The elements of  $\mathcal{D}_k$  are all functions

$$M : \lambda \rightarrow \mathcal{P}_k T$$

where  $\lambda < k$  is an ordinal, including the case  $\varepsilon : 0 \rightarrow \mathcal{P}_k T$ . The ordering is as follows: for  $N : \lambda' \rightarrow \mathcal{P}_k T$  put

$$M \leq N \text{ iff } \lambda \leq \lambda' \text{ and } M_i \subseteq N_i \text{ for all } i < \lambda.$$

We define, for every  $A \in \mathcal{A}_k$ , the diagram  $D_A : \mathcal{D}_k \rightarrow \mathcal{A}$ . The objects  $D_A(M) = A_M$  and the connecting morphisms  $a_{M,N} : A_M \rightarrow A_N$  ( $M \leq N$ ) are defined by transfinite induction on  $\lambda < k$ . For  $\lambda = 0$  we have  $A_\varepsilon = A$ . The isolated step is precisely as in 5.3, where for  $M : \lambda + 1 \rightarrow \mathcal{P}_k T$  we denote by  $M^* : \lambda \rightarrow \mathcal{P}_k T$  the domain-restriction. The limit steps are defined via colimits of smooth chains, see 2.10: if  $\lambda < k$  is a limit ordinal and  $M : \lambda \rightarrow \mathcal{P}_k T$  is given, then  $A_M$  is a colimit

of the chain  $A_{M/i}$  ( $i < \lambda$ ), where  $M/i$  is the domain restriction of  $M$  to  $i$ , with the connecting morphisms  $a_{M/i, M/j} : A_{M/i} \rightarrow A_{M/j}$  for all  $i \leq j < \lambda$ . The proof that these chains are smooth is an easy transfinite induction.

It is also easy to see that all the above results hold:  $\hat{A} = \text{colim} D_A$  is an  $\mathcal{H}$ -injective weak reflection of  $A$ , and all connecting morphisms of  $D_A$  are members of  $\overline{\mathcal{H}}$ . Consequently, the proof of the following proposition is analogous to that of 5.6:

**5.9. Proposition** *Let  $\mathcal{H}$  be a set of  $k$ -petty morphisms with  $\overline{\mathcal{H}_k} \subseteq \mathcal{M}$ . Then for every object  $A \in \mathcal{A}_k$  a colimit  $\gamma_M : A_M \rightarrow \hat{A}$  of  $D_A$  yields a weak reflection of  $A$  in  $\text{Inj}\mathcal{H}$  via  $r_A = \gamma_\varepsilon : A \rightarrow \hat{A}$ .*

## 6. Completeness in strongly locally ranked categories

**6.1. Assumption** Throughout this section  $\mathcal{A}$  denotes a strongly locally ranked category. We first prove the completeness of the finitary logic. Recall that the finitary morphisms are those where the domain and codomain are of finite  $\mathcal{M}$ -rank. Let us remark that whenever the class  $\mathcal{M}$  is closed under pushout, then the method of proof of Theorem 3.6 applies again. However, this excludes examples such as **Haus** (where strong monomorphisms are not closed under pushout).

**6.2. Theorem** *The Finitary Injectivity Logic is complete in every strongly locally ranked category. That is, given a set  $\mathcal{H}$  of finitary morphisms, every finitary morphism  $h$  which is an injectivity consequence of  $\mathcal{H}$  is a formal consequence (in the deduction system of 2.4). Shortly:  $\mathcal{H} \models h$  implies  $\mathcal{H} \vdash h$ .*

**6.3. Remark** We do not need the full strength of weak local presentation for this result. We are going to prove the completeness under the following milder assumptions on  $\mathcal{A}$ :

- (i)  $\mathcal{A}$  is cocomplete and has a left-proper factorization system  $(\mathcal{E}, \mathcal{M})$ ;
- (ii)  $\mathcal{A}_\omega$  is a set of objects of finite  $\mathcal{M}$ -rank, closed under finite colimits and  $\mathcal{E}$ -quotients;
- (iii)  $\mathcal{M}$  is closed under filtered colimits in  $\mathcal{A}$  (see 4.5 (iv)).

The statement we prove is, then, concerned with petty morphisms (see 5.2). We show that for every set  $\mathcal{H}$  of petty morphisms we have

$$\mathcal{H} \models h \text{ implies } \mathcal{H} \vdash h \text{ (for all } h \text{ petty).}$$

The choice of  $\mathcal{A}_\omega$  as a set of representatives of all objects of finite  $\mathcal{M}$ -rank yields the statement of the theorem.

**Proof of 6.2 and 6.3** Let then  $\mathcal{H}$  be a set of petty morphisms, and let


$$\overline{\mathcal{H}}$$

denote the closure of  $\mathcal{H}$  as in 5.2.

(1) We first prove that the theorem holds whenever  $\overline{\mathcal{H}} \subseteq \mathcal{M}$ . Moreover, we will show that for every petty injectivity consequence  $\mathcal{H} \models h$  we have a formal proof of  $h$  from assumptions in  $\mathcal{H}$  such that the use of **PUSHOUT** is always restricted to pushing out along petty morphisms.



To prove this, consider, for the given petty injectivity consequence  $h : A \rightarrow B$  of  $\mathcal{H}$ , the weak reflection  $r_A : A \rightarrow \hat{A}$  in  $\text{Inj } \mathcal{H}$  of 5.6. The object  $\hat{A}$  is injective w.r.t.  $h$ , thus  $r_A$  factors through  $h$  via some  $f : B \rightarrow \hat{A}$ :



Since  $B \in \mathcal{A}_\omega$ , it has finite  $\mathcal{M}$ -rank, and 5.4 implies that  $\text{hom}(B, -)$  preserves the colimit  $\hat{A} = \text{colim} D_A$ . Then  $f$  factors through one of the colimit morphisms  $\gamma_N : A_N \rightarrow \hat{A}$ :

$$f = \gamma_N \cdot g \text{ for some } g : B \rightarrow A_N.$$

We know that  $r_A = \gamma_\varepsilon$  is the composite of the connecting morphism  $a_{\varepsilon, N} : A \rightarrow A_N$  of  $D_A$  and  $\gamma_N$ , therefore,

$$\gamma_N \cdot a_{\varepsilon, N} = r_A = \gamma_N \cdot g \cdot h.$$

That is, the colimit morphism  $\gamma_N$  merges the parallel pair  $a_{\varepsilon, N}, g \cdot h : A \rightarrow A_N$ . Now the domain  $A$  has finite  $\mathcal{M}$ -rank, thus  $\text{hom}(A, -)$  also preserves  $\hat{A} = \text{colim} D_A$ . Consequently, by (ii) in 3.1(b) the parallel pair is also merged by some connecting morphism  $a_{N, M} : A_N \rightarrow A_M$  of  $D_A$ :

$$a_{N, M} \cdot a_{\varepsilon, N} = a_{N, M} \cdot g \cdot h : A \rightarrow A_M.$$

The left-hand side is simply  $a_{\varepsilon, M}$ , and this is a morphism of  $\overline{\mathcal{H}}$ , see Lemma 5.4. Recall that the definition of  $\overline{\mathcal{H}}$  implies that every morphism in  $\overline{\mathcal{H}}$  can be proved from  $\mathcal{H}$  using Finitary Injectivity Logic in which PUSHOUT is only applied to pushing out along petty morphisms. Thus, we have a proof of the right-hand side  $a_{N, M} \cdot g \cdot h$ . The last step is deriving  $h$  from this by CANCELLATION.

(2) Assuming  $\mathcal{H} \subseteq \mathcal{E}$ , then we prove that  $\text{Inj } \mathcal{H}$  is a reflective subcategory of  $\mathcal{A}$ , and for every object  $A \in \mathcal{A}_\omega$  the reflection map  $r_A : A \rightarrow \hat{A}$  is a formal consequence of  $\mathcal{H}$  lying in  $\mathcal{E}$ :

$$\mathcal{H} \vdash r_A \text{ and } r_A \in \mathcal{E}.$$

In fact, from  $\mathcal{H} \subseteq \mathcal{E}$  it follows that  $\overline{\mathcal{H}} \subseteq \mathcal{E}$  (since  $\mathcal{E}$  is closed under composition and pushout). Since  $A$  has only finitely many  $\mathcal{E}$ -quotients, see 4.2, we can form a finite wide pushout,  $r_A : A \rightarrow \hat{A}$ , of all  $\mathcal{E}$ -quotients of  $A$  lying in  $\overline{\mathcal{H}}$ . Clearly,  $\mathcal{H} \vdash r_A$ , in fact,  $r_A \in \overline{\mathcal{H}}$ .

The object  $\hat{A}$  is injective w.r.t.  $\mathcal{H}$ : given  $h : P \rightarrow P'$  in  $\mathcal{H}$  and  $f : P \rightarrow \hat{A}$ , form a pushout  $h'$  of  $h$  along  $f$ . This is an  $\mathcal{E}$ -quotient in  $\overline{\mathcal{H}}$ , then the same is true for

$h' \cdot r_A$ . Consequently,  $r_A$  factors through  $h' \cdot r_A$ , and the factorization,  $i : B \rightarrow \hat{A}$ , is an epimorphism split by  $h'$ , thus,  $f = i \cdot g \cdot h$ :

$$\begin{array}{ccccc} & & P & \xrightarrow{h} & P' \\ & & \downarrow f & & \downarrow g \\ A & \xrightarrow{r_A} & \hat{A} & \xrightarrow{h'} & B \\ & & & \xleftarrow{i} & \end{array}$$

The morphism  $r_A$  is a weak reflection: given a morphism  $u$  from  $A$  to an object  $C$  of  $\text{Inj } \mathcal{H}$ , then  $u$  factors through  $r_A$  because  $C$  is injective w.r.t.  $\overline{\mathcal{H}}$  and  $r_A \in \overline{\mathcal{H}}$ .

(3) Let  $\mathcal{H}$  be arbitrary. We begin our proof by defining an increasing sequence of sets  $\mathcal{E}_i \subseteq \mathcal{E}$  of petty morphisms ( $i \in \text{Ord}$ ). For every member  $f : A \rightarrow B$  of  $\overline{\mathcal{H}}$  we denote by  $f_i$  a reflection of  $f$  in  $\text{Inj } \mathcal{E}_i$ :

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ r_A \downarrow & & \downarrow r_B \\ \hat{A} & \xrightarrow{f_i} & \hat{B} \end{array}$$

*First step:*  $\mathcal{E}_0 = \{\text{id}_A; A \in \mathcal{A}_\omega\}$ . Here  $\text{Inj } \mathcal{E}_0 = \mathcal{A}$ , thus  $f_0 = f$ .

*Isolated step:* For each  $f \in \overline{\mathcal{H}}$ , let  $f_i = f'_i \cdot f''_i$  be the  $(\mathcal{E}, \mathcal{M})$ -factorization of the reflection  $f_i$  of  $f$  in  $\text{Inj } \mathcal{E}_i$ , and put

$$\mathcal{E}_{i+1} = \mathcal{E}_i \cup \{f'_i; f \in \overline{\mathcal{H}}\}.$$

*Limit step:*  $\mathcal{E}_j = \cup_{i < j} \mathcal{E}_i$  for limit ordinals  $j$ .

We prove that for every ordinal  $i$  we have

$$\mathcal{H} \vdash f'_i \quad \text{for every } f \in \overline{\mathcal{H}} \tag{6.1}$$

and

$$\text{Inj } \mathcal{H} = \text{Inj } \mathcal{E}_i \cap \text{Inj } \{f_i\}_{f \in \overline{\mathcal{H}}}. \tag{6.2}$$

For  $i = 0$ , (6.1) and (6.2) are trivial (use CANCELLATION for (6.1) and IDENTITY for (6.2)). Given  $i > 0$ , assuming that  $\mathcal{H} \vdash f'_j$  for all  $j < i$ , with  $f : A \rightarrow B$  in  $\overline{\mathcal{H}}$ , that is,  $\mathcal{H} \vdash \mathcal{E}_i$ , we have, by (2), that


$$\mathcal{H} \vdash r_B \tag{6.3}$$

where  $r_B$  is the reflection of  $B$  in  $\text{Inj } \mathcal{E}_i$ . Thus,  $\mathcal{H} \vdash f_i \cdot r_A$ . Moreover,  $r_A$  is an epimorphism, therefore the following square

$$\begin{array}{ccccc} A & \xrightarrow{r_A} & \hat{A} & \xrightarrow{f_i} & \hat{B} \\ r_A \downarrow & & & & \downarrow \text{id} \\ \hat{A} & \xrightarrow{f_i} & & & \hat{B} \end{array}$$

is a pushout, which proves  $\mathcal{H} \vdash f_i$  (via PUSHOUT).  $\mathcal{H} \vdash f'_i$  then follows by CANCELLATION.

To prove (6.2), observe that (6.1) implies  $\text{Inj } \mathcal{H} \subseteq \text{Inj } \mathcal{E}_i$ , and our previous argument yields  $\text{Inj } \mathcal{H} \subseteq \text{Inj } \{f_i\}_{f \in \overline{\mathcal{H}}}$ . Thus, it remains to prove the reverse inclusion: every object  $X$  injective w.r.t.  $\mathcal{E}_i \cup \{f_i\}_{f \in \overline{\mathcal{H}}}$  is injective w.r.t.  $\mathcal{H}$ . In fact, given  $f : A \rightarrow B$  in  $\mathcal{H}$  and a morphism  $u : A \rightarrow X$ , then since  $X \in \text{Inj } \mathcal{E}_i$  we have a factorization  $u = v \cdot r_A$ , and then the injectivity of  $X$  w.r.t.  $f_i$  yields the desired factorization of  $u$  through  $f$ .



(4) Since  $\mathcal{A}_\omega$  is a small category, there exists an ordinal  $j$  with

$$\mathcal{E}_j = \mathcal{E}_{j+1}.$$

We want to apply (1) to the category

$$\mathcal{A}' = \text{Inj } \mathcal{E}_j,$$

and the set


$$\mathcal{A}'_\omega = \mathcal{A}_\omega \cap \text{obj } \mathcal{A}'.$$

Let us verify that  $\mathcal{A}'$  satisfies the assumptions (i) – (iii) of Remark 6.3 w.r.t.

$$\mathcal{E}' = \mathcal{E} \cap \text{mor } \mathcal{A}' \quad \text{and} \quad \mathcal{M}' = \mathcal{M} \cap \text{mor } \mathcal{A}'.$$

Ad(i):  $\mathcal{A}'$  is cocomplete because it is reflective in  $\mathcal{A}$ . Moreover, since the reflection maps lie in  $\mathcal{E}$ , it follows that  $(\mathcal{E}', \mathcal{M}')$  is a factorization system: in fact,  $\mathcal{A}'$  is closed under factorization in  $\mathcal{A}$ . Since  $\mathcal{E} \subseteq \text{Epi}(\mathcal{A})$ , we have  $\mathcal{E}' \subseteq \text{Epi}(\mathcal{A}')$ .

Ad(iii): It is sufficient to prove that  $\mathcal{A}'$  is closed under filtered colimits of  $\mathcal{M}'$ -morphisms in  $\mathcal{A}$ . In fact, let  $D$  be a filtered diagram in  $\mathcal{A}'$  with connecting morphisms in  $\mathcal{M}$ , and let  $c_t : C_t \rightarrow C$  ( $t \in T$ ) be a colimit of  $D$  in  $\mathcal{A}$ . Then  $C \in \mathcal{A}'$ , i.e.,  $C$  is injective w.r.t.  $f_j : \hat{A} \rightarrow E$  for every  $f \in \overline{\mathcal{H}}$ . This follows from  $\hat{A}$  having finite  $\mathcal{M}$ -rank (because  $A \in \mathcal{A}_\omega$  implies  $\hat{A} \in \mathcal{A}_\omega$  due to the fact that  $r_A : A \rightarrow \hat{A}$  is an  $\mathcal{E}$ -quotient): since  $\text{hom}(\hat{A}, -)$  preserves the colimit of  $D$ , every morphism  $u : \hat{A} \rightarrow C$  factors through some of the colimit morphisms:




Since  $C_t \in \mathcal{A}'$  is injective w.r.t.  $f_j$ , we have a factorization of  $v$  through  $f_j$ , and therefore,  $u$  also factors through  $f_j$ . This proves  $C \in \mathcal{A}'$ .

Ad(ii): Due to the above, every object of  $\mathcal{A}'$  having a finite  $\mathcal{M}$ -rank in  $\mathcal{A}$  has a finite  $\mathcal{M}'$ -rank in  $\mathcal{A}'$ . Also, a finite colimit of objects of  $\mathcal{A}'$  in  $\mathcal{A}'$  is a reflection (thus, an  $\mathcal{E}$ -quotient) of the corresponding finite colimit in  $\mathcal{A}$ . Thus, it lies in  $\mathcal{A}'_\omega$ .


Next we claim that the set  $\mathcal{H}' = \{f_j; f \in \overline{\mathcal{H}}\}$  fulfils

$$\mathcal{H}' \subseteq \mathcal{M}'$$

and  $\mathcal{H}'$  is closed under petty identities, composition, and pushouts along petty morphisms. In fact, in the above  $(\mathcal{E}, \mathcal{M})$ -factorization of  $f_j$ :



we know that  $f'_j$  lies in  $\mathcal{E}_{j+1} = \mathcal{E}_j$  and  $\hat{A}$  is injective w.r.t.  $\mathcal{E}_j$ , thus,  $f'_j$  is a split monomorphism (as well as an epimorphism, since  $\mathcal{E} \subseteq \text{Epi}(\mathcal{A})$ ). Thus,  $f'_j$  is an isomorphism, which implies  $f_j \in \mathcal{M}$ .  $\mathcal{H}'$  contains  $\text{id}_A$  for every  $A \in \mathcal{A}'_\omega$  because  $\overline{\mathcal{H}}$  contains it;  $\mathcal{H}'$  is closed under composition because  $\overline{\mathcal{H}}$  is (and  $f \mapsto f_j$  is the action of the reflector functor from  $\mathcal{A}$  to  $\text{Inj } \mathcal{E}_j$ ). Finally,  $\mathcal{H}'$  is closed under pushout along petty morphisms. In fact, to form a pushout of  $f_j : \hat{A} \rightarrow \hat{B}$  along  $u : \hat{A} \rightarrow C$  in  $\mathcal{A}' = \text{Inj } \mathcal{E}_j$ , we form a pushout,  $g$ , of  $f$  along  $u \cdot r_A$  in  $\mathcal{A}$ , and compose it with the reflection map  $r_D$  of the codomain  $D$ :



Since  $C$  lies in  $\mathcal{A}'$ , we can assume  $r_C = \text{id}_C$ , and the reflection  $\hat{g} = r_D \cdot g$  of  $g$  in  $\mathcal{A}'$  is then a pushout of  $f_j$  along  $u$ . Now  $f \in \overline{\mathcal{H}}$  implies  $g \in \overline{\mathcal{H}}$ , and we have  $\hat{g} = g_j \in \mathcal{H}'$ .


(5) We are ready to prove that if a petty morphism  $h : A \rightarrow B$  is an injectivity consequence of  $\mathcal{H}$ , then  $\mathcal{H} \vdash h$  in  $\mathcal{A}$ . We write  $\mathcal{H} \vdash_{\mathcal{A}} h$  for the latter since we work within two categories: when we apply (1) to  $\mathcal{A}'$  we use  $\vdash_{\mathcal{A}'}$  for formal consequence in  $\mathcal{A}'$ . Analogously with  $\models_{\mathcal{A}}$  and  $\models_{\mathcal{A}'}$ . Let  $\hat{h} : \hat{A} \rightarrow \hat{B}$  be a reflection of  $h$  in  $\mathcal{A}'$ , then

$$\mathcal{H}' \models_{\mathcal{A}'} \hat{h}$$

because every object  $C \in \mathcal{A}' = \text{Inj } \mathcal{E}_j$  which is injective w.r.t.  $\mathcal{H}' = \{f_j\}_{f \in \overline{\mathcal{H}}}$  is, due to (6.2), injective w.r.t.  $\mathcal{H}$  in  $\mathcal{A}$ . Then  $C$  is injective w.r.t.  $h$ , and from  $C \in \mathcal{A}'$  it follows easily that  $C$  is injective w.r.t.  $\hat{h}$ . Due to (4) we can apply (1). Therefore,

$$\mathcal{H}' \vdash_{\mathcal{A}'} \hat{h}.$$

We thus have a proof of  $\hat{h}$  from  $\mathcal{H}'$  in  $\mathcal{A}'$ . We modify it to obtain a proof of  $h$  from  $\mathcal{H}$  in  $\mathcal{A}$ . We have no problems with a line of the given proof that uses one of the assumptions  $f_j \in \mathcal{H}'$ : we know from (6.1) that  $\mathcal{H} \vdash_{\mathcal{A}} f_j$ , and we substitute that line with a formal proof of  $f_j$  in  $\mathcal{A}$ . No problem is, of course, caused by the lines using COMPOSITION or CANCELLATION. But we need to modify the lines using PUSHOUT because  $\mathcal{A}'$  is not closed under pushout in  $\mathcal{A}$ . However, a pushout,  $g''$ , of a morphism  $g$  along a petty morphism  $u$  in  $\mathcal{A}'$



is obtained from a pushout,  $g'$ , of  $g$  along  $u$  in  $\mathcal{A}$  by composing it with a reflection map  $r_{Q'}$  of the pushout codomain. Recall that  $P, P', Q \in \mathcal{A}_\omega$  imply  $Q' \in \mathcal{A}_\omega$ . Thus, we can replace the line  $g''$  of the given proof by using PUSHOUT in  $\mathcal{A}$  (deriving  $g'$ ), followed by a proof of  $r_{Q'}$  (recall from (6.3) that  $\mathcal{H} \vdash_{\mathcal{A}} r_{Q'}$ ) and an application of COMPOSITION. We thus proved that

$$\mathcal{H} \vdash_{\mathcal{A}} \hat{h}.$$

Since  $r_B \cdot h = \hat{h} \cdot r_A$  and  $\mathcal{H} \vdash_{\mathcal{A}} r_A$  (see (6.3)), we conclude  $\mathcal{H} \vdash_{\mathcal{A}} \hat{h} \cdot r_A$ ; by CANCELLATION then  $\mathcal{H} \vdash_{\mathcal{A}} h$ .

**6.4. Corollary** (Compactness Theorem) *Let  $\mathcal{H}$  be a set of finitary morphisms in a strongly locally ranked category. Every finitary morphism which is an injectivity consequence of  $\mathcal{H}$  is an injectivity consequence of a finite subset of  $\mathcal{H}$ .*

**6.5. Remark** We proceed by generalizing the completeness result from finitary to  $k$ -ary, where  $k$  is an arbitrary infinite cardinal. The  $k$ -ary logic, then, deals with  $k$ -ary morphisms (i.e., those having both domain and codomain of  $\mathcal{M}$ -rank  $k$ ) and the  $k$ -ary Injectivity Deduction System of 2.18.

**6.6. Theorem** *The  $k$ -ary Injectivity Logic is complete in every strongly locally ranked category. That is, given a set  $\mathcal{H}$  of  $k$ -ary morphisms, then every  $k$ -ary morphism which is an injectivity consequence of  $\mathcal{H}$  is a formal consequence (in the  $k$ -ary Injectivity Deduction System).*

**Proof** The whole proof is completely analogous to that of Theorem 6.2. As described in Remark 6.3 we work under the following milder assumptions on the category  $\mathcal{A}$ :

- (i)  $\mathcal{A}$  is cocomplete and has a left-proper factorization system  $(\mathcal{E}, \mathcal{M})$ ;
- (ii)  $\mathcal{A}_k$  is a set of objects of  $\mathcal{M}$ -rank  $k$ , closed under colimits of less than  $k$  morphisms and under  $\mathcal{E}$ -quotients;
- (iii)  $\mathcal{M}$  is closed under  $k$ -filtered colimits in  $\mathcal{A}$ .

The statement we prove is concerned with  $k$ -petty morphisms (see 5.7). We denote by  $\overline{\mathcal{H}}_k$  the closure of  $\mathcal{H}$  as in 5.7. We write  $\mathcal{H} \vdash h$  for the  $k$ -ary Injectivity Logic.

(1) The theorem holds whenever  $\overline{\mathcal{H}}_k \subseteq \mathcal{M}$ . The proof, based on the construction of a weak reflection  $\hat{A} = \text{colim} D_A$  of 5.8, is completely analogous to that of (1) in 6.2.

(2) Assuming  $\mathcal{H} \subseteq \mathcal{E}$ , then  $\text{Inj } \mathcal{H}$  is a reflective subcategory, and the reflection maps  $r_A$  fulfil  $\mathcal{H} \vdash r_A$  and  $r_A \in \mathcal{E}$ . This is analogous to the proof of (2) of 6.2.

(3) The definition of  $\mathcal{E}_i$  is precisely as in the proof of 6.2.

(4) For the first ordinal  $j$  with  $\mathcal{E}_j = \mathcal{E}_{j+1}$  the category  $\mathcal{A}' = \text{Inj } \mathcal{E}_j$  fulfils the assumptions (i)-(iii) above, and the set  $\mathcal{H}' = \{f_j; f \in \overline{\mathcal{H}}\}$  fulfils  $\mathcal{H}' = \overline{\mathcal{H}'} \subseteq \mathcal{M}$ .

(5) The theorem is then proved by applying (1) to  $\mathcal{A}'$  and  $\mathcal{H}'$ : we get  $\mathcal{H}' \vdash \hat{h}$  in  $\mathcal{A}'$  and we derive  $\mathcal{H} \vdash h$  in  $\mathcal{A}$  precisely as in the proof of 6.2.

**6.7. Corollary** *The Injectivity Logic is sound and complete. That is, given a set  $\mathcal{H}$  of morphisms of a strongly locally ranked category, then the consequences of  $\mathcal{H}$  are precisely the formal consequences of  $\mathcal{H}$  (in the Injectivity Deduction System). Shortly:*

$$\mathcal{H} \models h \quad \text{iff} \quad \mathcal{H} \vdash h \quad (\text{for all morphisms } h)$$

In fact, soundness was proved in Section 2. Completeness follows from Theorem 6.6: since  $\mathcal{H}$  is a set, and since every object of  $\mathcal{A}$  has an  $\mathcal{M}$ -rank, see 4.5(ii), there exists  $k$  such that all domains and codomains of morphisms of  $\mathcal{H} \cup \{h\}$  have  $\mathcal{M}$ -rank  $k$ .

## 7. Counterexamples

**7.1. Example** In “nice” categories which are not strongly locally ranked the completeness theorem can fail. Here we refer to  $\vdash$  of the Deduction System 2.13 (and the logic concerning arbitrary morphisms). We denote by

$$\mathbf{CPO}(\mathbf{1})$$

the category of unary algebras defined on  $CPO$ 's. Recall that a  $CPO$  is a poset with directed joins, and the corresponding category,  $\mathbf{CPO}$ , has as morphisms the *continuous functions* (i.e., those preserving directed joins). The category  $\mathbf{CPO}(\mathbf{1})$  has as objects the triples  $(A, \sqsubseteq, \alpha)$  where  $(A, \sqsubseteq)$  is a  $CPO$  and  $\alpha : A \rightarrow A$  is a unary operation. Morphisms are the continuous algebra homomorphisms.

First let us observe that the assumption of cocompleteness is fulfilled.

**Lemma  $\mathbf{CPO}(\mathbf{1})$  is cocomplete.**

**Proof** The category  $\mathbf{CPO}$  is easily seen to be cocomplete. The category  $\mathbf{CPO}(\mathbf{1})^*$  of partial unary algebras on  $CPO$ 's (defined as above except that we allow  $\alpha : A' \rightarrow$

$A$  for any  $A' \subseteq A$ ) is monotopological over **CPO**, see [3], since for every monosource  $f_i : (A, \sqsubseteq) \rightarrow (A_i, \sqsubseteq_i, \alpha_i)$  ( $i \in I$ ) we define a partial operation  $\alpha$  on  $A$  at an element  $x \in A$  iff  $\alpha_i$  is defined at  $f_i(x)$  for every  $i$ , and then

$$\alpha x = y \text{ iff } f_i(y) = \alpha_i(f_i(x)) \text{ for all } i \in I.$$

Consequently, **CPO(1)\*** is cocomplete by [3], 21.42 and 21.15. Further, **CPO(1)** is a full reflective subcategory of **CPO(1)\***: form a free unary algebra on the given partial unary algebra, ignoring the ordering, and then extend the ordering trivially (i.e., the new elements are pairwise incomparable, and incomparable with any of the original elements). Thus, **CPO(1)** is cocomplete.

We will find morphisms  $h_1, h_2$  and  $k$  of **CPO(1)** with

$$\{h_1, h_2\} \models k \text{ but } \{h_1, h_2\} \not\models k.$$

(i) We define a morphism  $h_1$  that expresses, by injectivity, the condition

$$(h1) \quad x \sqsubseteq \alpha x \text{ for all } x \in A.$$

Let  $=$  denote the discrete order on the set  $\mathbf{N}$  of natural numbers, and  $\sqsubseteq$  that order enlarged by  $0 \sqsubseteq 1$ . Let  $s : \mathbf{N} \rightarrow \mathbf{N}$  be the successor operation. Then

$$h_1 = \text{id} : (\mathbf{N}, =, s) \rightarrow (\mathbf{N}, \sqsubseteq, s)$$

is a morphism such that an algebra is injective w.r.t.  $h_1$  iff it fulfils (h1) above.

(ii) The condition

$$(h2) \quad A \neq \emptyset$$

is expressed by the injectivity w.r.t.

$$h_2 : \emptyset \rightarrow (\mathbf{N}, =, s)$$

where  $\emptyset$  is the empty (initial) algebra. The following morphism  $k$  expresses the existence of a fixed point of  $\alpha$ :

$$k : \emptyset \rightarrow 1$$

where  $1$  is a one-element (terminal) algebra.

**Proposition**  $\{h_1, h_2\} \models k$  but  $\{h_1, h_2\} \not\models k$ .

**Proof** To prove  $\{h_1, h_2\} \models k$ , let  $(A, \sqsubseteq, \alpha)$  be injective w.r.t.  $h_1$  and  $h_2$ , i.e., fulfill  $x \sqsubseteq \alpha(x)$  and be nonempty. Define a smooth (see 2.10) chain  $(a_i)_{i \in Ord}$  in  $(A, \sqsubseteq)$  by transfinite induction:  $a_0 \in A$  is any chosen element. Given  $a_i$  put  $a_{i+1} = \alpha(a_i)$ ; we know that  $a_i \sqsubseteq a_{i+1}$ . Limit steps are given by (directed) joins,  $a_j = \bigsqcup_{i < j} a_i$ . Since  $A$  is small, there exist  $i$  with  $a_i = a_{i+1}$ , that is,  $a_i$  is a fixed point of  $\alpha$ . Thus,  $A$  is injective w.r.t.  $k$ .

To prove  $\{h_1, h_2\} \not\models k$ , it is sufficient to find an extension  $\mathcal{K}$  of the category **CPO(1)** in which **CPO(1)** is closed under colimits (therefore  $\vdash$  has the same meaning in **CPO(1)** and in  $\mathcal{K}$ ) and in which there exists an object which is injective w.r.t.  $h_1$  and  $h_2$  but not w.r.t.  $k$ . Thus  $k$  cannot be proved in  $\mathcal{K}$  from  $h_1, h_2$ ; consequently it cannot be proved in **CPO(1)** either.

We define  $\mathcal{K}$  by adding a single new object  $K$  to **CPO(1)**. The only morphism with domain  $K$  is  $\text{id}_K$ . For every algebra  $(A, \sqsubseteq, \alpha)$  of **CPO(1)** we call a function

$f : A \rightarrow Ord$  a coloring of  $A$  provided that it is continuous and fulfils  $f(\alpha(x)) = f(x) + 1$  for all  $x \in A$ .

The hom-object of  $A$  and  $K$  in  $\mathcal{K}$  is defined to be the class of all colorings of  $A$ . The composition in  $\mathcal{K}$  is defined “naturally”: given a continuous homomorphism  $h : (A, \sqsubseteq, \alpha) \rightarrow (B, \leq, \beta)$ , then for every coloring  $f : B \rightarrow Ord$  of  $B$  we have a coloring  $f \cdot h : A \rightarrow Ord$  of  $A$ . The category  $\mathbf{CPO}(\mathbf{1})$  is a full subcategory of  $\mathcal{K}$  closed under (small) colimits. In fact, given a colimit cocone  $a_i : A_i \rightarrow A$  ( $i \in I$ ) in  $\mathbf{CPO}(\mathbf{1})$ , then for every compatible cocone of colorings  $f_i : A_i \rightarrow Ord$  ( $i \in I$ ) there exists an ordinal  $j$  such that all ordinals in  $\cup_{i \in I} f_i[A_i]$  are smaller than  $j$ . Let  $B = (j^+, \leq, \bar{s})$  be the object of  $\mathbf{CPO}(\mathbf{1})$  where  $\leq$  is the usual linear ordering of  $j^+$  (the poset of all ordinals smaller or equal to  $j$ ), and  $\bar{s}$  is the successor map except  $\bar{s}(j) = j$ . Then the codomain restriction  $f'_i$  of each  $f_i$  defines a continuous homomorphism  $f'_i : A_i \rightarrow B$ , and we obtain a compatible cocone  $(f'_i)_{i \in I}$  for our diagram. The unique continuous homomorphism  $g : A \rightarrow B$  with  $g \cdot a_i = f'_i$  yields, by composing it with the inclusion  $j^+ \hookrightarrow Ord$ , a coloring  $f : A \rightarrow Ord$  with  $f \cdot a_i = f_i$  ( $i \in I$ ).

It is obvious that  $K$  is injective w.r.t.  $h_1$ : every coloring of  $(\mathbf{N}, =, s)$  is also a coloring of  $(\mathbf{N}, \sqsubseteq, s)$ . And  $K$  is injective w.r.t.  $h_2$  (because the inclusion  $\mathbf{N} \hookrightarrow Ord$  is a coloring of  $(\mathbf{N}, =, s)$ ). But  $K$  is not injective w.r.t.  $k$ , since  $1$  has no coloring.

**7.2. Example** None of the deduction rules of the Finitary Injectivity Deduction System can be left out. For each of them we present an example of a finite complete lattice  $\mathcal{A}$  in which the reduced deduction system is not complete (for finitary morphisms).

(1) **IDENTITY** The deduction system **CANCELLATION**, **COMPOSITION** and **PUSHOUT** is not complete because nothing can be derived from the empty set of assumptions, although  $\emptyset \models \text{id}_A$ .


(2) **CANCELLATION** In the poset



the only object injective w.r.t.  $\{0 \rightarrow 2\}$  is  $2$ , thus, we see that  $\{0 \rightarrow 2\} \models 0 \rightarrow 1$ . However,  $0 \rightarrow 1$  cannot be derived from  $0 \rightarrow 2$  by means of **IDENTITY**, **COMPOSITION** and **PUSHOUT** because the set of all morphisms of  $\mathcal{A}$  except  $0 \rightarrow 1$  is closed under composition and pushout.

(3) **COMPOSITION** In  $\mathcal{A}$  above we clearly have  $\{0 \rightarrow 1, 1 \rightarrow 2\} \models 0 \rightarrow 2$ . However, the set of all morphisms except  $0 \rightarrow 2$  is closed under left cancellation and pushout.

(4) **PUSHOUT** In the poset



we have  $\{0 \rightarrow a\} \models b \rightarrow 1$ , but we cannot derive  $b \rightarrow 1$  from  $0 \rightarrow a$  using **IDENTITY**, **COMPOSITION** and **CANCELLATION** because the set of all morphisms except  $b \rightarrow 1$  is closed under composition and cancellation.



**7.3. Example** Here we demonstrate that in the Finitary Injectivity Logic we cannot restrict the statement of the completeness theorem from the given strongly locally ranked category  $\mathcal{A}$  to its full subcategory  $\mathcal{A}_\omega$  on all objects of finite rank: although the relation  $\vdash$  works entirely in  $\mathcal{A}_\omega$ , the relation  $\models$  does not.

More precisely, let  $\mathcal{H} \models_\omega h$  mean that every  $\mathcal{H}$ -injective object of finite  $\mathcal{M}$ -rank is also  $h$ -injective. And let  $\vdash_\omega$  be the formal consequence w.r.t. Deduction System 2.4. Then the implication

$$\mathcal{H} \models_\omega h \text{ implies } \mathcal{H} \vdash_\omega h$$

does NOT hold in general for sets of finitary morphisms.

Indeed, let  $\mathcal{A} = \mathcal{Gra}$  be the category of graphs, i.e., binary relational structures  $(A, R)$ ,  $R \subseteq A \times A$ , and the usual graph homomorphisms. Recall that  $\mathcal{Gra}$  is locally finitely presentable, and the finitely presentable objects are precisely the finite graphs. Let us call a graph a *clique* if  $R = A \times A - \Delta_A$ . Denote by  $C_n$  a clique of cardinality  $n$ , and let  $\mathbf{0}$  be the initial object (empty graph).

For the set

$$\mathcal{H} = \{\mathbf{0} \rightarrow C_n\}_{n \in \mathbb{N}}$$

we have the following property:

every finite  $\mathcal{H}$ -injective graph  $G$  has a loop (i.e., a morphism from  $\mathbf{1}$  to  $G$ ).

In fact, if  $G$  has cardinality less than  $n$  and is injective w.r.t.  $\mathbf{0} \rightarrow C_n$ , then we have a homomorphism  $f : C_n \rightarrow G$ . Since  $f$  cannot be one-to-one, there exist  $x \neq y$  in  $C_n$  with  $f(x) = f(y)$  – and the last element defines a loop of  $G$  because  $(x, y)$  is an edge of  $C_n$ . Hence

$$\mathcal{H} \models_\omega (\mathbf{0} \rightarrow \mathbf{1}).$$

However,  $\mathbf{0} \rightarrow \mathbf{1}$  cannot be proved in the Finitary Injectivity Logic. In fact, the graph

$$G = \coprod_{n \in \mathbb{N}} C_n$$

demonstrates that  $\mathcal{H} \not\models (\mathbf{0} \rightarrow \mathbf{1})$ .

## References

- [1] J. Adámek, M. Hébert and L. Sousa, A Logic of Orthogonality, *Archivum Mathematicum* 42 (2006), 309-334.
- [2] J. Adámek, H. Herrlich, J. Rosický and W. Tholen, On a generalized small-object argument for the injective subcategory problem, *Cah. Topol. Géom. Différ. Catég.* 43 (2002), 83–106.
- [3] J. Adámek, H. Herrlich and G. E. Strecker, *Abstract and Concrete Categories*, John Wiley and Sons, New York 1990. Freely available at [www.math.uni-bremen.de/~dmb/acc.pdf](http://www.math.uni-bremen.de/~dmb/acc.pdf)
- [4] J. Adámek and J. Rosický: *Locally presentable and accessible categories*, Cambridge University Press, 1994.
- [5] J. Adámek and J. Rosický, On injectivity in locally presentable categories, *Trans. Amer. Math. Soc.* 336 (1993), 785–804.
- [6] J. Adámek, M. Sobral and L. Sousa, Logic of implications, Preprints of the Department of Mathematics of the University of Coimbra 05-24 (2005).
- [7] H. Andr eka and I. N emeti, A general axiomatizability theorem formulated in terms of cone-injective subcategories. In: *Universal Algebra* (Proc. Conf. Esztergom Hungary 1977), L. M arki (ed.), Colloq. Math. Soc. J. Bolyai – North-Holland, Vol. 29, 1981, 13-35.
- [8] H. Andr eka and I. N emeti, Injectivity in categories to represent all first order formulas, *I. Demonstratio Mathematica* XII, 3 (1979), 717-732.
- [9] B. Banaschewski and H. Herrlich, Subcategories defined by implications, *Houston J. Math.* 2 (1976), 149-171.
- [10] G. Birkhoff, On the structure of abstract algebras, *Proc. Cambridge Phil. Soc.* 31 (1935), 433–454.
- [11] R. Diaconescu, Completeness of category-based equational deduction, *Mathem. Str. in Comput. Sci.* 5, 1 (1995), 9-40.
- [12] P. Gabriel and F. Ulmer, *Lokal Praesentierbare Kategorien*, Lect. Notes in Math. 221, Springer-Verlag, Berlin (1971).
- [13] M. H ebert, Purity and injectivity in accessible categories, *J. Pure Appl. Algebra* 129 (1998), 143-147.
- [14] M.D. Kan, On c.s.s. complexes, *Amer. J. Math.* 79 (1957), 449-476.
- [15] I. N emeti and I. Sain, Cone-implicational subcategories and some Birkhoff-type theorems. In: *Universal Algebra* (Proc. Conf. Esztergom Hungary 1977), L. M arki (ed.), Colloq. Math. Soc. J. Bolyai – North-Holland, Vol. 29, 1981, 535-578.
- [16] R. Quackenbush, Completeness theorems for universal algebra and implicational logics of algebras via congruences, *Proc. Amer. Math. Society* 103 (1988) 1015-1021.
- [17] D. Quillen, *Homotopical Algebra*, Lect. Notes in Math. 43, Springer-Verlag, Berlin 1967.

- [18] J. Rosický, J. Adámek and F. Borceux, More on injectivity in locally presentable categories, *Theory Appl. Categ.* 10 (2002), 148-238.
- [19] G. Roşu, Complete Categorical Equational Deduction, *Lect. Notes Comput. Sci.* 2142 (2001), 528–538.

This article may be accessed via WWW at <http://jhrrs.rmi.acnet.ge>

J. Adámek  
[adamek@iti.cs.tu-bs.de](mailto:adamek@iti.cs.tu-bs.de)

Department of Theoretical Computer Science,  
Technical University of Braunschweig,  
Postfach 3329, 38023 Braunschweig,  
Germany

M. Hébert  
[mhebert@aucegypt.edu](mailto:mhebert@aucegypt.edu)

Department of Mathematics,  
American University in Cairo,  
P.O. Box 2511, Cairo 11511,  
Egypt

L. Sousa  
[sousa@mat.estv.ipv.pt](mailto:sousa@mat.estv.ipv.pt)

Departamento de Matemática,  
Escola Superior de Tecnologia de Viseu, Campus Politécnico,  
3504-510 Viseu,  
Portugal