

COCOMPACT LATTICES

Grigore Călugăreanu

*Department of Mathematics, University “Babeş-Bolyai”, str.
Kogălniceanu 1, 3400 Cluj, Romania*

Received: August 1995

MSC 1991: 06 C 05, 06 C 20

Keywords: Cocompact, algebraic, inductive, reducible lattices, compact, essential, pseudocomplement, superfluous elements, socle and radical of a lattice.

Abstract: A lattice L is called cocompact if its dual L^0 is compact. If M is a R -module the lattice $S_R(M)$ of all the submodules of M is cocompact iff M is finitely cogenerated. Most of the properties of these modules are proved in the latticial general setting.

1. Introduction

A complete lattice L is called *cocompact* if each discover of 0 has a finite subdiscover, i.e. for every subset X of L such that $\bigwedge X = 0$ there is a finite subset F of X such that $\bigwedge F = 0$. Obviously, L is cocompact iff the dual L^0 is compact. An element $a \in L$ is called *cocompact* if the sublattice $a/0$ is cocompact.

The following characterization is well-known: a lattice L is artinian iff for each subset A of L there is a finite subset F of A such that $\bigwedge F = \bigwedge A$. Hence

Remark 1.1. *Every artinian lattice is cocompact.*

Remark 1.2. *If L is a cocompact lattice, for each $0 \neq a \in L$ the sublattice $a/0$ is also cocompact.*

Our main result is Th. 2.2: *Let L be an algebraic lattice. L is cocompact iff the socle $s(L)$ is compact and essential in L .*

In the sequel we will use only complete lattices L and the following definitions: a non-zero element e is called *essential* if for every

element $a \in L$, $a \wedge e = 0$ implies $a = 0$ and *superfluous* dually; the *socle* $s(L)$ of a lattice is defined as the join of all the atoms of L and, dually, the *radical* $r(L)$ as the meet of all the maximal elements (dual atoms) of L ; a lattice L is called *atomic* if for every $0 \neq a \in L$ the sublattice $a/0$ contains atoms, *inductive* if for each $a \in L$ and every chain $\{b_i\}_{i \in I}$, $\forall i \in I$, $a \wedge b_i = 0 \implies a \wedge (\bigvee_{i \in I} b_i) = 0$ and every sublattice (interval) of L has this property, *(R3)* if for every $a \neq 1$, a essential in L , $1/a$ contains atoms, *reducible* if the socle $s(L) = 1$, and *torsion* if for each $a \neq 1$, $1/a$ contains atoms (see [1], [2] and [3]). As in [1] we use the following definitions: we say that a set $\{a_i\}_{i \in I}$ of elements of a lattice is *independent* if $a_i \wedge (\bigvee_{j \neq i} a_j) = 0$ for all $i \in I$; in this case we denote the join $\bigvee_{i \in I} a_i$ by $\bigoplus_{i \in I} a_i$ and call it the *direct sum (join)*. For all the notions (such as: compact element, essential, pseudocomplement in a lattice and algebraic, artinian, pseudocomplemented or upper continuous lattice) and notation we refer to [4], [5] and [6].

2. Results

Lemma 2.1. *Let a be an essential element of a lattice L . If $a/0$ is cocompact then L is also cocompact.*

Proof. Let $\{a_i\}_{i \in I}$ be a family of non-zero elements of L such that $\bigwedge_{i \in I} a_i = 0$. The element a being essential in L , we have $a \wedge a_i \neq 0$ and

$0 = a \wedge \left(\bigwedge_{i \in I} a_i \right) = \bigwedge_{i \in I} (a \wedge a_i)$. Hence $\{a \wedge a_i\}_{i \in I}$ is a discover of 0 in $a/0$, and $a/0$ being cocompact there is a finite subset $F \subseteq I$ such that $0 = \bigwedge_{i \in F} (a \wedge a_i) = a \wedge \left(\bigwedge_{i \in F} a_i \right)$. Finally, a being essential, $\bigwedge_{i \in F} a_i = 0$ and L is cocompact. \diamond

Lemma 2.2. *In an algebraic, modular, reducible lattice the radical $r(L) = 0$.*

Proof. We verify that for each atom s , $s \wedge r(L) = 0$ (this suffices in a reducible lattice, which is also atomic). Reducible, inductive lattices being complemented (each algebraic lattice is upper continuous, each upper continuous lattice is inductive), let m be a complement of s . Using modularity, one easily proves that m is maximal in L . Hence $s \wedge m = 0$

implies $s \wedge r(L) = 0$. \diamond

Lemma 2.3. *In an algebraic cocompact lattice L the socle $s(L)$ is essential in L (more can be proved; see the last theorem).*

Proof. Let $a \in L$ be such that $s(L) \wedge a = 0$ or, equivalently, $s(a/0) = 0$. The sublattice $a/0$ being algebraic, the socle is also the join of all the essential elements (of $a/0$) and so, being cocompact $0 = \bigwedge_{i \in F} e_i$ for a

finite family of essential elements $\{e_i\}_{i \in F}$ of $a/0$. Hence 0 is essential in $a/0$ and so $a = 0$. \diamond

Remark 2.1. *In every atomic lattice the socle is essential. If the lattice L is inductive then the converse is also true. Indeed, if $a \neq 0$ then $0 \neq s(L) \wedge a \in s(L)/0$ an inductive and reducible lattice. Using Th. 9.2 from [1], each element of L is a direct sum of atoms. Hence $a/0$ contains atoms.*

So, *cocompact algebraic lattices are atomic*. Moreover, one can prove that *algebraic cocompact (R3) lattices are torsion lattices* (cf.[2]).

Proposition 2.1. *A lattice L is artinian iff for every $a \neq 1$ the sublattice $1/a$ is cocompact.*

Proof. Each sublattice of an artinian lattice is clearly artinian and so, by the Remark 1.1, is cocompact. Conversely, let $\dots \leq a_n \leq \dots \leq a_2 \leq a_1$ be an ascending chain of elements in L . If $a = \bigwedge_{n \in \mathbb{N}} a_n$ then

$\{a_n\}_{n \in \mathbb{N}}$ is surely a discover of a in $1/a$. The sublattice $1/a$ being cocompact there is a finite subset $F \subset \mathbb{N}$ such that $a = \bigwedge_{n \in F} a_n$. Hence

$a = a_m$ where $m = \min(F)$ and $a_{m+l} = a_m$ for each $l \in \mathbb{N}$, so the chain is finite and L is artinian. \diamond

Proposition 2.2. *If for an element a of an modular inductive lattice L the sublattices $a/0$ and $1/a$ are cocompact then the lattice L is cocompact.*

Proof. If $a = 0$ nothing remains to be proved. If $a \neq 0$ let $0 = \bigwedge_{i \in I} b_i$ a discover of 0 in L . Then $\bigwedge_{i \in I} (a \wedge b_i) = a \wedge (\bigwedge_{i \in I} b_i) = a \wedge$

$\bigwedge_{i \in I} b_i = 0$, is a discover of 0 in $a/0$. By cocompacity, there is a finite subset F of I such that $0 = \bigwedge_{i \in F} (a \wedge b_i) = a \wedge (\bigwedge_{i \in F} b_i)$. If $\bigwedge_{i \in F} b_i =$

0 (e.g. if a is essential in L) the proof is complete. If $\bigwedge_{i \in F} b_i \neq 0$ then let c be a pseudocomplement of a which contains $\bigwedge_{i \in F} b_i$. We have

$\bigwedge_{i \in F} b_i \in c/0 = c/(a \wedge c) \cong (a \vee c)/a \subseteq 1/a$ (the isomorphism is given

by modularity). The sublattice $1/a$ being cocompact, $(a \vee c)/a$ and hence $c/0$ are also cocompact. $0 = \bigwedge_{i \in I} (c \wedge b_i)$ being a discover of 0 in $c/0$ there is a finite subset G of I such that $0 = \bigwedge_{i \in G} (c \wedge b_i) = c \wedge \bigwedge_{i \in G} b_i$. Now, for $b = \bigwedge_{i \in F \cup G} b_i$ we have $b \leq \bigwedge_{i \in F} b_i \leq c$ and $c \wedge b \leq c \wedge \bigwedge_{i \in G} b_i = 0$ so that $b = 0$, and we have the required finite discover of 0. \diamond

This is a purely laticial proof which avoids the injective hull, a non-laticial notion (see [7]).

Consequence 2.1. *A direct sum of cocompact elements in an inductive modular lattice is cocompact.*

Proof. If $a/0, b/0$ are cocompact and $a \oplus b = 1$ (b is a complement of a) then by modularity $b/0 = b/(a \wedge b) \cong (a \vee b)/a = 1/a$ and we use the previous Prop. 2.2. \diamond

Proposition 2.3. *Let L be an algebraic cocompact lattice with the radical $r(L) = 0$. Then L is reducible and compact.*

Proof. From the third lemma we already know that L is atomic. The lattice L being algebraic the radical is also the union of all the superfluous elements. Hence the condition $r(L) = 0$ implies that the only superfluous element of L is 0. Equivalently, for each $0 \neq a \in L$ there is an $x \neq 1$ such that $a \vee x = 1$. In particular, each atom has a complement (maximal if L is also modular). Indeed, if s is an atom, as mentioned, there is an $m \neq 1$ such that $s \vee m = 1$. But $s \wedge m \in \{0, s\}$ and $s \wedge m = s$ implies $s \leq m$ or $m = 1$. Hence $s \wedge m = 0$ and s has a complement.

Now if the socle $s(L) \neq 1$ then let $x \neq 1$ be such that $s(L) \vee x = 1$ ($L \neq 0$ atomic implies $s(L) \neq 0$). One gets an atom which would not be contained in $s(L)$, contradiction. Hence L is reducible.

Finally, L being cocompact, the radical $r(L)$, which is the intersection of the maximal elements, and so is a discover of 0, must give a finite subdiscover of 0 by, say n maximal elements. The compactness of L follows now by induction on n . One verifies that each cover of 1 has a finite subcover. The dual analogon of this proof is detailed in the proof of the next theorem. \diamond

Theorem 2.1. *Let L be an algebraic, reducible and modular lattice. Then the following conditions are equivalent: (a) L is compact; (b) L is cocompact; (c) 1 is a finite direct sum of atoms.*

Proof. (a) \implies (c): L being reducible and inductive we have $1 = \bigoplus_{i \in I} s_i$, with s_i atoms (see [1]). But $\{s_i\}_{i \in I}$ is a cover for 1, compact element, so a finite subset $F \subseteq I$ exists such that $1 = \bigoplus_{i \in F} s_i$.

(c) \implies (b): If $\bigoplus_{i=1}^n s_i = 1$ we prove that every discover of $0 = \bigwedge_{i \in I} a_i$ has a finite subdiscover by induction on n . If $n = 1$ the assertion is obvious. We assume that the assertion is true for each lattice such that 1 is a direct sum of at most $n - 1$ atoms. First, observe that there is a $k \in I$ such that $a_k \wedge s_n = 0$. Indeed, otherwise $a_i \wedge s_n = s_n$ for every $i \in I$ or $s_n \leq \bigwedge_{i \in I} a_i$, contradiction. The element a_k is also a direct sum of at most $n - 1$ atoms (the modularity is needed for the use of the Jordan–Hölder theorem). By the induction hypothesis a finite subset of the family $\{a_i \wedge a_k\}_{i \in I}$ has the intersection 0. Hence L is cocompact.

(b) \implies (a) follows from Lemma 2.2 (which assures $r(L) = 0$) and Prop. 2.3. \diamond

Remark 2.2. *The implication (c) \implies (a) follows easily: in an upper continuous lattice every atom is compact and finite unions of compact elements are compact.*

Theorem 2.2. *Let L be an algebraic lattice. Then L is cocompact iff the socle $s(L)$ is compact and essential in L .*

Proof. If L is cocompact and $a \neq 0$ then clearly $a/0$ is also cocompact. Hence the sublattice $s(L)/0$ is cocompact and reducible. By Th. 2.1 $a/0$ is also compact, i.e. $s(L)$ is compact in L . The essentialness follows from Lemma 2.3. Conversely, if $s(L)$ is compact then $s(L)/0$ is reducible and compact and hence cocompact, again by the above theorem. The socle $s(L)$ being also essential in L , L is cocompact by Lemma 2.1. \diamond

References

- [1] BENABDALLAH, K. and PICHÉ, C.: Lattices related to torsion abelian groups, *Mitteilungen aus dem Math. Seminar Giessen*, Heft 197, Giessen 1990, 118 p.
- [2] CĂLUGĂREANU, G.: Torsion in lattices, *Mathematica* **25(48)** (1983), 127–129.
- [3] CĂLUGĂREANU, G.: Restricted socle conditions in lattices, *Mathematica* **28(51)** (1986), 27–29.

- [4] CRAWLEY, P. and DILWORTH, R.: *Algebraic Theory of Lattices*, Prentice-Hall, Englewood Cliffs, N.J., 1973.
- [5] GRÄTZER, G.: *General Lattice Theory*, Akademie-Verlag, Berlin, 1978.
- [6] STENSTRÖM, B.: *Rings of Quotients*, Springer Verlag, Berlin 1975.
- [7] WISBAUER, R.: *Foundation of Module and Ring Theory*, Gordon and Breach, Philadelphia, 1991.