

POLYHEDRA DUAL TO THE WEYL CHAMBER DECOMPOSITION: A PRÉCIS

KYOJI SAITO

ABSTRACT. Let $V_{\mathbb{R}}$ be a real vector space with an irreducible action of a finite reflection group W . We study the semi-algebraic geometry of the W -quotient affine variety $V//W$ with the discriminant divisor D_W in it and the τ -quotient affine variety $V//W//\tau$ with the bifurcation set B_W in it, where τ is the \mathbb{G}_a -action on $V//W$ obtained by the integration of the primitive vector field D on $V//W$ and B_W is the discriminant divisor of the induced projection $:D_W \rightarrow V//W//\tau$.

Our goal is the construction of a *one-parameter family of the semi-algebraic polyhedra* $K_W(\lambda)$ in $V_{\mathbb{R}}$ which are dual to the Weyl chamber decomposition of $V_{\mathbb{R}}$.

As an application, we obtain *geometric descriptions of generators for $\pi_1((V//W)_{\mathbb{C}}^{reg})$, satisfying the Artin braid relations.*

The key of the proof is a theorem on a linearization of the tube domain in $(V//W)_{\mathbb{R}}$ over the simplicial cone E_W in $T_{W,\mathbb{R}}$.

CONTENTS

1. Parallelotopes $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and polyhedra $K_W^{\varepsilon}(\lambda^{[\varepsilon]})$	9
1.1. Finite reflection group W	9
1.2. Simplicial cone decomposition of $V_{\mathbb{R}}$	9
1.3. Polyhedron dual to the simplicial cone decomposition	10
1.4. Invariants for W and the quotient variety S_W	11
1.5. Discriminant divisor and the central component $\mathcal{C}^{\{\varepsilon\}}$	12
1.6. Primitive vector field D and \mathbb{G}_a -action τ on S_W	13
1.7. The opposite components $\mathcal{C}_{\pm}^{[\varepsilon]}$ of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$	14
1.8. Semi-algebraic sets $\bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ in $S_{W,\mathbb{R}}^{[\varepsilon]}$ and $\bar{K}_W^{\varepsilon}(\lambda^{[\varepsilon]})$ in $V_{\mathbb{R}}^{\varepsilon}$	14
2. The central region $E_W^{\{\varepsilon\}}$ in $T_{W,\mathbb{R}}^{[\varepsilon]}$	16
2.1. τ -quotient space T_W and τ -quotient morphism π_{τ}	16
2.2. Bifurcation divisor $B_W = \bigcup_{p=2}^{\infty} B_{W,p}$	16
2.3. Real forms of the τ -action and the τ -quotient space	17
2.4. Subspace $S_{W(I_2(h))}$ of S_W	18

The present paper is an expanded version of an unpublished note of the author "The geometric generators for Artin groups of finite type, 1983", which was a sketch of the proof for the case $\varepsilon = +1$. Professor Egbert Brieskorn, at a conference Oberwolfach (1996), suggested the author to publish it. It is a great pleasure to the author to realize his suggestion in the occasion of the 40th anniversary of RIMS. The complete version including a proof of Theorem C shall appear in [S4].

2.5.	V.o. axis, v.o. line and the sign factor $\sigma(D, \{\Pi_1, \Pi_2\})$	19
2.6.	Algebroid functions φ_α and ϕ_α for $\alpha \in \Pi$	21
3.	Linearization map c_W	24
3.1.	Linear model spaces \widehat{V}_Π and V_Π	25
3.2.	$\Gamma(W)$ -cone $E_{\Gamma(W)}$ in V_Π	26
3.3.	Covering spaces $T_{W, \text{odd}, \mathbb{C}}$ and $S_{W, \text{odd}, \mathbb{C}}$	26
3.4.	Linearization morphism c_W on $S_{W, \text{odd}, \mathbb{C}}$	27
3.5.	Theorem C	28
3.6.	Proof of Theorem A	31
3.7.	Proof of Theorem B	32
3.8.	Examples of type A_3	36
4.	Fundamental group of $S_{W, \mathbb{C}} \setminus D_{W, \mathbb{C}}$	37
4.1.	1-skeleton of the polyhedron $J_W^{\{\varepsilon\}}(\lambda)$	37
4.2.	Proof of Theorem 4.1	38
4.3.	Zariski-van Kampen generator system	39
4.4.	Proof of Theorem 4.2	39
4.5.	Comparison of generator systems for $\varepsilon \in \{\pm 1\}$	40
	Appendix. Dihedral Group of Type $I_2(h)$	42
	References	45

Table of Figures.

Fig. 1.	Four base point loci in $T_{W, \text{odd}, \mathbb{C}}$	27
Fig. 2.	The linearization maps of type A_3	36
Fig. 3.	The generator on 1-skeleton of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$	37
Fig. 4.	The 2-facet $[p_\alpha, p_\beta, *]$	38
Fig. 5.	Pencil close to $\overline{\alpha\beta}$ -edge	38
Fig. 6.	The Zariski-van Kampen generators on a τ -orbit	39
Fig. 7.	The complexification of the vertex orbit axis AO	41
Fig. 8.	Polyhedra $J_{A_2}^{\{+1\}}(\lambda^{[+1]})$ and $K_{A_2}^+(\lambda^{[+1]})$ for $\lambda^{[+1]} = 1$	43
Fig. 9.	Polyhedra $J_{A_2}^{\{-1\}}(\lambda^{[-1]})$ and $K_{A_2}^-(\lambda^{[-1]})$ for $\lambda^{[-1]} = 1$	43
Fig. 10.	Positions of $S_{A_2, \mathbb{R}}^{[+1]}$ and $S_{A_2, \mathbb{R}}^{[-1]}$ inside $S_{A_2, \mathbb{C}} \cap \{\text{Im}(R) = 0\}$	43
Fig. 11.	Polyhedra $J_{B_2}^{\{\pm 1\}}(\lambda^{[\pm 1]})$ and $K_{B_2}^{\pm 1}(\lambda^{[\pm 1]})$ for $\lambda^{[\pm 1]} = 1$	44
Fig. 12.	Positions of $S_{B_2, \mathbb{R}}^{[\pm 1]}$ and $S_{B_2, \mathbb{R}}^{[\beta]}$ inside $S_{B_2, \mathbb{C}} \cap \{\text{Im}(R) = 0\}$	44

Introduction

Let $V_{\mathbb{R}}$ be a finite-dimensional real vector space and W a finite group acting irreducibly on $V_{\mathbb{R}}$ generated by reflections. We denote by V the associated scheme over \mathbb{R} and by $S_W := V//W$ the quotient scheme¹. Let $D_W \subset S_W$ be the discriminant divisor defined by the zero locus of $\Delta :=$ the square of the fundamental anti-invariant of W . The open regular part $(V//W)^{reg}$, defined as the complement $S_W \setminus D_W$, is a simple geometric object where several different areas of mathematics (e.g., Lie group theory, complex and differential geometries, ..., etc.) intersect.

We recall two basic results on the topology of the complexification $(V//W)_{\mathbb{C}}^{reg} := S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ of the regular orbit space:

- a) *the fundamental group of $(V//W)_{\mathbb{C}}^{reg}$ is an Artin group (generalized braid group)* (Brieskorn [Br1],[Br2] and [BS]), and
- b) *the universal covering of $(V//W)_{\mathbb{C}}^{reg}$ is contractible* (Deligne [D1]).

Interestingly, for the both results, the polyhedron K_W which is dual to the simplicial cone decomposition of $V_{\mathbb{R}}$ plays an essential role. Namely, a) the 1-skeleton and the 2-skeleton of K_W determine the generators and relations for the fundamental group of $(V//W)_{\mathbb{C}}^{reg}$, and b) the contractibility of K_W is a key step in the proof ([D1]) of the contractibility of the nerve of a simple covering of the universal covering of $(V//W)_{\mathbb{C}}^{reg}$. We remark further that c) *the dual polyhedron K_W also describes the Stiefel-Whitney class of a related vector bundle* ([Hu],[M] and [N]).

A goal of the present paper is to reconstruct the dual polyhedron K_W from a completely different viewpoint. The quotient variety $S_W := V//W$ carries a differential geometric structure, called the flat structure (Saito [S1,3]). Then we shall make use of a part of the real flat structure to construct the polyhedron as follows.

A principal ingredient of the flat structure is the vector field D on S_W of the lowest degree, which is unique up to a constant factor, called the *primitive vector field* (1.6.1). The integration $\exp(\lambda D)$ of D induces a \mathbb{G}_a -action τ on S_W (1.6.2), transversal to the discriminant divisor D_W (see [S2,3] for the role of D in the theory of primitive forms).

For $\varepsilon \in \{\pm 1\}$, consider the real form $S_{W,\mathbb{R}}^{[\varepsilon]}$ of $S_{W,\mathbb{C}}$ (the “quotient” of the real form $V_{\mathbb{R}}^{\varepsilon} := \sqrt{\varepsilon} \otimes V_{\mathbb{R}}$ of $V_{\mathbb{C}} := \mathbb{C} \otimes V$, see (1.4.8)). The \mathbb{G}_a -action τ induces the one-parameter group action $\tau^{[\varepsilon]} : \mathbb{R} \times S_{W,\mathbb{R}}^{[\varepsilon]} \rightarrow S_{W,\mathbb{R}}^{[\varepsilon]}$ (see

¹We mean by “ $V//W$ ” the categorical quotient scheme (1.4.5) of V by the W -action. Even though W is a finite group, it is convenient to use scheme-theoretic concepts and notation, since we study mainly over the real number field \mathbb{R} . The set theoretic quotient space $V_{\mathbb{R}}/W$ is not sufficient to describe structures we study. The \mathbb{R} or \mathbb{C} -rational point set of a scheme is indicated by the subscript \mathbb{R} or \mathbb{C} .

(1.6.4)). For each fixed $\lambda^{[\varepsilon]} \in \mathbb{R}_{>0}$, consider three real hypersurfaces in $S_{W,\mathbb{R}}^{[\varepsilon]}$: a) the real discriminant locus : $D_{W,\mathbb{R}}^{[\varepsilon]}$ and b) $^\pm$ the positive and negative translations of the real discriminant locus: $\tau^{[\varepsilon]}(\lambda^{[\varepsilon]})(D_{W,\mathbb{R}}^{[\varepsilon]})$ and $\tau^{[\varepsilon]}(-\lambda^{[\varepsilon]})(D_{W,\mathbb{R}}^{[\varepsilon]})$. Then, for $\varepsilon \in \{\pm 1\}$ and each $\lambda^{[\varepsilon]} \in \mathbb{R}_{>0}$, one has:

Theorem A (§1.8). *There exists an open semi-algebraic parallelootope $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ in $S_{W,\mathbb{R}}^{[\varepsilon]}$, which is surrounded² by the hypersurfaces a) and b) $^\pm$. It is adjacent to the origin $o \in S_{W,\mathbb{R}}^{[\varepsilon]}$, and the faces adjacent to the origin are indexed by the set Π of simple generators of W .*

Theorem B (§1.8). *The inverse image $K_W^\varepsilon(\lambda^{[\varepsilon]})$ in $V_{\mathbb{R}}^\varepsilon$ of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ in $S_{W,\mathbb{R}}^{[\varepsilon]}$ is an open polyhedron which is dual to the simplicial cone decomposition of $V_{\mathbb{R}}^\varepsilon$ by the Weyl chambers.*

See Appendix Fig. 8–12 for illustrated examples of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and $K_W^\varepsilon(\lambda^{[\varepsilon]})$ of type A_2 and B_2 .

It was asked by Brieskorn, Deligne, and others (including the author) to find some descriptions of the generator system of $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, *)$ as an Artin group in terms of the geometry of S_W . Let us give two answers to this question as an application of Theorems A and B (see §4 for details and proofs).

1. Let $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ be the vertex of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ antipodal to the origin o . Due to Theorem A, the edges of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ adjacent to $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ are indexed by the set Π in such a manner that the α th edge for $\alpha \in \Pi$ intersects the α th face of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ transversally at a point, say p_α , in $D_{W,\mathbb{R}}^{[\varepsilon]}$ (see Fig. 4). Inside a complexification of the α th edge (an open complex curve in $S_{W,\mathbb{C}}$ containing the α th edge), take a path, say γ_α , based at $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and turning counter-clockwise once around the discriminant divisor $D_{W,\mathbb{C}}$ at p_α (Fig. 3). Here, the class of γ_α in $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ is uniquely determined by the index $\alpha \in \Pi$.

Corollary 1 (§4.1 and §4.2). *The 1-homotopy classes of γ_α for $\alpha \in \Pi$ give a system of generators for $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]}))$, which satisfy the Artin braid relations as the system of fundamental relations.*

2. Next, we choose an arbitrary point $* \in J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and consider the orbit $\tau^{[\varepsilon]}(\mathbb{R}) \cdot *$ which is a real line in $S_{W,\mathbb{R}}^{[\varepsilon]}$. If $*$ is generic, the real line intersects l distinct points of the real discriminant locus $D_{W,\mathbb{R}}^{[\varepsilon]}$ ³ (Fig.

²By the word “surrounded”, we mean that $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ is a connected component of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus (D_{W,\mathbb{R}}^{[\varepsilon]} \cup \tau^{[\varepsilon]}(\lambda^{[\varepsilon]})(D_{W,\mathbb{R}}^{[\varepsilon]}) \cup \tau^{[\varepsilon]}(-\lambda^{[\varepsilon]})(D_{W,\mathbb{R}}^{[\varepsilon]}))$.

³This fact is a non-trivial consequence of Theorem C stated below.

5). One chooses paths inside the complex line $\tau^{[\varepsilon]}(\mathbb{C}) \cdot *$ as in Fig. 6, whose homotopy classes are called the Zariski-van Kampen generators.

Corollary 2 (§4.3 and §4.4). *The system of the Zariski-van Kampen generators is homotopic to the generator system in Corollary 1.*

Theorems A and B and their corollaries are direct applications of another basic Theorem C on the real bifurcation set which we explain below.

The quotient space $T_W := S_W // \tau$ by the τ -action is a smooth $(l-1)$ -dimensional affine variety, and the quotient map $\pi_\tau : S_W \rightarrow T_W$ is a linear projection in the direction of the primitive vector field. The restriction $\pi_\tau|_{D_W}$ of π_τ to the discriminant divisor is a finite covering over T_W . The ramification divisor B_W , i.e., the discriminant divisor of $\pi_\tau|_{D_W}$, is called the *bifurcation set*. Decompose it as $B_W = \cup_{p=2}^\infty B_{W,p}$ according to the ramification index p , where $B_{W,1}$ does not appear due to the transversality property of the primitive vector field D to D_W . We split the bifurcation set B_W into the ordinary part $B_{W,2}$ and the higher part $B_{W,\geq 3}$ (called the stratum of Maxwell's convention and the caustics, respectively, in [T2]).

For each $\varepsilon \in \{\pm 1\}$, we introduce some closed subset O^ε in $T_{W,\mathbb{R}}^{[\varepsilon]} \setminus B_{W,\geq 3,\mathbb{R}}^{[\varepsilon]}$ (resp. AO^ε in $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$), which are defined by the help of regular eigenvectors of the Coxeter element of W (see 2.5). They shall play two basic roles: i) to single out particular connected components of $T_{W,\mathbb{R}}^{[\varepsilon]} \setminus B_{W,\geq 3,\mathbb{R}}^{[\varepsilon]}$ (resp. $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$) containing them, and ii) to be chosen as a base point for the fundamental group of the complexification $T_{W,\mathbb{C}} \setminus B_{W,\geq 3,\mathbb{C}}$ (resp. $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$). On the other hand, they are related with the vertex of the polyhedra $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ as: $AO^\varepsilon = \{ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]}) \mid \lambda^{[\varepsilon]} \in \mathbb{R}_{>0}\}$ and $O^\varepsilon = \pi_\tau(AO^\varepsilon)$. We, therefore, call AO^ε the *half vertex orbit axis* and O^ε the *half vertex orbit line* (here “half” indicates that they are isomorphic to the half line $\mathbb{R}_{>0}$).

The connected component $\mathcal{C}^{\{\varepsilon\}}$ of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$ containing AO^ε is nothing but the image of a Weyl chamber in $V_{\mathbb{R}}^\varepsilon$, called the central component. The connected component $E_W^{\{\varepsilon\}}$ of $T_{W,\mathbb{R}}^{[\varepsilon]} \setminus B_{W,\geq 3,\mathbb{R}}^{[\varepsilon]}$ containing O^ε , called the *central region*, is a key object in the present paper. The fact which makes the situation non trivial is that although the region $E_W^{\{\varepsilon\}}$ contains the image $\pi_\tau(\mathcal{C}^{\{\varepsilon\}})$, the gap $E_W^{\{\varepsilon\}} \setminus \pi_\tau(\mathcal{C}^{\{\varepsilon\}})$ is “growing exponentially” as the rank l grows.

Theorem C of the present paper concerns the central region $E_W^{\{\varepsilon\}}$ and its inverse image $\pi_\tau^{-1}(E_W^{\{\varepsilon\}})$ (called the *tube domain*) in $S_{W,\mathbb{R}}^{[\varepsilon]}$.

Let $\widehat{V}_\Pi := \bigoplus_{\alpha \in \Pi} \mathbb{R}v_\alpha$ be the vector space with basis v_α attached to the set Π of simple generators for W , and let $V_\Pi := \widehat{V}_\Pi / \mathbb{R}v_\Pi$ be the quotient space for $v_\Pi := \sum_{\alpha \in \Pi} v_\alpha$, and let $\pi_\Pi : \widehat{V}_\Pi \rightarrow V_\Pi$ be the projection.

Theorem C (§3.5). *There exist i) an open simplicial cone $E_{\Gamma(W)} \subset V_\Pi$ depending only on the Coxeter diagram $\Gamma(W)$ in such a manner that its faces are indexed by the edges of $\Gamma(W)$, and ii) real algebroid maps c_W and b_W with the commutative diagram:*

$$\begin{array}{ccc} (\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}}) & \xrightarrow{c_W} & (\pi_\Pi)^{-1}(E_{\Gamma(W)}) \\ \pi_\tau^{[\varepsilon]} \downarrow & & \pi_\Pi \downarrow \\ E_W^{\{\varepsilon\}} & \xrightarrow{b_W} & E_{\Gamma(W)} \end{array}$$

where we mean by \simeq a semi-algebraic isomorphism. The map c_W induces a bijection

$$D_{W, \mathbb{R}}^{[\varepsilon]} \cap (\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}}) \simeq (\cup_{\alpha \in \Pi} H_\alpha) \cap (\pi_\Pi)^{-1}(E_{\Gamma(W)})$$

where H_α is the coordinate hyperplane in \widehat{V}_Π .

Corollary. *The real discriminant locus $D_{W, \mathbb{R}}^{[\varepsilon]}$ cut by the tube domain $(\pi_\Pi)^{-1}(E_{\Gamma(W)})$ decomposes into the union of hyperplanes H_α indexed by $\alpha \in \Pi$.*

The linearization maps c_W and b_W of type A_3 are illustrated in Fig. 2. Precise statements of Theorems A, B and C are given in §1.8 and §3.5.

Theorems A and B and their corollaries are proved in §3 as the direct consequences of Theorem C. However, Theorem C is not proved in the present article, since Theorem C is a part of consequences of a general study of the linearization maps c_W and b_W , whose comprehensive treatment shall appear in [S4].

Before we go further, we explain a motivation of the present paper. The quotient variety S_W appears as the base space of the universal unfolding $X_W \rightarrow S_W$ of a simple singularity [Br3]. On the total space X_W there is a special de Rham cohomology class relative to S_W , called the primitive form $\zeta_W^{(0)}$ [S2]. The period integral $\int \zeta_W^{(0)}$ over cycles in the fibers of the unfolding gives a multivalued map, called the period map, defined on $S_W \setminus D_W$ to the period domain. For the study of the period map, we need to understand the homotopy groups of the space $S_W \setminus D_W$. This gives one motivation.

The primitive form induces the flat structure on S_W , where $\zeta_W^{(0)}$ is identified with the primitive vector field D on S_W ([S2]). In the present paper, we employ not only D but the basic framework of the theory of primitive forms such as the τ -orbit space T_W with its bifurcation

divisor B_W , the characteristic variety C_W and the finite morphism $q_W : C_W \rightarrow T_W$. Therefore, it does not seem an accident that the polyhedron K_W is reconstructed through the action τ , the integral of the primitive vector field. However, we still need to clarify the relation of the period map for $\zeta_W^{(0)}$ with the polyhedron $K_W(\lambda)$. Some natural questions are the followings. *Can one reconstruct Deligne's proof [D1] in terms of the semi-algebraic geometry of the spaces V and $V//W$ as in the present work? Is $T_{W,\mathbb{C}} \setminus B_{W,\geq 3,\mathbb{C}}$ an Eilenberg-MacLane space? Determine the fundamental relations for its fundamental group with respect to the natural generators indexed by the edges of $\Gamma(W)$.*

There are many precedent works on the semi-algebraic geometry of the space S_W with the discriminant divisor D_W in it, among others, by Hilbert [H], Thom [T1,2], Arnold [Ar1,2], Looijenga [Lo1,2], Springer [Sp1,2] and Tits [?]. In particular, Thom's idea on the universal unfolding ([T2]) influenced either directly or indirectly on the idea of the primitive form and the primitive vector field. We also note an article on the semi-algebraic geometry of the orbit spaces of compact Lie groups by Procesi-Schwarz [P-S], though we do not know yet its direct relation with the present paper.

Let us explain the construction of the present paper.

The first half of §1 is an elementary preparation on the quotient variety $S_W := V//W$ by the finite reflection group W . Then, we introduce the τ -action on S_W and on its real forms. After these preparations, we formulate Theorems A and B in §1.8.

§2 studies the τ -quotient variety T_W with its bifurcation set B_W . After introducing the base point loci O^ε in $T_{W,\mathbb{C}} \setminus B_{W,\geq 3,\mathbb{C}}$, we introduce the central regions $E_W^{\{\varepsilon\}}$ in §2.5, and algebroid functions $\varphi_{\alpha,\varepsilon}$ in §2.6. This section is an extract from §2-§9 of the forthcoming paper [S4]. Leaving a general treatment to [S4], we restrict our attention only to the real structures $[\varepsilon]$. We also omitted the study of the characteristic variety C_W (which plays an important role to understand the discriminant divisor D_W).

In §3, we study the linearization map c_W . The target spaces \widehat{V}_Π , V_Π and the simplicial cone $E_{\Gamma(W)}$ are introduced in §3.1 and §3.2. The map c_W is introduced as an algebroid map in §3.4. Using them, Theorem C is formulated in §3.5. As its application, Theorems A and B are proved in §3.6 and §3.7. The proof of Theorem C is not given in the present paper but it is given in [S4], where we formulate c_W as an algebraic correspondence, which is more appropriate for our purpose.

§4 studies the generator systems of the fundamental group of the space $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$. A pair of generator systems depending on $\varepsilon \in \{\pm 1\}$ is constructed by use of the polyhedra $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ in §4.1 and is identified with Brieskorn's generator system in §4.2. A pair of the Zariski-van Kampen generator systems depending on ε by the use of τ -pencil is described in §4.3. It is identified in §4.4 with the one in §4.1. The relationship between the generator systems for $\varepsilon = +1$ and for $\varepsilon = -1$ is given in §4.5.

Appendix studies the rank two case in detail. The polyhedra $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and $K_W^\varepsilon(\lambda^{[\varepsilon]})$ of types A_2 and B_2 are illustrated in Fig. 8, 9 and 11.

Concluding Remarks: The study of the polyhedra $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$, $K_W^\varepsilon(\lambda^{[\varepsilon]})$ and the real region $E_W^{\{\varepsilon\}}$ has just started. The proofs are rather involved. On the other hand, we have observed a new aspect of the geometry of V , $V//W$ and $V//W//\tau$: the interaction between the semi-algebraic geometry of their real forms and the topology of their complexification, where the flat structure combines them. We may briefly summarize the present work as *a combinatorial aspect of the flat structure on the quotient variety by a finite reflection group*. These new features of the geometry seem to the author quite attractive and worthwhile to be studied further. Perhaps (and hopefully), the study in the present paper is the first fortunate model case ⁴ of a certain new mathematical research subject.

The author would like to express his hearty gratitude to Professors Masaki Kashiwara and Takahiro Kawai for their supports and helps during the preparation of the present paper, to Professor Hiroaki Terao for careful reading of the manuscripts and many useful pieces of advice, and to Mrs. Kumiko Matsumura for her beautiful drawing of figures.

The author would like to express his deep sorrow to the early death of the late Professor Nobuo Sasakura (March 5, 1941 – June 16, 1997), who constantly showed interests in the present work when it was in a preparatory form.

⁴One next model may be the case of elliptic root systems, which admit again the flat structure. Since the complement of the complex discriminant divisor may have 2-homotopy classes, we need to study the non-simply connected polyhedra.

1. PARALLELOTOPE $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ AND POLYHEDRA $K_W^\varepsilon(\lambda^{[\varepsilon]})$

We construct our main objects $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and $K_W^\varepsilon(\lambda^{[\varepsilon]})$ of the present paper, and give precise statements of Theorems A and B announced in the introduction.

In 1.1–1.5, we recall basic results on a finite reflection group W and its invariants from [B,Ch.4,5]. In 1.6 and 1.7, we introduce the new concept: the τ -action on the W -quotient varieties $S_{W,\mathbb{R}}^{[\pm 1]}$. By the use of the τ -action, Theorems A and B in §1.8 describe the polyhedra $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ and $K_W^\varepsilon(\lambda^{[\varepsilon]})$.

1.1. Finite reflection group W .

Let $V_{\mathbb{R}}$ be an \mathbb{R} -vector space of rank l equipped with the classical topology. An element $\alpha \in \text{GL}(V_{\mathbb{R}})$ is a *reflection* if there exist $e_\alpha \in V_{\mathbb{R}}$ and $f_\alpha \in V_{\mathbb{R}}^* := \text{Hom}_{\mathbb{R}}(V_{\mathbb{R}}, \mathbb{R})$ with $\langle f_\alpha, e_\alpha \rangle = 2$ such that $\alpha(x) = x - f_\alpha(x)e_\alpha$ for $x \in V_{\mathbb{R}}$. Two vectors e_α and f_α are not unique but $e_\alpha \otimes f_\alpha$ is uniquely determined by α . If I is an α -invariant symmetric bilinear form on $V_{\mathbb{R}}$ such that $I(e_\alpha, e_\alpha) \neq 0$, then $f_\alpha(x) = I(e_\alpha^\vee, x)$ for $e_\alpha^\vee := 2e_\alpha/I(e_\alpha, e_\alpha)$. The kernel $H_\alpha := \ker(f_\alpha) = \ker(1 - \alpha)$ is called the *reflection hyperplane* of α .

Let W be a finite group generated by reflections on $V_{\mathbb{R}}$ and I a W -invariant positive-definite symmetric bilinear form on V . Assume that W acts irreducibly on $V_{\mathbb{R}}$. Then, I is unique up to a positive constant. Put

$$(1.1.1) \quad R(W) := \{\alpha \in W \mid \alpha \text{ is a reflection}\}.$$

We recall some basic facts on W in [B].

1. A connected component of $V_{\mathbb{R}} \setminus \bigcup_{\alpha \in R(W)} H_\alpha$, called a *Weyl chamber*, is a simplicial cone. The group W acts simply transitively on the set of chambers.

2. Put $\Pi(C) := \{\alpha \in R(W) \mid H_\alpha \text{ is a wall of } C\}$ for a chamber C . Then $(W, \Pi(C))$ is a Coxeter system with respect to the Coxeter matrix $M_W := (m_{\alpha\beta})_{\alpha, \beta \in \Pi(C)}$ with $m_{\alpha\beta} := \text{ord}(\alpha\beta)$ (see [B, Ch.IV, §1 n°1.3.]).

3. The closure \bar{C} of a chamber C is a fundamental domain of the action of W on $V_{\mathbb{R}}$, that is, $\bar{C} \rightarrow V_{\mathbb{R}}/W$ is a homeomorphism.

4. The vectors $\{e_\alpha \mid \alpha \in \Pi(C)\}$ form a basis of $V_{\mathbb{R}}$. Choose e_α so that $C = \{x \in V_{\mathbb{R}} \mid \langle f_\alpha, x \rangle > 0 \text{ for } \alpha \in \Pi(C)\}$. Then i) $I(e_\alpha, e_\beta) \leq 0$ for $\alpha \neq \beta \in \Pi(C)$, and ii) the coefficients of the expression $e_\gamma = \sum_{\alpha \in \Pi(C)} c_\alpha e_\alpha$ for any $\gamma \in R(W)$ are either all non-negative or all non-positive.

1.2. Simplicial cone decomposition of $V_{\mathbb{R}}$.

For a subset $F \subset R(W)$, consider the subspace $H_F := \bigcap_{\beta \in F} H_\beta$ of $V_{\mathbb{R}}$ and the set of hyperplanes of H_F induced by reflection hyperplanes:

$$(1.2.1) \quad A(H_F) := \{H_F \cap H_\alpha \mid \alpha \in R(W), H_\alpha \not\subset H_F\}.$$

A point in H_F is called *generic* if it lies in $\dot{H}_F := H_F \setminus \bigcup_{G \in A(H_F)} G$. A connected component of \dot{H}_F is called a *facet* of $V_{\mathbb{R}}$. Let Γ be the index set of all facets of $V_{\mathbb{R}}$ and let us denote by V_γ the facet corresponding to $\gamma \in \Gamma$. Then the vector space $V_{\mathbb{R}}$ decomposes into a disjoint union:

$$(1.2.2) \quad V_{\mathbb{R}} = \sqcup_{\gamma \in \Gamma} V_\gamma.$$

Put $\gamma \leq \delta$ for $\gamma, \delta \in \Gamma$ iff $V_\gamma \subset \bar{V}_\delta$. The decomposition is a stratification, i.e., it satisfies the *boundary condition*: if $V_\gamma \cap \bar{V}_\delta \neq \emptyset$ then $V_\gamma \subset \bar{V}_\delta$. The minimal element of Γ is denoted by 0 (i.e., $V_0 = \{0\}$). The maximal elements of Γ correspond to *chambers*. Any stratum is a cone over a simplex, and hence (1.2.2) is called the *simplicial cone decomposition*.

1.3. Polyhedron dual to the simplicial cone decomposition.

Definition. 1. A compact subset \bar{P} in \mathbb{R}^l with a fixed semi-algebraic stratification (a finite decomposition of \bar{P} into smooth semi-algebraic sets satisfying the boundary condition) is called a *semi-algebraic polyhedron*, if there is a semi-algebraic diffeomorphism, say φ , from \bar{P} to a polyhedron in \mathbb{R}^l (a convex hull of finite points in \mathbb{R}^l which has non-trivial interior points). More precisely, φ induces an isomorphism from each stratum to a facet of the polyhedron. A stratum of \bar{P} corresponding to a face, facet or vertex is called a face, facet or vertex of \bar{P} , respectively. The set P of interior points of \bar{P} is called an open semi-algebraic polyhedron. We say *the faces of \bar{P} are crossing normally at a point $x \in \bar{P}$* , if there is a real-analytic diffeomorphism from a neighborhood of x in \mathbb{R}^l to a neighborhood of the origin of \mathbb{R}^l which maps locally (\bar{P}, x) to $(\mathbb{R}_{\geq 0}^k \times \mathbb{R}^{l-k}, 0)$ for some $0 \leq k \leq l$.

2. A semi-algebraic polyhedron \bar{K} in $V_{\mathbb{R}}$ is called *dual* to the simplicial cone decomposition (1.2.2), if it has the facet decomposition:

$$(1.3.1) \quad \bar{K} = \sqcup_{\gamma \in \Gamma} K_\gamma$$

indexed by the same index set Γ as in (1.2.2) such that

- i) $\bar{K}_\gamma \supset \bar{K}_\delta$ if and only if $\gamma \leq \delta$,
- ii) $K_\gamma \cap V_\delta \neq \emptyset$ if and only if $\gamma \leq \delta$, for any $\gamma, \delta \in \Gamma$,
- iii) if $\gamma \leq \delta$, then K_γ and V_δ intersects transversally at each point of $K_\gamma \cap V_\delta$. There exists a real analytic diffeomorphism from a neighborhood of $\overline{K_\gamma \cap V_\delta}$ to a neighborhood of the cube $[0, 1]^k$ of dimension $k = \dim(V_\delta) - \dim(V_\gamma)$, which induces a homeomorphism from $\overline{K_\gamma \cap V_\delta}$ to $[0, 1]^k$. In particular, $\dim(K_\gamma) + \dim(V_\gamma) = l$ for $\gamma \in \Gamma$.

The last condition iii) implies the following property:

iv) the faces of \overline{K} are crossing normally everywhere on \overline{K} .

The definition implies that K_0 is an open cell in $V_{\mathbb{R}}$ containing $0 \in V_{\mathbb{R}}$ such that $K = K_0$. The simpliciality of the cone decomposition (1.2.2) implies that \overline{K} is a *manifold with corners*.

1.4. Invariants for W and the quotient variety S_W .

We recall basic facts on W -invariants $S(V_{\mathbb{R}}^*)^W$ ([B, Ch.v, §5]) and fix notation on the W -quotient space.

1. A product $c := \prod_{\alpha \in \Pi(C)} \alpha$ is called a *Coxeter element*. Its conjugacy class in W is independent of the order of the product. The order h of c is called the *Coxeter number*. The eigenvalues of c are given by $\exp(2\pi\sqrt{-1}m_i/h)$ ($i = 1, \dots, l$) where $0 < m_i < h$ are called the *exponents* of W and are ordered as $m_1 = 1 < m_2 \leq \dots \leq m_{l-1} < m_l = h-1$.

2. Let $S(V_{\mathbb{R}}^*)$ be the symmetric tensor algebra of $V_{\mathbb{R}}^*$. We denote by $S(V_{\mathbb{R}}^*)^W$ the subring consisting of W -invariants in $S(V_{\mathbb{R}}^*)$. Chevalley's Theorem [Ch] states that $S(V_{\mathbb{R}}^*)^W$ is generated by l algebraically independent homogeneous elements of degrees $m_i + 1$ ($i = 1, \dots, l$). In the rest of the paper, we fix a homogeneous generator system (P_1, \dots, P_l) with $d_i := \deg P_i = m_i + 1$. Therefore, we have $S(V_{\mathbb{R}}^*)^W \simeq \mathbb{R}[P_1, \dots, P_l]$.

3. The module of anti-invariants $S(V_{\mathbb{R}}^*)^{-W} := \{P \in S(V_{\mathbb{R}}^*) \mid g(P) = \det(g)^{-1} P \text{ for all } g \in W\}$ is a free $S(V_{\mathbb{R}}^*)^W$ -module of rank one generated by

$$(1.4.1) \quad \delta_W := \prod_{\alpha \in R(W)} f_{\alpha}.$$

The Jacobian of the generator system (P_1, \dots, P_l) of invariants with respect to a linear coordinates (X_1, \dots, X_l) of $V_{\mathbb{R}}$ is a basic anti-invariant:

$$(1.4.2) \quad \det\left(\frac{\partial(P_1, \dots, P_l)}{\partial(X_1, \dots, X_l)}\right) = c \delta_W \quad \text{for } c \in \mathbb{R}_{\neq 0}.$$

4. Let $\Omega := \exp(\pi\sqrt{-1}/h)$ be a primitive $(2h)$ th root of unity. The eigenvector ξ of a Coxeter element belonging to the eigenvalue Ω^2 in the complexification $V_{\mathbb{C}} = \mathbb{C} \otimes V_{\mathbb{R}}$ is regular, i.e., $\delta_W(\xi) \neq 0$ ([B, Ch.V, §6]). This implies an equality (c.f. §2.4 Fact 1):

$$(1.4.3) \quad \#R(W) = h \cdot l/2.$$

5. The square $\Delta_W := \delta_W^2$ is a W -invariant called the *discriminant*. Express Δ_W as a polynomial in P_l . In view of the degree counting: $\deg(P_l) = h$ and $\deg(\Delta) = hl$, we know that it is of the form:

$$(1.4.4) \quad \Delta_W = A_0 P_l^l + A_1 P_l^{l-1} + \dots + A_l$$

where A_i is a homogeneous polynomial of degree ih in P_1, \dots, P_{l-1} . Then $A_0 \neq 0$ (since, by the degree condition, one has $P_1(\xi) = \dots = P_{l-1}(\xi) = 0$). Then $\Delta_W(\xi) \neq 0$ implies $A_0 \neq 0$ and $P_l(\xi) \neq 0$.

6. The categorical quotient variety $V//W$ as a scheme over \mathbb{R} is given by

$$(1.4.5) \quad S_W := V//W := \text{Spec}(S(V_{\mathbb{R}}^*)^W),$$

and its \mathbb{C} -rational point set is given by

$$(1.4.6) \quad S_{W,\mathbb{C}} := \text{Hom}_{\mathbb{R}}^{alg}(S(V_{\mathbb{R}}^*)^W, \mathbb{C}) = \text{Hom}_{\mathbb{C}}^{alg}(S(V_{\mathbb{C}}^*)^W, \mathbb{C}),$$

where Hom^{alg} is the set of algebra homomorphisms. The image in $S_{W,\mathbb{C}}$ of the origin of $V_{\mathbb{C}}$ is denoted by o and is called the origin of $S_{W,\mathbb{C}}$.

For $\varepsilon \in \{\pm 1\}$, we consider the real form $V_{\mathbb{R}}^{\varepsilon}$ of $V_{\mathbb{C}} := V_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ where

$$(1.4.7) \quad V_{\mathbb{R}}^{+1} := V_{\mathbb{R}} \quad \text{and} \quad V_{\mathbb{R}}^{-1} := \sqrt{-1}V_{\mathbb{R}}.$$

The \mathbb{C} -linear W -action on $V_{\mathbb{C}}$ leaves the real forms invariant such that $S((V_{\mathbb{R}}^{\varepsilon})^*)^W \otimes_{\mathbb{R}} \mathbb{C} \simeq S(V_{\mathbb{C}}^*)^W$. Thus, we introduce two real forms of $S_{W,\mathbb{C}}$:

$$(1.4.8) \quad S_{W,\mathbb{R}}^{[\varepsilon]} := \text{Hom}_{\mathbb{R}}^{alg}(S((V_{\mathbb{R}}^{\varepsilon})^*)^W, \mathbb{R})$$

for $\varepsilon \in \{\pm 1\}$. These two real forms coincide if $-\text{id}_{V_{\mathbb{R}}} \in W$. Note that a real coordinate system of $S_{W,\mathbb{R}}^{[\varepsilon]}$ is given by $(P_i/\sqrt{\varepsilon^{m_i+1}})_{i=1}^l$ so that

$$(1.4.9) \quad (P_1/\sqrt{\varepsilon^2}, \dots, P_l/\sqrt{\varepsilon^h}) : S_{W,\mathbb{R}}^{[\varepsilon]} \xrightarrow{\sim} \mathbb{R}^l,$$

where we put $\sqrt{1} := 1$ and $\sqrt{-1} :=$ the unit of pure imaginary number.

7. For any point $x \in V_{\mathbb{C}}$, the evaluation homomorphism: $S(V_{\mathbb{C}})^W \ni P \mapsto P(x) \in \mathbb{C}$ induces the W -invariant morphisms:

$$(1.4.10) \quad \pi_{W,\mathbb{C}} : V_{\mathbb{C}} \rightarrow S_{W,\mathbb{C}} \quad \text{and} \quad \pi_{W,\mathbb{R}}^{\varepsilon} : V_{\mathbb{R}}^{\varepsilon} \rightarrow S_{W,\mathbb{R}}^{[\varepsilon]} \quad (\varepsilon \in \{\pm 1\}).$$

These morphisms are finite and closed maps with respect to the classical topology. The morphism $\pi_{W,\mathbb{C}}$ induces a homeomorphism $V_{\mathbb{C}}/W \simeq S_{W,\mathbb{C}}$, and $\pi_{W,\mathbb{R}}^{\varepsilon}$ induces an embedding $V_{\mathbb{R}}^{\varepsilon}/W \subset S_{W,\mathbb{R}}^{[\varepsilon]}$ onto a closed semi-algebraic set (see Assertion 1.1 (4)).

1.5. Discriminant divisor and the central component $\mathcal{C}^{\{\varepsilon\}}$.

The *discriminant divisor* D_W in S_W is defined by $\Delta_W = 0$. Its \mathbb{C} -rational point set in $S_{W,\mathbb{C}}$ or \mathbb{R} -rational point set in $S_{W,\mathbb{R}}^{[\varepsilon]}$ for $\varepsilon \in \{\pm 1\}$ (called also the *complex or real discriminant locus*) are given by

$$(1.5.1) \quad D_{W,\mathbb{C}} := \{t \in S_{W,\mathbb{C}} \mid \Delta_W(t) = 0\} \quad \text{and} \quad D_{W,\mathbb{R}}^{[\varepsilon]} := D_{W,\mathbb{C}} \cap S_{W,\mathbb{R}}^{[\varepsilon]}.$$

The equalities (1.4.1) and (1.4.2) imply:

- i) The critical values of $\pi_{W,\mathbb{C}}$ lie in the discriminant divisor $D_{W,\mathbb{C}}$.
- ii) The inverse image $\pi_{W,\mathbb{C}}^{-1}D_{W,\mathbb{C}}$ is the union $\bigcup_{\alpha \in R(W)} H_{\alpha,\mathbb{C}}$.

Assertion 1.1. (1) *The stabilizer subgroup of W at any point $x \in V_{\mathbb{C}}$ is generated by the reflections whose reflection hyperplanes contain x .*

(2) *The complement of the discriminant locus $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ is the space of regular orbits (i.e., stabilizer free) of the W -action on $V_{\mathbb{C}}$.*

(3) $\pi_{W,\mathbb{C}} : V_{\mathbb{C}} \setminus \bigcup_{\alpha \in R(W)} H_{\alpha,\mathbb{C}} \rightarrow S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ *is a normal covering whose covering transformation group is W .*

(4) *For $\varepsilon \in \{\pm 1\}$, there exists a connected component $\mathcal{C}^{\{\varepsilon\}}$ of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$ such that for any connected component (chamber) C of $V_{\mathbb{R}} \setminus \bigcup_{\alpha \in R(W)} H_{\alpha}$, the morphism $\pi_{W,\mathbb{R}}^{\varepsilon}$ induce the homeomorphisms:*

$$(1.5.2) \quad \sqrt{\varepsilon}C \simeq \mathcal{C}^{\{\varepsilon\}} \quad \text{and} \quad \sqrt{\varepsilon}\overline{C} \simeq \overline{\mathcal{C}}^{\{\varepsilon\}}.$$

We call $\mathcal{C}^{\{\varepsilon\}}$ the central component of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$.

(5) *As a consequence of (4), $\overline{\mathcal{C}}^{\{\varepsilon\}}$ is a semi-algebraic simplicial cone with the vertex at o , whose faces are indexed by $\Pi = \Pi(C)$.*

1.6. Primitive vector field D and \mathbb{G}_a -action τ on S_W .

We fix a particular vector field D on S_W , which we shall call the *primitive vector field* ([S3,(2.2)]). The vector field D is transversal to the discriminant divisor D_W and plays a basic role throughout the present paper.

Let Der_{S_W} be the module of derivations of the algebra $S(V_{\mathbb{R}}^*)^W$ over \mathbb{R} , which is a graded $S(V_{\mathbb{R}}^*)^W$ -module. Using the generator system P_1, \dots, P_l for $S(V_{\mathbb{R}}^*)^W$ (see 2. of §1.4), its free basis are given by ∂_{P_i} ($i = 1, \dots, l$) with $\partial_{P_i} P_j = \delta_{ij}$ and $\deg(\partial_{P_i}) = -\deg(P_i)$. The maximality $\deg(P_l) > \deg(P_i)$ for $i = 1, \dots, l - 1$, implies that the lowest graded piece of Der_{S_W} is a vector space of dimension one spanned by

$$(1.6.1) \quad D := \partial_{P_l}.$$

In the rest of the paper, we fix a basis (1.6.1) and call it the *primitive vector field*.

Remark 1. The primitive vector field is one of the basic building blocks for the flat structure on S_W , but we do not go into details ([S1,3]).

Integrating D , we introduce a group action

$$(1.6.2) \quad \tau : \mathbb{G}_a \times S_W \longrightarrow S_W,$$

whose co-action τ^* on $S(V_{\mathbb{R}}^*)^W$ is given by

$$(1.6.3) \quad \begin{aligned} \tau^* : S(V_{\mathbb{R}}^*)^W &\longrightarrow S(V_{\mathbb{R}}^*)^W \otimes \mathbb{R}[\lambda], \\ P_i &\mapsto P_i \quad (i = 1, \dots, l - 1) \quad \text{and} \quad P_l \mapsto P_l + \lambda. \end{aligned}$$

Note that $(\tau(\mathbb{C}) \cdot o) \cap D_{W,\mathbb{C}} = \{o\}$ where o is the origin of $S_{W,\mathbb{C}}$, since the leading coefficient A_0 in (1.4.4) does not vanish.

For each $\varepsilon \in \{\pm 1\}$, let us choose and fix the real valued function

$$\lambda^{[\varepsilon]} := \lambda / \sqrt{\varepsilon^h}$$

on the real form $\mathbb{G}_a^{\varepsilon^h} = \sqrt{\varepsilon^h} \mathbb{R} \subset \mathbb{G}_{a,\mathbb{C}} = \mathbb{C}$ as its real coordinate. Then, recalling (1.4.9), one obtains the real one-parameter group action:

$$(1.6.4) \quad \begin{array}{ccc} \tau^{[\varepsilon]} & : & \mathbb{R} \times S_{W,\mathbb{R}}^{[\varepsilon]} \rightarrow S_{W,\mathbb{R}}^{[\varepsilon]} \\ \lambda^{[\varepsilon]} \times (P_1/\sqrt{\varepsilon^2}, \dots, P_l/\sqrt{\varepsilon^h}) & \mapsto & (P_1/\sqrt{\varepsilon^2}, \dots, P_l/\sqrt{\varepsilon^h} + \lambda^{[\varepsilon]}). \end{array}$$

A domain in $S_{W,\mathbb{R}}^{[\varepsilon]}$ is called a *tube domain* if it is $\tau^{[\varepsilon]}$ -invariant.

1.7. The opposite components $\mathcal{C}_\pm^{[\varepsilon]}$ of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$.

Since the half lines $\tau^{[\varepsilon]}(\mathbb{R}_{>0}) \cdot o$ and $\tau^{[\varepsilon]}(-\mathbb{R}_{>0}) \cdot o$ do not intersect the discriminant locus, we have the following definition.

Definition. The *opposite components* of $S_{W,\mathbb{R}}^{[\varepsilon]} \setminus D_{W,\mathbb{R}}^{[\varepsilon]}$ are

$$(1.7.1) \quad \begin{array}{l} \mathcal{C}_+^{[\varepsilon]} := \text{the connected component which contains } \tau^{[\varepsilon]}(\mathbb{R}_{>0}) \cdot o, \\ \mathcal{C}_-^{[\varepsilon]} := \text{the connected component which contains } \tau^{[\varepsilon]}(\mathbb{R}_{<0}) \cdot o. \end{array}$$

One has: $\mathcal{C}_+^{[\varepsilon]} \neq \mathcal{C}^{\{\varepsilon\}} \neq \mathcal{C}_-^{[\varepsilon]}$ (except for type A_1), since the eigenvectors for $\exp(2\pi\sqrt{-1}/h)$ of the Coxeter element do not belong to $V_{\mathbb{R}}^\varepsilon$. Each of the opposite components $\mathcal{C}_\pm^{[\varepsilon]}$ is the interior of the quotient of a certain twisted real form of $V_{\mathbb{C}}$. We shall give another expression of opposite components in (2.5.5) by determining the twisted real form.

1.8. Semi-algebraic sets $\bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ in $S_{W,\mathbb{R}}^{[\varepsilon]}$ and $\bar{K}_W^{\varepsilon}(\lambda^{[\varepsilon]})$ in $V_{\mathbb{R}}^\varepsilon$.

We state Theorem A announced in the introduction.

Theorem A. For $\lambda^{[\varepsilon]} \in \mathbb{R}_{>0}$ and for $\varepsilon \in \{\pm 1\}$, put

$$(1.8.1) \quad J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]}) := \mathcal{C}^{\{\varepsilon\}} \cap \tau^{[\varepsilon]}(-\lambda^{[\varepsilon]})\mathcal{C}_+^{[\varepsilon]} \cap \tau^{[\varepsilon]}(\lambda^{[\varepsilon]})\mathcal{C}_-^{[\varepsilon]}.$$

Then $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ is an open semi-algebraic polyhedron in $S_{W,\mathbb{R}}^{[\varepsilon]}$ isomorphic to the l -dimensional parallelotope $(0, \lambda^{[\varepsilon]})^l$ adjacent to the origin $o \in S_{W,\mathbb{C}}$. Let $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ be the vertex of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ which is antipodal to the origin. Then faces in $\bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ are crossing normally at any point of any closed edge adjacent to $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$.

Remark 2. The explicit identification $c_W : \bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]}) \simeq [0, -\lambda^{[\varepsilon]}]^{l_1} \times [0, \lambda^{[\varepsilon]}]^{l_2}$ with $l = l_1 + l_2$ is given in Theorem C in §3.5.

In an assertion in §3.4, we prove a stronger normal crossing property of faces of $\bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$, which implies that the inverse image in $V_{\mathbb{R}}^\varepsilon$ of any facet of $\bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ adjacent to $ao^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ is smooth. This gives the next theorem, stated as Theorem B in the introduction.

Theorem B. For $\lambda^{[\varepsilon]} \in \mathbb{R}_{>0}$ and for $\varepsilon \in \{\pm 1\}$, put

$$\begin{aligned} K_W^\varepsilon(\lambda^{[\varepsilon]}) : &= (\pi_{W,\mathbb{R}}^\varepsilon)^{-1} \left(J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]}) \right) \\ &= (\pi_{W,\mathbb{R}}^\varepsilon)^{-1} \left(\tau^{[\varepsilon]}(-\lambda^{[\varepsilon]})\mathcal{C}_+^{[\varepsilon]} \cap \tau^{[\varepsilon]}(\lambda^{[\varepsilon]})\mathcal{C}_-^{[\varepsilon]} \right). \end{aligned}$$

Then $K_W^\varepsilon(\lambda^{[\varepsilon]})$ is an open semi-algebraic polyhedron in $V_{\mathbb{R}}^\varepsilon$ dual to the simplicial cone decomposition (1.2.2). The W -action induces

$$(1.8.3) \quad \bar{K}_W^\varepsilon(\lambda^{[\varepsilon]})/W \cong \pi_{W,\mathbb{R}}^\varepsilon(\bar{K}_W^\varepsilon(\lambda^{[\varepsilon]})) = \bar{J}_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]}).$$

Remark 3. The change D to ωD for $\omega \in \mathbb{R}^\times$ induces the change $K_W^\varepsilon(\lambda^{[\varepsilon]}, D) = |\frac{\lambda^{[\varepsilon]}}{\lambda^{[\varepsilon]'}} \omega|^{1/h} K_W^\varepsilon(\lambda^{[\varepsilon]',} \omega D)$ for $\lambda^{[\varepsilon]}, \lambda^{[\varepsilon]'} \in \mathbb{R}_{>0}$, i.e. the polyhedra for any scale ω and any parameter $\lambda^{[\varepsilon]}$ are homothetic to each other.

2. THE CENTRAL REGION $E_W^{\{\varepsilon\}}$ IN $T_{W,\mathbb{R}}^{[\varepsilon]}$

The concept of a universal unfolding and its bifurcation set is due to Thom [T2] and is studied by several authors (e.g. [Ar],[Lo1],[Ly],[Te]). We re-introduce the bifurcation set in the setting of W - τ -invariant theory, and then introduce the central region $E_W^{\{\varepsilon\}}$, which is a key concept in the present paper. Several results are extracted from [S4]. For a comprehensive study of them, one is referred to [S4].

2.1. τ -quotient space T_W and τ -quotient morphism π_τ .

We first introduce the τ -quotient space and the τ -quotient morphism. Recall the co-action (1.6.3) and consider the ring of its invariants:

$$(2.1.1) \quad \begin{aligned} S(V_{\mathbb{R}}^*)^{W,\tau} &:= \{f \in S(V_{\mathbb{R}}^*)^W \mid \tau^*(\lambda)f = f\} \\ &= \{f \in S(V_{\mathbb{R}}^*)^W \mid Df = 0\}. \end{aligned}$$

The associated τ -quotient variety is denoted by

$$(2.1.2) \quad T_W := \text{Spec}(S(V_{\mathbb{R}}^*)^{W,\tau}).$$

One has the τ -quotient morphism

$$(2.1.3) \quad \pi_\tau : S_W \longrightarrow T_W$$

induced by the inclusion $S(V_{\mathbb{R}}^*)^{W,\tau} \subset S(V_{\mathbb{R}}^*)^W$. Using the coordinates P_1, \dots, P_l , one has the explicit expressions:

$$S(V_{\mathbb{R}}^*)^{W,\tau} = \mathbb{R}[P_1, \dots, P_{l-1}] \quad \text{and} \quad S(V_{\mathbb{R}}^*)^W = S(V_{\mathbb{R}}^*)^{W,\tau}[P_l].$$

Namely, T_W is an affine variety with the coordinates (P_1, \dots, P_{l-1}) and π_τ is the projection forgetting the last coordinate P_l .

2.2. Bifurcation divisor $B_W = \cup_{p=2}^\infty B_{W,p}$.

We introduce the bifurcation set B_W in T_W as the ramification divisor of the finite cover $\pi_\tau|D_W : D_W \rightarrow T_W$. Recall that the discriminant Δ_W is a monic polynomial in P_l (1.4.4). The resultant of Δ_W and $D\Delta_W = \partial_{P_l}\Delta_W$ with respect to P_l is an element in $S(V_{\mathbb{R}}^*)^{W,\tau}$. Decompose it as

$$(2.2.1) \quad \delta(\Delta_W, D\Delta_W) = \prod_{p \geq 2} \omega_{W,p}^p,$$

according to its multiplicity (=ramification index), where $\omega_{W,p}$ are multiplicity-free polynomials in $S(V_{\mathbb{R}}^*)^{W,\tau}$. Using the p -th factor $\omega_{W,p}$, the p -th bifurcation divisor is defined by the equation:

$$(2.2.2) \quad B_{W,p} := \text{the divisor in } T_W \text{ defined by } \omega_{W,p} = 0.$$

We define the ordinary part $B_{W,2}$, the odd part $B_{W,odd} := \cup_{p: \text{odd}} B_{W,p}$ and the higher part $B_{W,\geq 3} := \cup_{p \geq 3} B_{W,p}$ of the bifurcation divisor.

Note. The p th bifurcation divisor $B_{W,p}$ is the image of the union of two-codimensional subspaces of V where p reflection hyperplanes are intersecting (see the last formula $(*)$ at the end of Appendix in order to justify the decomposition (2.2.1)).

One basic formula which plays a key role in the sequel is the following:

$$(2.2.3) \quad \det \left(\frac{\partial(\Delta_W, D\Delta_W, \dots, D^{l-1}\Delta_W)}{\partial(P_1, P_2, \dots, P_l)} \right) = c \cdot \prod_{p \geq 2} \omega_{W,p}^{p-1},$$

The proof uses the degree of $\omega_{W,p}$, obtained by the case by case study (see [S4,(3.6.1)]). The author does not know any other proof.

2.3. Real forms of the τ -action and the τ -quotient space.

Let $u \in \text{GL}(V_{\mathbb{R}})$ be an element of the normalizer $N(W)$ of W . We denote by $[u]$ its W -coset class in $N(W)/W$. Assume $[u]^2 = 1$ and define an anti- \mathbb{C} -linear automorphism $[u]^{a*} : S(V_{\mathbb{C}}^*)^W \rightarrow S(V_{\mathbb{C}}^*)^W$ by $[u]^{a*}P := \overline{P} \circ u$. Then the twisted real form $S_{W,\mathbb{R}}^{[u]} := \text{Hom}_{\mathbb{R}}(S(V_{\mathbb{C}}^*)^{W,[u]^{a*}}, \mathbb{R})$ is given by the subalgebra $S(V_{\mathbb{C}}^*)^{W,[u]^{a*}}$ of $[u]^{a*}$ -invariants (see [Lo2],[S4]).

Assertion 2.1. *There exists $b[u] \in \{\pm 1\}$ making the following diagram commutative:*

$$(2.3.1) \quad \begin{array}{ccc} S(V_{\mathbb{C}}^*)^W & \xrightarrow{\tau^*} & S(V_{\mathbb{C}}^*)^W \otimes_{\mathbb{C}} \mathbb{C}[\lambda] \\ \downarrow [u]^{a*} & & \downarrow [u]^{a*} \otimes (b[u] \circ \text{complex conjugation}) \\ S(V_{\mathbb{C}}^*)^W & \xrightarrow{\tau^*} & S(V_{\mathbb{C}}^*)^W \otimes_{\mathbb{C}} \mathbb{C}[\lambda] \end{array} .$$

One can choose a generator P_l satisfying $[u]^{a*}P_l = b[u]P_l$ so that $P_l^{[u]} := P_l / \sqrt{b[u]}$ is a $[u]^{a*}$ -invariant. The co-action τ^* (1.6.3) turns out to be

$$(2.3.2) \quad \tau^* : S(V_{\mathbb{C}}^*)^{W,[u]^{a*}} \rightarrow S(V_{\mathbb{C}}^*)^{W,[u]^{a*}} \otimes_{\mathbb{R}} \mathbb{R}[\lambda / \sqrt{b[u]}].$$

Accordingly, we introduce a new real variable $\lambda^{[u]} := \lambda / \sqrt{b[u]}$ so that we obtain a twisted real τ -action

$$(2.3.3) \quad \begin{array}{ccc} \tau^{[u]} : \mathbb{R} \times S_{W,\mathbb{R}}^{[u]} & \longrightarrow & S_{W,\mathbb{R}}^{[u]} \\ \lambda^{[u]} \times (P_1^{[u]}, \dots, P_l^{[u]}) & \longmapsto & (P_1^{[u]}, \dots, P_l^{[u]} + \lambda^{[u]}). \end{array}$$

The $\tau^{[u]}$ -invariants $S(V_{\mathbb{C}}^*)^{W,[u]^{a*},\tau}$ defines a twisted real form

$$(2.3.4) \quad T_{W,\mathbb{R}}^{[u]} := \text{Hom}_{\mathbb{R}}^{alg}(S(V_{\mathbb{C}}^*)^{W,[u]^{a*},\tau}, \mathbb{R})$$

of the space $T_{W,\mathbb{C}}$ together with the twisted real quotient morphism

$$(2.3.5) \quad \pi_{\tau}^{[u]} : S_{W,\mathbb{R}}^{[u]} \longrightarrow T_{W,\mathbb{R}}^{[u]}.$$

One sees from the coordinate expression that the morphism (2.3.5) is an honest set-theoretical quotient map. In the present article, we are interested in the case $[u] = [\varepsilon]$ for $\varepsilon \in \{\pm 1\}$. In such a case, $b[u] = \varepsilon^h$.

2.4. Subspace $S_{W(I_2(h))}$ of S_W .

We introduce a canonical two-dimensional subspace $S_{W(I_2(h))}$ of S_W based on the study of regular eigenvectors of a Coxeter element due to Coleman [C] and Kostant [K] (cf. [B, Ch.5, §6, n°2, lemma 2]).

Let C be a Weyl chamber in $V_{\mathbb{R}}$ of W and let $\Pi = \Pi(C)$ be the set of simple reflections. Recall the Coxeter diagram structure $\Gamma(W)$ on Π , where one puts an edge between two vertices α and β if $m_{\alpha\beta} \geq 3$. Since $\Gamma(W)$ is a tree, one has a unique decomposition up to a transposition:

$$(2.4.1) \quad \Pi = \Pi_1 \sqcup \Pi_2$$

where each Π_j is a totally disconnected subset in $\Gamma(W)$. Put

$$(2.4.2) \quad c_i := \prod_{\alpha \in \Pi_i} \alpha, \quad c := c_1 c_2 \quad \text{and} \quad d := c_1 + c_2,$$

where c and d are called the Coxeter element and the Killing element. $c = c_1 c_2$ is of order h , and d has only real eigenvalues $2 \cos(\pi m_i/h)$. Let e be an eigenvector of d belonging to the largest eigenvalue $2 \cos(\pi/h)$ (which has multiplicity 1). We can choose e such that its expression with respect to the basis e_α ($\alpha \in \Pi$) (see 1.1, 4.) has all positive coefficients, so that e is unique up to a positive constant multiple. Put

$$(2.4.3) \quad U := \mathbb{R}e_1 + \mathbb{R}e_2,$$

where we use the decomposition $e = e_1 + e_2$ with $e_j \in \sum_{\alpha \in \Pi_j} \mathbb{R}_{>0} e_\alpha$.

Assertion 2.2. c_i leaves the space U invariant and $c_i|U$ is a reflection with respect to e_i . The restriction homomorphism $\langle c_1, c_2 \rangle \rightarrow \langle c_1|U, c_2|U \rangle$ is an isomorphism and defines a faithful $W(I_2(h))$ -action on U .

Here $W(I_2(h))$ is the dihedral group of order $2h$ generated by reflections $c_1|U$ and $c_2|U$. Therefore, we introduce

- i) the quotient space $S_{W(I_2(h))} := U//W(I_2(h))$,
- ii) a primitive vector field $\frac{\partial}{\partial S}$ (which, we fix explicitly below),
- iii) the τ -action on $S_{W(I_2(h))}$ as the integration of $\frac{\partial}{\partial S}$,
- iv) the τ -quotient variety $T_{W(I_2(h))} := S_{W(I_2(h))} // \tau$.

The embedding $U \subset V_{\mathbb{R}}$ induces the morphism $S_{W(I_2(h))} \rightarrow S_W$. The following facts are not difficult, but are not obvious (cf. [S4]).

Assertion 2.3. *i) The morphism $S_{W(I_2(h))} \rightarrow S_W$ is a closed embedding. Its image is independent of the choices of a chamber C , a decomposition (2.4.1) and an eigenvector e .*

ii) The primitive vector field on S_W is tangent to $S_{W(I_2(h))}$, and induces a primitive vector field on the subspace $S_{W(I_2(h))}$.

iii) The restriction of the τ -action on S_W coincides with the τ -action on $S_{W(I_2(h))}$ (up to a scaling constant).

iv) One obtains a canonical embedding $T_{W(I_2(h))} \rightarrow T_W$ which makes the following diagram commutative and Cartesian:

$$(2.4.4) \quad \begin{array}{ccc} S_{W(I_2(h))} & \longrightarrow & S_W \\ \downarrow \pi_\tau & & \downarrow \pi_\tau \\ T_{W(I_2(h))} & \longrightarrow & T_W \end{array}$$

The image of $S_{W(I_2(h))}$ in S_W is called the *vertex orbit plane* and that of $T_{W(I_2(h))}$ is called the *vertex orbit line*. In the sequel, we write v.o. for vertex orbit for short (see the introduction for the naming).

2.5. V.o. axis, v.o. line and the sign factor $\sigma(D, \{\Pi_1, \Pi_2\})$.

Depending on a choice of the vector e and the partition $\{\Pi_1, \Pi_2\}$, we obtain some particular generators of $S[U^*]^{W(I_2(h))}$ (called the flat coordinates), which leads to some new concepts and constructions.

First, we identify the vector space U with \mathbb{C} regarded as a real vector space $\mathbb{R} \oplus \mathbb{R}i$, where, in order to avoid the confusion with the complex number field as the coefficient field, we use notation i for the unit of pure imaginary number instead of $\sqrt{-1}$.

i) The identification $U \simeq \mathbb{R} \oplus \mathbb{R}i$ is given by the basis correspondence: $e_1 \leftrightarrow i$ and $e_2 \leftrightarrow -i\omega$, where $\omega := \exp(\pi i/h) = \cos(\pi/h) + \sin(\pi/h)i$.

ii) For $z_1, z_2 \in \mathbb{C} \simeq U$, put $I(z_1, z_2) = \text{Re}(z_1 \bar{z}_2)$.

iii) The generators $c_1|U$ and $c_2|U$ of $W(I_2(h))$ are the reflections with respect to e_1 and e_2 : $c_1(z) = \bar{z}$ and $c_2(z) = \omega^2 \bar{z}$ on \mathbb{C} .

iv) The Coxeter element $c|U$ of $W(I_2(h))$ is identified with the multiplication by ω^{-2} .

v) We choose the generators R and S of $S[U^*]^{W(I_2(h))}$:

$$(2.5.1) \quad \begin{aligned} R &:= R(\{\Pi_1, \Pi_2\}, e) := z\bar{z} = x^2 + y^2, \\ S &:= S(\{\Pi_1, \Pi_2\}, e) := \text{Re}(z^h) = \sum_{k=0}^{\lfloor h/2 \rfloor} (-1)^k C_{2k}^h x^{h-2k} y^{2k}. \end{aligned}$$

vi) The changes $e \mapsto re$ ($r \in \mathbb{R}_{\geq 0}$) and $\{\Pi_1, \Pi_2\} \mapsto \{\Pi_2, \Pi_1\}$ induce:

$$(2.5.2) \quad \begin{aligned} R(\{\Pi_1, \Pi_2\}, re) &= r^2 R(\{\Pi_1, \Pi_2\}, e) \\ S(\{\Pi_1, \Pi_2\}, re) &= r^h S(\{\Pi_1, \Pi_2\}, e) \end{aligned}$$

$$(2.5.3) \quad \begin{aligned} R(\{\Pi_2, \Pi_1\}, e) &= R(\{\Pi_1, \Pi_2\}, e) \\ S(\{\Pi_2, \Pi_1\}, e) &= -S(\{\Pi_1, \Pi_2\}, e). \end{aligned}$$

After the preparation above, we can now clarify several sign problems as follows.

1. **Sign factor.** The derivation $\frac{\partial}{\partial S}$ is a primitive vector field on $S_{W(I_2(h))}$. Due to Assertion 2.3, ii), the proportion $\frac{\partial}{\partial S} : D|S_{W(I_2(h))}$ is

in \mathbb{R}^\times . Depending on the choice of the primitive vector field D (1.6.1) and the decomposition $\{\Pi_1, \Pi_2\}$ (2.4.1), we introduce the sign factor:

$$(2.5.4) \quad \sigma(D, \{\Pi_1, \Pi_2\}) := \text{sign}\left(\frac{\partial}{\partial S} : D|_{S_{W(I_2(h))}}\right) \in \{\pm 1\}.$$

A typical use of the sign factor is the following [S4,5.1,Sign Theorem]:

Assertion 2.4. *For $i \in \{1, 2\}$ and $\varepsilon \in \{\pm 1\}$, define the twisted real vector space with respect to εc_i by $V_{\mathbb{R}}^{\varepsilon c_i} := \{v \in V_{\mathbb{C}} \mid c_i(v) = \varepsilon \bar{v}\}$. Then $\pi_W(V_{\mathbb{R}}^{\varepsilon c_i})$ is the closure of a connected component of $S_{W, \mathbb{R}}^{[\varepsilon]} \setminus D_{W, \mathbb{R}}^{[\varepsilon]}$ denoted by $\mathcal{C}^{\{\varepsilon c_i\}}$. The components $\mathcal{C}^{\{\varepsilon c_i\}}$ ($i = 1, 2$) and the opposite components $\mathcal{C}_+^{[\varepsilon]}$ and $\mathcal{C}_-^{[\varepsilon]}$ (1.7.1) are related by the formula:*

$$(2.5.5) \quad \mathcal{C}^{\{\varepsilon c_i\}} = \mathcal{C}_{\varepsilon^{\lfloor h/2 \rfloor} (-1)^{i-1} \sigma(D, \{\Pi_1, \Pi_2\})}^{[\varepsilon]} \quad \text{for } i = 1, 2.$$

The sign factor appears again in Theorem C in §3.5.

2. Vertex orbit axis AO . Let us call the coordinate axis defined by $S = 0$ in the v.o. plane $S_{W(I_2(h))}$ the *vertex orbit axis*. It is a one-dimensional line in S_W . Note the fact that the coordinate R (c.f. (2.5.1)) is unique up to a positive constant multiple. Thus, the real v.o. axis and the half v.o. axis

$$(2.5.6) \quad \begin{aligned} AO &:= \{(R, S) \in S_{W(I_2(h)), \mathbb{C}} \mid R \in \mathbb{R} \text{ and } S = 0\}, \\ AO^\pm &:= \{(R, S) \in S_{W(I_2(h)), \mathbb{C}} \mid R \in \pm \mathbb{R}_{>0} \text{ and } S = 0\} \end{aligned}$$

are a real line and real half lines in $S_{W, \mathbb{C}}$ independent of the choices of a chamber C , a partition $\{\Pi_1, \Pi_2\}$ or a vector e .

Assertion 2.5. *i) For any $\varepsilon \in \{\pm 1\}$, the twisted real form $S_{W, \mathbb{R}}^{[\varepsilon]}$ contains the full real v.o. axis AO (see Remark 4. below).*

ii) For $\varepsilon \in \{\pm 1\}$, the connected component of $S_{W, \mathbb{R}}^{[\varepsilon]} \setminus D_{W, \mathbb{R}}^{[\varepsilon]}$ containing the half v.o. axis AO^ε is the central component \mathcal{C}^ε (recall (1.5.2)).

In the present article, we use AO^+ and AO^- as the base point locus for the fundamental group of $S_{W, \mathbb{C}} \setminus D_{W, \mathbb{C}}$.

3. Vertex orbit line O . Recall the one-dimensional subspace $T_{W(I_2(h))}$ of T_W (called the v.o. line) which is the projection image of the vertex orbit axis by π_τ (c.f. (2.4.4)). Similarly to the case of the vertex orbit axis, its coordinate R is unique up to a positive constant multiple. Thus, the real v.o. line and the half v.o. lines are given by

$$(2.5.7) \quad \begin{aligned} O &:= \{(R) \in T_{W(I_2(h))} \mid R \in \mathbb{R}\}, \\ O^\pm &:= \{(R) \in T_{W(I_2(h))} \mid R \in \pm \mathbb{R}_{>0}\}. \end{aligned}$$

They are a well-defined real line and real half lines in $T_{W, \mathbb{C}}$.

Assertion 2.6. *i) For any $\varepsilon \in \{\pm 1\}$, the twisted real form $T_{W,\mathbb{R}}^{[\varepsilon]}$ contains the real v.o. line O (see Remark 4. below).*

ii) For $\varepsilon \in \{\pm 1\}$, the half v.o. line O^ε does not intersect the higher bifurcation locus $B_{W,\geq 3,\mathbb{C}}$ in $T_{W,\mathbb{C}}$.

Definition. We introduce

$$(2.5.8) \quad E_W^{\{\varepsilon\}} := \begin{array}{l} \text{the connected component of } T_{W,\mathbb{R}}^{[\varepsilon]} \setminus B_{W,\geq 3,\mathbb{R}}^{[\varepsilon]} \\ \text{containing the half v.o. line } O^\varepsilon \end{array}$$

and call it the *central region* in $T_{W,\mathbb{R}}^{[\varepsilon]}$.

Remark 4. The statements i) in Assertion 2.5 and i) in 2.6 are valid for any twisted real structures $[u]$ on S_W and T_W , respectively. We study the connected component $E_W^{[u],\varepsilon}$ of $T_{W,\mathbb{R}}^{[u]} \setminus B_{W,\geq 3,\mathbb{R}}^{[u]}$ containing O^ε in [S4,§10].

Remark 5. Actually, the v.o. line O is contained in the ordinary bifurcation set $B_{W,2}$ for $l \geq 3$. Hence, the ordinary bifurcation set intersects the central region for $l \geq 3$. The description of the intersection $E_W^{\{\varepsilon\}} \cap B_{W,2,\mathbb{R}}^{[\varepsilon]}$ is reduced to certain real linear inequalities by use of Theorem C in §3.5. It proposes a quite interesting and important combinatorial geometric problem.

2.6. Algebroid functions φ_α and ϕ_α for $\alpha \in \Pi$.

Once we have fixed the base point locus, either O^+ or O^- , in $T_{W,\mathbb{C}}$, we are able to discuss multi-valued functions and, in particular, algebroid functions defined on $T_{W,\mathbb{C}} \setminus B_{W,\geq 3,\mathbb{C}}$.

Recall the discriminant (1.4.4). It is a monic polynomial of degree l in the indeterminate P_l . Let us consider the equation in P_l :

$$(2.6.1) \quad \Delta_W = A_0 P_l^l + A_1 P_l^{l-1} + \dots + A_{l-1} P_l + A_l = 0,$$

where the coefficients A_i are polynomial functions on $T_{W,\mathbb{C}}$. One can show that i) for any point of $T_{W,\mathbb{C}} \setminus B_{W,\text{odd},\mathbb{C}}$, there exist a neighborhood \mathcal{V} and l holomorphic functions f_1, \dots, f_l on \mathcal{V} such that $f_1(t), \dots, f_l(t)$ are the solutions of the equation (2.6.1) at $t \in \mathcal{V}$, and ii) the functions f_1, \dots, f_l can be analytically continued to everywhere in $T_{W,\mathbb{C}} \setminus B_{W,\text{odd},\mathbb{C}}$.

As for an initial system, let us choose l functions indexed by Π :

$$(2.6.2) \quad \varphi_{\alpha,\varepsilon} \quad (\alpha \in \Pi)$$

on a neighborhood of the base point locus O^ε as follows (see [S4,§8,9]).

For $\alpha \in \Pi_i$, let $\zeta_i \in U$ ($i = 1, 2$) be a point ($\neq 0$) fixed by the action of $c_i|U$ on the real 2-space U . Since $c_i|U$ is a reflection with respect to e_i (recall Assertion 2.2), we see $e_i \cdot \zeta_i = 0$, where $x \cdot y$ is a $W(I_2(h))$ -invariant positive symmetric bilinear form on U . Taking the fact $e_1 \cdot e_2 < 0$ into account, the coefficients $a, b \in \mathbb{R}$ in the expression

$\zeta_i = ae_1 + be_2$ are simultaneously positive or negative. We choose ζ_i such that $a, b > 0$. Then ζ_i is unique up to a positive constant multiple (see Remark 7).

For $\varepsilon \in \{\pm 1\}$, put $\zeta_i^\varepsilon := \sqrt{\varepsilon}\zeta_i \in U^\varepsilon := \sqrt{\varepsilon} \otimes U$ (see Remark 8). It projects by π_W to a point p_i on the discriminant locus $D_{W(I_2(h)),\mathbb{R}}^{[\varepsilon]} \subset S_{W(I_2(h)),\mathbb{R}}^{[\varepsilon]}$, and p_i projects further to a point $q_i \in O^\varepsilon \subset T_{W,\mathbb{R}}^{[\varepsilon]}$ by π_τ (c.f. (2.4.4)). On the other hand, the reflection hyperplane H_α contains ζ_i . Hence, $\zeta_i^\varepsilon \in H_{\alpha,\mathbb{C}}$. Since H_α ($\alpha \in \Pi_i$) are normally crossing at ζ_i , the stabilizer subgroup $W(\zeta_i)$ of ζ_i is an abelian group $\langle \beta, \beta \in \Pi_i \rangle \simeq (\mathbb{Z}_2)^{\#\Pi_i}$ and preserves each hyperplane H_α for $\alpha \in \Pi$. Then, it is easy to see that a neighborhood of ζ_i^ε in $H_{\alpha,\mathbb{C}}$ projects (by $2^{\#\Pi_i-1}$ to one) onto a neighborhood \mathcal{U}_α of p_i in one of $\#\Pi_i$ -number of local irreducible components of $D_{W,\mathbb{C}}$ at p_i . We can show that $q_i = \pi_\tau(p_i) \notin B_{W,\text{odd},\mathbb{C}}$ and that the projection $\pi_\tau|_{\mathcal{U}_\alpha}$ is a locally homeomorphism onto a neighborhood \mathcal{V}_α of q_i in $T_{W,\mathbb{C}}$. Then, we reverse the map $\pi_\tau|_{\mathcal{U}_\alpha}$ to a map $\varpi_\alpha : (\mathcal{V}_\alpha, q_i) \rightarrow (\mathcal{U}_\alpha, p_i)$, and put

$$\varphi_{\alpha,\varepsilon} := P_l \circ \varpi_\alpha.$$

By definition, $\varphi_{\alpha,\varepsilon}$ is a solution to the discriminant equation (2.6.1) on a neighborhood of O^ε . Thus we obtain the system of algebroid functions indexed by Π (2.6.2). By use of characteristic variety C_W , we observe that these give the full system of solutions of the discriminant equation. That is: one has the ‘‘local factorization’’ of the discriminant:

$$(2.6.3) \quad \Delta_W = A_0 \prod_{\alpha \in \Pi} (P_l - \varphi_{\alpha,\varepsilon}),$$

on a neighborhood of the base point locus O^ε . We set

$$(2.6.4) \quad \phi_{\alpha,\varepsilon} := P_l - \varphi_{\alpha,\varepsilon} \quad \text{for } \alpha \in \Pi.$$

Since $D^i \Delta_W$ is, up to the factor A_0 , equal to the $(l-i)$ -th elementary symmetric function of $\{\phi_{\alpha,\varepsilon}\}_{\alpha \in \Pi}$, the formula (2.2.3) can be rewritten as

$$(2.6.5) \quad \wedge_{\alpha \in \Pi} d\phi_{\alpha,\varepsilon} = c \prod_{p \geq 3} \omega_{W,p}^{p/2-1} \cdot \wedge_{i=1}^l dP_i \quad \text{for some } c \in \mathbb{R}^\times.$$

Remark 6. In the next §3.4, we introduce a largest covering space of $T_{W,\mathbb{C}} \setminus \mathbb{B}_{W,\text{odd},\mathbb{C}}$ with liftings of the base point loci O^+ and O^- such that $\varphi_{\alpha,+}$ and $\varphi_{\alpha,-}$ are lifted to functions defined on the neighborhoods of the base point loci and are analytically continued to the same univalent function.

Remark 7. In the above construction of $\varphi_{\alpha,\varepsilon}$, we have chosen ζ_i in such a manner that the coefficients with respect to the basis e_i are positive. However, we may choose $-\zeta_i$ as the starting point of the construction.

Then, for $\varepsilon \in \{\pm 1\}$, the function $\varphi_{\alpha,\varepsilon}$ changes to $\varphi_{\chi_W(\alpha),\varepsilon}$, where χ_W is the bijection of Π induced by the adjoint action of the longest element of W (see [S4,8.11], c.f. also (3.3.1)). This change is caused by the change of the “reference” chamber from C to $-C$, which covers the central component \mathcal{C} .

Remark 8. In the above construction of $\varphi_{\alpha,\varepsilon}$, we have chosen $\zeta_i^\varepsilon \in H_\alpha^\varepsilon$ to be $\zeta_i\sqrt{\varepsilon}$. However, we may choose its *complex conjugate* $\zeta_i/\sqrt{\varepsilon}$ for ζ_i^ε as the starting point of the construction. Then the $\varphi_{\alpha,+1}$ is unchanged, but the $\varphi_{\alpha,-1}$ changes to $\varphi_{\chi_W(\alpha),-1}$ (see Remark 6 for notation and reference). This change is caused by the change of the sign of the unit $\sqrt{-1}$ of the pure imaginary number.

3. LINEARIZATION MAP c_W

The system of algebroid functions $\phi_{\alpha,\varepsilon}$ and $\varphi_{\alpha,\varepsilon}$ introduced in the previous section define, by analytic continuation, multivalued holomorphic maps:

$$c_W = (\phi_\alpha)_{\alpha \in \Pi} : S_{W,\mathbb{C}} \dashrightarrow \mathbb{C}^\Pi \quad \& \quad b_W = (-\varphi_\alpha)_{\alpha \in \Pi} : T_{W,\mathbb{C}} \dashrightarrow \mathbb{C}^\Pi / \mathbb{C}v_\Pi$$

where i) $v_\Pi = (1_\alpha)_{\alpha \in \Pi}$ is the diagonal element, and ii) “ \dashrightarrow ” means that the “maps” are not univalent on $S_{W,\mathbb{C}}$ or on $T_{W,\mathbb{C}}$ but are defined on their suitable (branched) covering spaces. In order to clarify this multivaluedness and to give a clear setting, there may be two approaches.

1. Transcendental method: introduce a topological covering space $T_{W,\text{odd},\mathbb{C}}$ of $T_{W,\mathbb{C}} \setminus B_{W,\text{odd},\mathbb{C}}$ and lift b_W to a univalent holomorphic map $b_{W,\text{odd}}$ defined on it, and similarly for $c_{W,\text{odd}}$.

2. Algebraic method: introduce a suitable finite covering variety $\tilde{T}_W \rightarrow T_W$ (branching along $B_{W,\text{odd}}$) and introduce \tilde{b}_W as a scheme morphism from \tilde{T}_W to the affine space V_Π , and similarly for \tilde{c}_W .

The first approach is naive and easily understandable. However, there is a disadvantage that the “boundary points $B_{W,\text{odd}}$ ” is excluded from the domain of definition. In the second approach those boundary points are naturally included in the domain of definition. Furthermore, it has another advantage that we can discuss about the twisted real forms of the maps (which plays a basic role in our study). For these reasons, we employ the second approach in [S4,§10]. However, the second approach is technically more involved, and we use the first approach to formulate Theorem C in §3.5 in the present paper.

An important role of the maps c_W and b_W is that they identify certain area in S_W and in T_W with certain area in a linear space \widehat{V}_Π and in V_Π , respectively. In particular, the map c_W identifies the twisted real discriminant locus in a tube domain of the source space with a system of real hyperplanes in a tube domain of the target space. This has several fruitful consequences, since *the study of configurations among branches of the real discriminant locus is reduced to a study of a certain system of linear inequalities*. By this reason, we call these maps c_W and b_W the *linearization maps*.

Let us explain the contents of this section.

The linear model spaces \widehat{V}_Π and V_Π , which will be the target spaces of the linearization maps, are described in §3.1. Depending only on the Coxeter graph $\Gamma(W)$, we introduce a simplicial cone $E_{\Gamma(W)}$ in V_Π in §3.2. In §3.3, we introduce the covering space $T_{W,\text{odd},\mathbb{C}}$, on which the two algebroid functions $\varphi_{\alpha,\varepsilon}$ for $\varepsilon \in \{\pm 1\}$ in §2.6 lift to the same globally defined univalent function, denoted by φ_α . By the use of them

in §3.4, the linearization maps c_W and b_W are defined. In §3.5, we formulate Theorem C, which states about the comparison of the real spaces $S_W^{[e]}$ and \widehat{V}_Π obtained by the linearization maps. As applications of Theorem C, Theorems A and B are proved in §3.6 and §3.7. We illustrate in §3.8 the linearization maps for the type A_3 .

3.1. Linear model spaces \widehat{V}_Π and V_Π .

We introduce two linear model spaces \widehat{V}_Π and V_Π , which will be the target spaces of the linearization maps c_W and b_W , respectively.

Define the real vector space with the basis $\{v_\alpha\}_{\alpha \in \Pi}$:

$$(3.1.1) \quad \widehat{V}_\Pi := \bigoplus_{\alpha \in \Pi} \mathbb{R}v_\alpha.$$

Translation by constant multiples of the diagonal element:

$$(3.1.2) \quad v_\Pi := \sum_{\alpha \in \Pi} v_\alpha$$

defines a \mathbb{G}_a -action on \widehat{V}_Π :

$$(3.1.3) \quad (\lambda, \tilde{v}) \in \mathbb{G}_a \times \widehat{V}_\Pi \mapsto \tilde{v} + \lambda \cdot v_\Pi \in \widehat{V}_\Pi.$$

The quotient space V_Π and the quotient map π_Π are introduced by

$$(3.1.4) \quad \pi_\Pi : \widehat{V}_\Pi \longrightarrow V_\Pi := \widehat{V}_\Pi / \mathbb{R} \cdot v_\Pi.$$

The symmetric group $\mathfrak{S}(\Pi)$ acts linearly on \widehat{V}_Π by permuting the basis v_α . Since v_Π is fixed by $\mathfrak{S}(\Pi)$, it induces an action of $\mathfrak{S}(\Pi)$ on V_Π .

Let $\{\lambda_\alpha\}_{\alpha \in \Pi}$ be the dual basis of $\{v_\alpha\}_{\alpha \in \Pi}$, i.e., the coordinate system of \widehat{V}_Π . The infinitesimal action of the \mathbb{G}_a -action (3.1.4) is then given by

$$(3.1.5) \quad \sum_{\alpha \in \Pi} \frac{\partial}{\partial \lambda_\alpha}.$$

Consider the coordinate hyperplane in \widehat{V}_Π :

$$(3.1.6) \quad H_\alpha := \{ \sum_{\beta \in \Pi} \lambda_\beta v_\beta \in \widehat{V}_\Pi \mid \lambda_\alpha = 0 \}$$

for $\alpha \in \Pi$. The projection π_Π induces an isomorphism from H_α to V_Π for each $\alpha \in \Pi$, and also an isomorphism from the intersections $H_\alpha \cap H_\beta$ for $\alpha, \beta \in \Pi$ ($\alpha \neq \beta$) to the hyperplane in V_Π :

$$(3.1.7) \quad H_{\alpha\beta} := \{ v \in V_\Pi \mid \lambda_{\alpha\beta}(v) = 0 \}.$$

Here $\lambda_{\alpha\beta} := \lambda_\alpha - \lambda_\beta$ is a linear form on V_Π , which satisfies

$$(3.1.8) \quad \lambda_{\alpha\beta} + \lambda_{\beta\gamma} = \lambda_{\alpha\gamma} \quad \text{for } \alpha, \beta, \gamma \in \Pi.$$

Note. The set of linear forms $\{\lambda_{\alpha\beta} \mid \alpha, \beta \in \Pi, \alpha \neq \beta\}$ on V_Π forms a root system of type A_{l-1} , where $H_{\alpha\beta}$ are the reflection hyperplanes of the group $\mathfrak{S}(\Pi)$. What is different from the usual setting is the fact that the reflection hyperplane $H_{\alpha\beta}$ is labeled by the positive integer $\text{ord}(\alpha\beta)$.

3.2. $\Gamma(W)$ -cone $E_{\Gamma(W)}$ in V_{Π} .

Depending on the graph $\Gamma(W)$ (see §2.4) and on the partition $\{\Pi_1, \Pi_2\}$ (2.4.1), we introduce an open simplicial cone $E_{\Gamma(W)}$ in V_{Π} . Put

$$(3.2.1) \quad E_{\Gamma(W)} := \text{the connected component of } V_{\Pi} \setminus \{\Omega_{\Gamma(W)} = 0\} \\ \text{containing the half line } \mathbb{R}_{>0}(v_{\Pi_1} - v_{\Pi_2})$$

where $\Omega_{\Gamma(W)} := \prod_{\overline{\alpha\beta} \in \text{Edge}(\Gamma(W))} \lambda_{\alpha\beta}^2$ and $v_{\Pi_i} := \sum_{\alpha \in \Pi_i} v_{\alpha}$ ($i = 1, 2$). We call $E_{\Gamma(W)}$ the $\Gamma(W)$ -cone. As an immediate consequence of the definition, we have:

Assertion 3.1. *The $\Gamma(W)$ -cone is given by inequalities:*

$$(3.2.2) \quad E_{\Gamma(W)} = \left\{ v \in V_{\Pi} \mid \lambda_{\alpha\beta}(v) > 0 \right. \\ \left. \text{for } \alpha \in \Pi_1, \beta \in \Pi_2 \text{ such that } \overline{\alpha\beta} \in \text{Edge}(\Gamma(W)) \right\}.$$

Therefore, $E_{\Gamma(W)}$ is an open simplicial cone.

The transposition of Π_1 and Π_2 induces the change of the $\Gamma(W)$ -cone $E_{\Gamma(W)}$ to $-E_{\Gamma(W)}$. This dependence of the $\Gamma(W)$ -cone on the partition of Π is subtle but important. However, for the sake of simplicity, we omit $\{\Pi_1, \Pi_2\}$ in the notation $E_{\Gamma(W)}$ unless explicitly mentioned.

3.3. Covering spaces $T_{W, \text{odd}, \mathbb{C}}$ and $S_{W, \text{odd}, \mathbb{C}}$.

We introduce a covering space $T_{W, \text{odd}, \mathbb{C}}$ of $T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}}$, where we lift the two base point loci O^{ε} . It turns out that two germs of algebroid functions $\varphi_{\alpha, +1}$ and $\varphi_{\alpha, -1}$, lifted in their neighborhoods, are analytically continued to the same univalent function, denoted by φ_{α} , on $T_{W, \text{odd}, \mathbb{C}}$.

Consider the complexified vertex orbit line $T_{W(I_2(h)), \mathbb{C}}$ (recall §2.4 and §2.5), and let $\gamma^{[\varepsilon]}$ be the generator of $\pi_1(T_{W(I_2(h)), \mathbb{C}} \setminus \{0\}, O^{\varepsilon}) \simeq \mathbb{Z}$ turning once around the origin counter-clockwise. Then one has ([S4, 9.2]):

- i) $(\gamma^{[\varepsilon]})^2$ belongs to the center of $\pi_1(T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}}, O^{\varepsilon})$.
- ii) the monodromy action of $\gamma^{[\varepsilon]}$ on $\{\varphi_{\alpha, \varepsilon}\}_{\alpha \in \Pi}$ (2.6.2) is given by

$$(3.3.1) \quad \varphi_{\alpha, \varepsilon}(\tilde{t} \cdot \gamma^{[\varepsilon]}) = \varphi_{\chi_W(\alpha), \varepsilon}(\tilde{t}),$$

where $\chi_W \in \mathfrak{S}(\Pi)$ is the involution of the set Π obtained by the adjoint action of the longest element of W .

The fundamental group $\pi_1(T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}}, O^{\varepsilon})$ acts on the universal covering space $(T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}})^{\sim}$ by choosing a base point locus \tilde{O}^{ε} (i.e., a closed set in the covering which projects homeomorphically onto O^{ε}). Let us introduce the *central quotient space*


$$(3.3.2) \quad T_{W, \text{odd}, \mathbb{C}} := (T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}})^{\sim} / \langle (\gamma^{[\varepsilon]})^2 \rangle$$

with the base point locus $O_{\text{odd}}^{\varepsilon} :=$ the image of \tilde{O}^{ε} . One has the natural covering map: $\varpi_{\text{odd}} : T_{W, \text{odd}, \mathbb{C}} \rightarrow T_{W, \mathbb{C}} \setminus B_{W, \text{odd}, \mathbb{C}}$ with $\varpi_{\text{odd}} : O_{\text{odd}}^{\varepsilon} \simeq O^{\varepsilon}$.

Remark 9. The $T_{W,odd,\mathbb{C}}$ contains pull-backs of evenly labeled bifurcation set $B_{W,even,\mathbb{C}} \setminus B_{W,odd,\mathbb{C}}$ and, in particular, the ordinary part $B_{W,2,\mathbb{C}}$.

To be precise, we have constructed two covering spaces depending on $\varepsilon \in \{\pm 1\}$. In the following, we identify them and consider only one space, denoted again by $T_{W,odd,\mathbb{C}}$, by choosing the two base point locus O_{odd}^ε ($\varepsilon \in \{\pm 1\}$) simultaneously as follows: The inverse image by ϖ_{odd} of the complexified v.o. line $T_{W(I_2(h)),\mathbb{C}}$ decomposes into the connected components, each of which is a double cover of the complex v.o. line and is isomorphic to \mathbb{C}^\times and admits a natural \mathbb{C}^\times -action. Choose one component and fix the base point locus inside it as follows:

Fig.1. Four base point loci in $T_{W,odd,\mathbb{C}}$



where O_{odd*}^ε are some auxiliary base point loci (see Remark 10).

The germ of an algebroid function $\varphi_{\alpha,\varepsilon}$ (2.6.2) is lifted to a germ of holomorphic function, again denoted by $\varphi_{\alpha,\varepsilon}$, on a neighborhood of O_{odd}^ε . We observe that:

- i) the germ $\varphi_{\alpha,\varepsilon}$ for $\alpha \in \Pi$ and $\varepsilon \in \{\pm 1\}$ is analytically continued to a unique univalent holomorphic function on $T_{W,odd,\mathbb{C}}$,
- ii) for each $\alpha \in \Pi$, the two univalent functions defined in i) for $\varepsilon \in \{\pm 1\}$ define the same function on $T_{W,odd,\mathbb{C}}$. Let us denote it by

$$(3.3.3) \quad \varphi_\alpha \quad \text{for } \alpha \in \Pi.$$

Remark 10. The fact i) is an immediate consequence of (3.3.1). The fact ii) is based on the choices of the base point loci in Fig. 1 and of the sign of ζ_i^ε in the construction of $\varphi_{\alpha,\varepsilon}$ in §2.6. Namely, for $\varepsilon = -1$, if we choose ζ_i^ε to be the complex conjugate $\zeta_i/\sqrt{\varepsilon}$ of $\zeta_i\sqrt{\varepsilon}$, we have to take O_{odd*}^{-1} instead of O_{odd}^{-1} as the base point locus. That is: the generator of the Galois group $\text{Gal}(\mathbb{C}/\mathbb{R}) \simeq \mathbb{Z}/2\mathbb{Z}$ acts on the index set Π of $\{\varphi_{\alpha,-1}\}_{\alpha \in \Pi}$ by the formula (3.3.1) (see [S4,8.4 & (9.3.8)]).

Let us introduce the product space: $S_{W,odd,\mathbb{C}} := S_{W,\mathbb{C}} \times_{T_{W,\mathbb{C}}} T_{W,odd,\mathbb{C}}$. On $S_{W,odd,\mathbb{C}}$, we introduce a system of univalent holomorphic functions:

$$(3.3.4) \quad \phi_\alpha := P_l - \varphi_\alpha \quad \text{for } \alpha \in \Pi.$$

3.4. Linearization morphism c_W on $S_{W,odd,\mathbb{C}}$.

We introduce the linearization map b_W as the map from the covering space $T_{W,odd,\mathbb{C}}$ of $T_{W,\mathbb{C}}$ to the complexified model vector spaces $V_{\Pi,\mathbb{C}}$, and similarly c_W from $S_{W,odd,\mathbb{C}}$ to $\widehat{V}_{\Pi,\mathbb{C}}$.

Definition. Using the functions φ_α (3.3.3) and ϕ_α (3.3.4) as for the coefficients of v_α , we consider the maps:

$$(3.4.1) \quad \begin{aligned} c_{W,odd,\mathbb{C}} &= \omega^{-1} \sum_{\alpha \in \Pi} \phi_\alpha v_\alpha : S_{W,odd,\mathbb{C}} \times \mathbb{C}^\times \longrightarrow \widehat{V}_{\Pi,\mathbb{C}} := \widehat{V}_\Pi \otimes_{\mathbb{R}} \mathbb{C} \\ b_{W,odd,\mathbb{C}} &= -\omega^{-1} \sum_{\alpha \in \Pi} \varphi_\alpha v_\alpha : T_{W,odd,\mathbb{C}} \times \mathbb{C}^\times \longrightarrow V_{\Pi,\mathbb{C}} := V_\Pi \otimes_{\mathbb{R}} \mathbb{C} \end{aligned}$$

and call it the *linearization map*, where $\omega \in \mathbb{C}^\times$ is a scaling factor which shall take a special value depending on the situation.

The push-forward of the primitive vector field D is given by

$$(3.4.2) \quad (c_W)_*(D) = \omega^{-1} \sum_{\alpha \in \Pi} \frac{\partial}{\partial \lambda_\alpha}.$$

namely, $c_{W,\mathbb{C}}$ is equivariant with respect to the two \mathbb{G}_a -actions: the τ -action on S_W and the diagonal translation on \widehat{V}_Π (shifted by ω). One has

$$(3.4.3) \quad c_{W,odd,\mathbb{C}}(\tau(\lambda)z, \omega) = c_{W,odd,\mathbb{C}}(z, \omega) + \omega^{-1} \lambda \cdot v_\Pi,$$

and, hence, the following diagram is commutative:

$$(3.4.4) \quad \begin{array}{ccc} S_{W,odd,\mathbb{C}} \times \mathbb{C}^\times & \xrightarrow{c_{W,odd,\mathbb{C}}} & \widehat{V}_{\Pi,\mathbb{C}} \\ \downarrow (\pi_\tau, \text{id}) & & \downarrow \pi_\Pi \\ T_{W,odd,\mathbb{C}} \times \mathbb{C}^\times & \xrightarrow{b_{W,odd,\mathbb{C}}} & V_{\Pi,\mathbb{C}} \end{array} .$$

The formula (2.6.5) can be reformulated as the Jacobian formula

$$(3.4.5) \quad \begin{aligned} Jac(b_{W,odd,\mathbb{C}}) &= c \cdot \omega^{-l} \prod_{p \geq 3} \omega_{W,p}^{p/2-1}, \\ Jac(c_{W,odd,\mathbb{C}}) &= c \cdot \omega^{1-l} \pi_\tau^* \left(\prod_{p \geq 3} \omega_{W,p}^{p/2-1} \right). \end{aligned}$$

for some constants $c \in \mathbb{R}^\times$. We observe that the factor $\omega_{W,2}$ does not appear in the right hand side. Therefore,

Assertion 3.2. *The maps $c_{W,odd,\mathbb{C}}$ and $b_{W,odd,\mathbb{C}}$ are not ramifying along the ordinary bifurcation set $(\pi_\tau)^{-1}(B_{W,2,\mathbb{C}})$ and $B_{W,2,\mathbb{C}}$, but are ramifying along $(\pi_\tau)^{-1}(B_{W,2p,\mathbb{C}})$ and $B_{W,2p,\mathbb{C}}$ for $p \geq 2$, respectively.*

3.5. Theorem C.

In the previous paragraph, the linearization maps are introduced as holomorphic maps from the complex manifolds $T_{W,odd,\mathbb{C}}$ and $S_{W,odd,\mathbb{C}}$ to the linear model spaces. In this paragraph, we restrict the domain of the definitions of the linearization maps $b_{W,odd,\mathbb{C}}$ and $c_{W,odd,\mathbb{C}}$ to the central region $E_W^{\{\varepsilon\}}$ introduced in (2.5.8) and the tube domain $(\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}})$ over the central region with a fixed scaling parameter

$\omega \in \{\pm\sqrt{\varepsilon^h}\}$ (see Remark 12 below). The linearization maps for a fixed ω are denoted by $b_{W,\omega D}$ and $c_{W,\omega D}$, respectively:

$$b_{W,\omega D}(x) := b_{W,\text{odd},\mathbb{C}}(x, \omega) \quad \text{and} \quad c_{W,\omega D}(x) := c_{W,\text{odd},\mathbb{C}}(x, \omega).$$

It is a straightforward calculation that for these choices, the image of the linearization maps are contained in the real forms V_Π and \widehat{V}_Π (see Remarks 11). Thus, (3.4.4) gives rise to the following commutative diagram of semi-algebraic maps (see Remarks 12):

$$(3.5.1) \quad \begin{array}{ccc} (\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}}) & \xrightarrow{c_{W,\omega D}} & \widehat{V}_\Pi \\ \downarrow \pi_\tau^{[\varepsilon]} & & \downarrow \pi_\Pi \\ E_W^{\{\varepsilon\}} & \xrightarrow{b_{W,\omega D}} & V_\Pi \end{array}$$

Remark 11. Generally, we have the following result on the real form of linearization map ([S4]). *For any twisted real structure $T_{W,\mathbb{R}}^{[u]}$ and for any connected component E of $T_{W,\mathbb{R}}^{[u]} \setminus B_{W,\text{odd},\mathbb{R}}^{[u]}$, there exists an involution $\chi \in \mathfrak{S}(\Pi)$ such that the linearization map $b_{W,\omega D}$ induces a (real multivalued) map from E to the twisted real space V_Π^χ , where the scaling constant ω is chosen in the twisted real form: $\mathbb{C}^{\times, b[u]} = \mathbb{R}_{>0} \sqrt{b[u]} \sqcup (-\mathbb{R}_{>0} \sqrt{b[u]})$ with respect to the sign $b[u] \in \{\pm 1\}$ introduced in Assertion 2.1 in §2.3.*

Actually, in the present paper, we take $[u] = [\varepsilon]$, $E = E_W^{\{\varepsilon\}}$, $b[\varepsilon] = \varepsilon^h$, and $\chi = \text{id} \in \mathfrak{S}(\Pi)$.

Remark 12. So far in the present paper, the linearization maps are defined on the covering spaces $T_{W,\text{odd},\mathbb{C}}$ and $S_{W,\text{odd},\mathbb{C}}$. Therefore, one should have, first, introduced the map $b_{W,\omega D}$ on a certain covering space $\tilde{E}_W^{\{\varepsilon\}}$ embedded in $T_{W,\text{odd},\mathbb{C}}$, namely on the connected component of the inverse image of $E_W^{\{\varepsilon\}}$ in $T_{W,\text{odd},\mathbb{C}}$ which contains the base point locus $O_{\text{odd}}^\varepsilon$, and similarly for $c_{W,\omega D}$, and then formulate Theorem C in terms of the mapped defined on the covering spaces. Actually, as a consequence of Theorem C, $E_W^{\{\varepsilon\}}$ become homeomorphic to a simplicial cone in a real vector space so that it is simply connected. Also, $(\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}})$, as a tube domain over $E_W^{\{\varepsilon\}}$, is simply connected. Thus the covering spaces reduce to trivial covering spaces, and $b_{W,\omega D}$ and $c_{W,\omega D}$ are well defined as univalent maps on $E_W^{\{\varepsilon\}}$ and $(\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}})$. Therefore, in the formulation of Theorem C in the present paper, we assume the knowledge of the simply connectedness beforehand.

We state Theorem C, announced in the introduction.

Theorem C. *Depending on the choices of the sign $\varepsilon \in \{\pm 1\}$, the scaling factor $\omega \in \{\pm \sqrt{\varepsilon^h}\}$ and the partition $\{\Pi_1, \Pi_2\}$, take a sign factor*

$$(3.5.2) \quad \sigma := -\omega \sigma(D, \{\Pi_1, \Pi_2\})(\sqrt{\varepsilon})^h \in \{\pm 1\}$$

(recall (2.5.4) for $\sigma(D, \{\Pi_1, \Pi_2\})$). Then the following (1)–(7) hold.

(1) *The linearization map $b_{W, \sigma \cdot \omega D}^{[\varepsilon]}$ induces a semi-algebraic homeomorphism:*

$$(3.5.3) \quad b_{W, \sigma \cdot \omega D}^{[\varepsilon]} : E_W^{\{\varepsilon\}} \xrightarrow{\simeq} E_{\Gamma(W)},$$

which extends to their closures homeomorphically.

(2) *The linearization map $c_{W, \omega D}^{[\varepsilon]}$ induces a semi-algebraic homeomorphism:*

$$(3.5.4) \quad c_{W, \sigma \cdot \omega D}^{[\varepsilon]} : (\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}}) \xrightarrow{\simeq} (\pi_\Pi)^{-1}(E_{\Gamma(W)}),$$

which extends to their closures homeomorphically.

(3) *The linearization map (3.5.4) is \mathbb{G}_a -equivariant so that we obtain the commutative Cartesian diagram:*

$$(3.5.5) \quad \begin{array}{ccc} (\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}}) & \xrightarrow[c_{W, \sigma \cdot \omega D}^{[\varepsilon]}]{\simeq} & (\pi_\Pi)^{-1}(E_{\Gamma(W)}) \\ \pi_\tau^{[\varepsilon]} \downarrow & & \pi_\Pi \downarrow \\ E_W^{\{\varepsilon\}} & \xrightarrow[b_{W, \sigma \cdot \omega D}]{\simeq} & E_{\Gamma(W)} \end{array}$$

(4) *The linearization map $b_{W, \sigma \cdot \omega D}^{[\varepsilon]}$ maps the ordinary bifurcation set $B_{W, 2, \mathbb{R}}^{[\varepsilon]}$ to the union of the 2-labeled reflection hyperplanes in V_Π :*

$$(3.5.6) \quad b_{W, \sigma \cdot \omega D}^{[\varepsilon]} : E_W^{\{\varepsilon\}} \cap B_{W, 2, \mathbb{R}} \xrightarrow{\simeq} E_{\Gamma(W)} \cap (\cup_{\alpha\beta \notin \text{Edge}(\Gamma(W))} H_{\alpha\beta}),$$

(5) *The linearization map $c_{W, \sigma \cdot \omega D}^{[\varepsilon]}$ maps $D_{W, \mathbb{R}}^{[\varepsilon]}$ to the union of the hyperplanes in \widehat{V}_Π :*

$$(3.5.7) \quad c_{W, \sigma \cdot \omega D}^{[\varepsilon]} : (\pi_\tau^{[\varepsilon]})^{-1}(\overline{E_W^{\{\varepsilon\}}}) \cap D_W \xrightarrow{\simeq} (\pi_\Pi)^{-1}(E_{\Gamma(W)}) \cap (\cup_{\alpha \in \Pi} H_\alpha),$$

(6) *The linearization map $c_{W, \omega D}^{[\varepsilon]}$ maps the central component $\mathcal{C}^{\{\varepsilon\}}$ $S_{W, \mathbb{R}}^{[\varepsilon]}$ to the coordinate hyperquadrant in \widehat{V}_Π :*

$$(3.5.8) \quad \widehat{\mathcal{C}}_{\Gamma(W)} := \{\sum_{\alpha \in \Pi} \lambda_\alpha v_\alpha \in \widehat{V}_\Pi \mid (-1)^i \lambda_\alpha > 0 \text{ for } \alpha \in \Pi_i, i = 1, 2\}$$

homeomorphically:

$$(3.5.9) \quad c_{W, \sigma \cdot \omega D}^{[\varepsilon]} : \mathcal{C}^{\{\varepsilon\}} \xrightarrow{\simeq} \widehat{\mathcal{C}}_{\Gamma(W)}.$$

The map extends to their closures homeomorphically. The hyper-quadrant satisfies $\pi_\Pi(\widehat{\mathcal{C}}_{\Gamma(W)}) = E_{\Gamma(W)}$.

(7) The linearization map $c_{W,\omega D}^{[\varepsilon]}$ maps the opposite components $\mathcal{C}_{\pm}^{[\varepsilon]}$ in $S_{W,\mathbb{R}}^{[\varepsilon]}$ (cf. (1.7.1) and (2.5.5)) to the hyper-quadrants in \widehat{V}_{Π} :

$$(3.5.10) \quad \widehat{\mathcal{C}}_{\Gamma(W),\pm} := \left\{ \sum_{\alpha \in \Pi} \lambda_{\alpha} v_{\alpha} \in \widehat{V}_{\Pi} \mid \begin{array}{l} (-1)^{i-1} \lambda_{\alpha} > 0 \text{ for } i \text{ with} \\ (-1)^{i-1} = \pm 1 \text{ and } \alpha \in \Pi_i \end{array} \right\},$$

homeomorphically:

$$(3.5.11) \quad c_{W,\sigma \cdot \omega D}^{[\varepsilon]} : \mathcal{C}_{\pm}^{[\varepsilon]} \cap (\pi_{\tau}^{[\varepsilon]})^{-1} E_W^{\{\varepsilon\}} \xrightarrow{\simeq} \widehat{\mathcal{C}}_{\Gamma(W),\pm} \cap (\pi_{\Pi})^{-1} E_{\Gamma(W)}.$$

The map extends to their closures homeomorphically.

(8) For a subset Σ of Π , let F_{Σ} be the facet of $\mathcal{C}^{\{\varepsilon\}}$ corresponding to the facet $\cap_{\alpha \in \Sigma} \{\lambda_{\alpha} = 0\} \cap \widehat{\mathcal{C}}_{\Gamma(W)}$ by (3.5.9). Then ϕ_{β} for $\beta \in \Sigma^c := \Pi \setminus \Sigma$ is regular on a neighborhood of F_{Σ} in $S_{W,\mathbb{R}}^{\{\varepsilon\}}$, and one obtains a semi-algebraic isomorphism:

$$(3.5.12) \quad (\sigma \omega \cdot \phi_{\beta})_{\beta \in \Sigma^c} : F_{\Sigma} \xrightarrow{\simeq} \left(\prod_{\beta \in \Pi_1 \cap \Sigma^c} \mathbb{R}_{>0} \right) \times \left(\prod_{\beta \in \Pi_2 \cap \Sigma^c} \mathbb{R}_{<0} \right).$$

Remark 13. In Theorem C, the scaling factor ω and the sign factor σ appear always as the product $\sigma \cdot \omega$. In this paper we distinguished them because of their different origins.

3.6. Proof of Theorem A.

Recall the \mathbb{R} -equivariant isomorphism (3.5.5). Using (3.5.8)–(3.5.11), for a positive real number $\lambda \in \mathbb{R}_{>0}$, the map $c_{W,\sigma \cdot \omega D}^{[\varepsilon]}$ induces a semi-algebraic diffeomorphism from $\bar{J}_W^{\{\varepsilon\}}(\lambda) := \bar{\mathcal{C}}^{\{\varepsilon\}} \cap \tau^{[\varepsilon]}(-\lambda) \bar{\mathcal{C}}_+^{[\varepsilon]} \cap \tau^{[\varepsilon]}(\lambda) \bar{\mathcal{C}}_-^{[\varepsilon]}$ to

$$(3.6.1) \quad \begin{aligned} & \widehat{\mathcal{C}}_{\Gamma(W)} \cap (\widehat{\mathcal{C}}_{\Gamma(W),+} - \lambda v_{\Pi}) \cap (\widehat{\mathcal{C}}_{\Gamma(W),-} + \lambda v_{\Pi}) \\ & = \{(\lambda_{\alpha})_{\alpha \in \Pi} \in \widehat{V}_{\Pi} \mid 0 \leq (-1)^i \lambda_{\alpha} \leq \lambda \text{ for } \alpha \in \Pi_i, i = 1, 2\} \end{aligned}$$

where the right hand side is a parallelotope of dimension l in \widehat{V}_{Π} . It is the intersection of two simplicial cones $\widehat{\mathcal{C}}_{\Gamma(W)}^{\varepsilon}$ and $\widehat{\mathcal{C}}_{\Gamma(W)}^{\varepsilon}(\lambda)$, where

$$(3.6.2) \quad \begin{aligned} \widehat{\mathcal{C}}_{\Gamma(W)}^{\varepsilon}(\lambda) & := (\widehat{\mathcal{C}}_{\Gamma(W),+} - \lambda v_{\Pi}) \cap (\widehat{\mathcal{C}}_{\Gamma(W),-} + \lambda v_{\Pi}) \\ & = \{(\lambda_{\alpha})_{\alpha \in \Pi} \in \widehat{V}_{\Pi} \mid (-1)^i \lambda_{\alpha} \leq \lambda \text{ for } \alpha \in \Pi_i, i = 1, 2\}. \end{aligned}$$

Let us show a slightly stronger transversality between the two cones in order to apply it to the proof of Theorem B in the next subsection.

Theorem A addendum. *The faces of $\bar{J}_W^{\{\varepsilon\}}(\lambda)$ are crossing normally at any point of $\bar{J}_W^{\{\varepsilon\}}(\lambda) \setminus (\pi_{\tau}^{[\varepsilon]})^{-1}(B_{W,\geq 3,\mathbb{R}}^{[\varepsilon]})$ (recall §1.3 Definition 1).*

Proof. Recall the formula (3.4.5) for the Jacobian $\frac{\partial(\phi_{\alpha_1}, \dots, \phi_{\alpha_l})}{\partial(P_1, \dots, P_l)}$ of the map c_W . The right hand expression means that it does not vanish on the complement of $(\pi_\tau)^{-1}(B_{W, \geq 3, \mathbb{R}}^{[\varepsilon]})$. This means that the hyperplanes $\phi_\alpha = \text{const}$ for $\alpha \in \Pi$, which define faces of the polyhedra $J_W^{\{\varepsilon\}}(\lambda)$, are normally crossing on the complement of $(\pi_\tau)^{-1}(B_{W, \geq 3, \mathbb{R}}^{[\varepsilon]})$. \square

This addendum proves the transversality stated in Theorem A and hence completes the proof of Theorem A.

Remark 14. Let $ao^{\{\varepsilon\}}(\lambda)$ be the vertex of $\bar{J}_W^{\{\varepsilon\}}(\lambda)$ which is antipodal to the origin. By the definition, it is on the axis $\{S=0\} \subset S_{W(I_2(h)), \mathbb{R}}$. For each $\varepsilon \in \{\pm 1\}$, one has $AO^\varepsilon = \{ao^{\{\varepsilon\}}(\lambda) \mid \lambda \in \mathbb{R}_{>0}\}$. This is the reason why AO^ε introduced in (2.5.6), is called the vertex orbit axis.

3.7. Proof of Theorem B.

We shall show that

$$(3.7.1) \quad \bar{K}_W^\varepsilon(\lambda) := (\pi_W)^{-1}(\bar{J}_W^{\{\varepsilon\}}(\lambda))$$

is a semi-algebraic polyhedron dual to the Weyl chamber decomposition of $V_{\mathbb{R}}^\varepsilon$. Since the proofs for $\varepsilon=1$ and for $\varepsilon=-1$ are completely parallel, we prove only the case $\varepsilon=1$ and omit the upperscripts ε , $\{\varepsilon\}$ and $[\varepsilon]$. The proof is divided into two parts: 1. local analytic part and 2. global combinatorial part.

1. We study the local analytic property of $\bar{K}_W(\lambda)$. In this paragraph, we mean by $(X, x) \simeq (Y, y)$ that there exists an isomorphism from a neighborhood of x in X to a neighborhood of y in Y bringing x to y .

Assertion 3.3. *Let $\tilde{x} \in \bar{K}_W(\lambda)$ and $m := \dim_{\mathbb{R}} V_{\mathbb{R}}^{W(\tilde{x})}$, where $W(\tilde{x})$ is the stabilizer subgroup of W at \tilde{x} and $V_{\mathbb{R}}^{W(\tilde{x})}$ is the fixed point subspace by the $W(\tilde{x})$ -action. Then, there exist an integer k with $0 \leq k \leq m$ and a local real analytic isomorphism from a neighborhood of \tilde{x} in $V_{\mathbb{R}}$ to a neighborhood of the origin of $(V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}) \times \mathbb{R}^m$ which makes the following diagram commutative:*

$$(3.7.2) \quad \begin{array}{ccc} (V_{\mathbb{R}}, \tilde{x}) & \simeq & ((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}) \times \mathbb{R}^m, 0) \\ \cup & & \cup \\ (\bar{K}_W(\lambda), \tilde{x}) & \simeq & ((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}) \times \mathbb{R}_{\geq 0}^k \times \mathbb{R}^{m-k}, 0), \end{array}$$

Furthermore, the isomorphism induces the following isomorphisms:

i) the isomorphism of the subspaces

$$(3.7.3) \quad (V_{\mathbb{R}}^{W(\tilde{x})}, x) \simeq (\{0\} \times \mathbb{R}^m, 0)$$

ii) for any facet G of $\bar{J}_W(\lambda)$ which is adjacent to $ao(\lambda)$

$$(3.7.4) \quad (\pi_W^{-1}(\bar{G}), \tilde{x}) \simeq ((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}) \times \bar{K}, 0)$$

where \bar{K} is the closure of a facet K of $(\mathbb{R}_{\geq 0}^k \times \mathbb{R}^{m-k}, 0)$ (which may be empty).

Proof. Let Q_1, \dots, Q_l be a system of generators of the ring of invariants $S(V_{\mathbb{R}}^*)^{W(\tilde{x})}$, and let us consider a $W(\tilde{x})$ -invariant map $\bar{Q} = (Q_1, \dots, Q_l) : (V_{\mathbb{R}}, \tilde{x}) \rightarrow (\mathbb{R}^l, \bar{x})$ for $\bar{x} := \bar{Q}(\tilde{x})$. Then, there is a local analytic isomorphism $\varpi : (\mathbb{R}^l, \bar{x}) \simeq (S_{W, \mathbb{R}}, x)$ for $x := \pi_W(\tilde{x})$ such that $\pi_W = \varpi \circ \bar{Q}$. We may choose the first Q_1, \dots, Q_{l-m} which form the generators of the ring of invariant polynomials on $V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}$ by the $W(\tilde{x})$ -action, and the last Q_{l-m+1}, \dots, Q_l which are $W(\tilde{x})$ -invariant linear functions on $V_{\mathbb{R}}$ whose restrictions on $V_{\mathbb{R}}^{W(\tilde{x})}$ give its linear coordinate system such that $\bar{x} = 0$. By this choice of the Q_i 's, the local analytic isomorphism ϖ induces a local splitting of the set \mathcal{C} at x :

$$(3.7.5) \quad (\mathcal{C}, x) \simeq \left(((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})})/W(\tilde{x})) \times \mathbb{R}^m, 0 \right).$$

We shall denote by F the stratum of \mathcal{C} containing x in the left hand side, which is locally the image of $(V_{\mathbb{R}}^{W(\tilde{x})}, \tilde{x})$ by π_W . Then (F, \tilde{x}) is mapped to the subspace $(\mathbb{R}^m, 0)$ in the right hand side.

On the other hand, the linearization map c_W maps the central region \mathcal{C} to the cone $\widehat{\mathcal{C}}_{\Gamma(W)}$ (3.5.8), and hence the stratum F to a stratum of $\widehat{\mathcal{C}}_{\Gamma(W)}$, which is an open cone in $\cap_{i=m+1}^l H_{\alpha_i}$ for some $\{\alpha_{m+1}, \dots, \alpha_l\} \subset \Pi$. Then, $\phi_{\alpha_1}, \dots, \phi_{\alpha_m}$ for the remaining index set $\{\alpha_1, \dots, \alpha_m\} = \Pi \setminus \{\alpha_1, \dots, \alpha_{l-m}\}$ form a local coordinate system of F at x (Theorem C (8)). Therefore, replacing Q_{l-m+1}, \dots, Q_l with $\phi_{\alpha_1} - \phi_{\alpha_1}(x), \dots, \phi_{\alpha_m} - \phi_{\alpha_m}(x)$, we obtain a local analytic expression similar to (3.7.5), where the subspace $(\{0\} \times \mathbb{R}^m, 0)$ of the right hand side is still the image of F by $(\phi_{\alpha_i} - \phi_{\alpha_i}(x))$.

The parallelotope \bar{J}_W , locally at x , is defined as the subset of the central component \mathcal{C} given by inequalities $\pm \phi_{\alpha_i} \leq \lambda$ for some $i \in \{1, \dots, m\}$ and suitable signs (depending on i , recall (3.6.1)). Then after a suitable renumbering of $\{1, \dots, m\}$, we obtain further a local isomorphism:

$$(3.7.6) \quad (\bar{J}_W(\lambda), x) \simeq \left(((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})})/W(\tilde{x})) \times \mathbb{R}_{\geq 0}^k \times \mathbb{R}^{m-k}, 0 \right).$$

The facet decomposition of $\bar{J}_W(\lambda)$ as a parallelotope at x coincides with the natural facet decomposition of $\mathbb{R}_{\geq 0}^k \times \mathbb{R}^{m-k}$ in the right hand side.

Taking the inverse images of the both sides of (3.7.6) in their covering spaces, i.e., a neighborhood of \tilde{x} in $V_{\mathbb{R}}$ and a neighborhood of the origin in $(V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}) \times \mathbb{R}^m$, respectively, we obtain the local analytic isomorphism (3.7.2). Then (3.7.3) follows from the construction.

Let us consider a facet G of \bar{J}_W which is adjacent to $ao(\lambda)$. Since $ao(\lambda) \in \bar{G}$, G is contained in the interior of \mathcal{C} . This implies the image of G in the right hand side of (3.7.5) is contained in $\left(((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})})/W(\tilde{x}))^{\circ} \times \mathbb{R}^m, 0 \right)$ where $((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})})/W(\tilde{x}))^{\circ}$ is the unique open facet. Then, by the isomorphism (3.7.6), the closure \bar{G} of the stratum G is mapped to $\left(((V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})})/W(\tilde{x})) \times \bar{K}, 0 \right)$ for the closure of a suitable facet K of $(\mathbb{R}_{\geq 0}^k \times \mathbb{R}^{m-k}, 0)$ (including empty case). By taking their inverse images, the isomorphism in the first line of (3.7.2) induces (3.7.4). \square

Corollary. *Let G be a facet of $\bar{J}_W(\lambda)$ which is adjacent to $ao(\lambda)$. Then $(\pi_W)^{-1}(\bar{G})$ is a submanifold with corners in $V_{\mathbb{R}}$, which is transversal to the system of hyperplanes $\{H_{\alpha, \mathbb{R}}\}_{\alpha \in R(W)}$.*

Proof. Since $\pi_W|_{(V_{\mathbb{R}} \setminus \cup_{\alpha \in R(W)} H_{\alpha, \mathbb{R}})}$ is locally biregular and $\bar{G} \setminus D_{W, \mathbb{R}}$ is a manifold with corner due to Theorem A addendum in §3.6, we only have to show the property of $(\pi_W)^{-1}(\bar{G})$ at a point $\tilde{x} \in (\cup_{\alpha \in R(W)} H_{\alpha, \mathbb{R}}) \cap (\pi_W)^{-1}(\bar{G})$. Apply Assertion 3.3 at the point \tilde{x} .

The fact ii) in Assertion 3.3 implies that $(\pi_W)^{-1}(\bar{G})$ is a locally closed manifold with corners. Furthermore, the fact that $((\pi_W)^{-1}(\bar{G}), \tilde{x})$ contains the factor $(V_{\mathbb{R}}/V_{\mathbb{R}}^{W(\tilde{x})}, 0)$ implies that it is transversal to the submanifold $(\mathbb{R}^m, 0)$. Since $V_{\mathbb{R}}^{W(\tilde{x})}$ is the intersection of the reflection hyperplanes pathing through \tilde{x} , i) in Assertion 3.3 implies that $((\pi_W)^{-1}(\bar{G}), \tilde{x})$ is transversal to every facet V_{γ} of $V_{\mathbb{R}}$ (recall (1.2.2)). \square

2. We describe the facet decomposition of $\bar{K}_W(\lambda) := (\pi_W)^{-1}(\bar{J}_W(\lambda))$. We first prepare terminology on the facet decomposition of \bar{J}_W .

Let $\mathcal{F}(o) = \{F_{\bar{\gamma}}\}_{\bar{\gamma} \in \Gamma(o)}$ and $\mathcal{F}(ao) = \{G_{\bar{\delta}}\}_{\bar{\delta} \in \Gamma(o)}$ be the sets of facets of \bar{J}_W which are adjacent to o and to ao , respectively. Here we use the same index set $\Gamma(o)$ for the two sets by the reason i) below, and put an overline on the index by the reason iv) below.

i) There is a one-to-one correspondence $\mathcal{F}(o) \leftrightarrow \mathcal{F}(ao)$ in such a manner that $F_{\bar{\gamma}} \leftrightarrow G_{\bar{\delta}}$ if and only if $\bar{F}_{\bar{\gamma}} \cap \bar{G}_{\bar{\delta}}$ consists of a single point.

ii) The set $\Gamma(o)$ is partially ordered such that for $\bar{\gamma}, \bar{\delta} \in \Gamma(o)$ one has

$$(3.7.7) \quad \bar{\gamma} \leq \bar{\delta} \Leftrightarrow F_{\bar{\gamma}} \subset \bar{F}_{\bar{\delta}} \Leftrightarrow \bar{G}_{\bar{\gamma}} \supset G_{\bar{\delta}}.$$

iii) $\bar{G}_{\bar{\gamma}} \cap \bar{F}_{\bar{\delta}} \neq \emptyset$ if and only if $\bar{\gamma} \leq \bar{\delta}$ for $\bar{\gamma}, \bar{\delta} \in \Gamma(o)$. The intersection is a closed facet of \bar{J}_W ($\simeq [0, 1]^k$) of dimension $k = \dim(F_{\bar{\delta}}) - \dim(F_{\bar{\gamma}})$.

iv) Recall the index set Γ (1.2.2), on which W acts in the obvious manner. Then, there is a bijection $\Gamma/W \simeq \Gamma(o)$ (we denote by $\bar{\delta} \in \Gamma(o)$ the image of $\delta \in \Gamma$) such that $\pi_W(\bar{V}_{\delta}) \cap \bar{J}_W(\lambda) = \bar{F}_{\bar{\delta}}$ for $\delta \in \Gamma$.

Definition. A semi-algebraic set K in $V_{\mathbb{R}}$ is called a *facet of $\bar{K}_W(\lambda)$* if there is $G_{\bar{\gamma}} \in \mathcal{F}(ao)$ such that K is the interior of a connected component of $(\pi_W)^{-1}(\bar{G}_{\bar{\gamma}})$.

Let us show that the set of all facets of $\bar{K}_W(\lambda)$ is indexed by Γ . For any $\gamma \in \Gamma$, by the definition, $\pi_W(V_{\gamma})$ and $\bar{G}_{\bar{\gamma}}$ intersects at a single point transversally. Therefore, there exists a unique connected component of $(\pi_W)^{-1}(\bar{G}_{\bar{\gamma}})$ that intersects V_{γ} . Its interior is, by the definition, a facet of $\bar{K}_W(\lambda)$, which we shall denote by K_{γ} . Conversely, let K be the interior of a connected component of $(\pi_W)^{-1}(\bar{G}_{\bar{\gamma}})$ for some $\bar{\gamma} \in \Gamma(o)$. Since $\bar{F}_{\bar{\gamma}}$ intersects $\bar{G}_{\bar{\gamma}}$ transversally at a point, there exists a connected component of $(\pi_W)^{-1}(\bar{F}_{\bar{\gamma}})$ whose closure intersects K at a single point. Let V_{γ} for $\gamma \in \Gamma$ be the cone in (1.2.2) which support the component such that γ projects to $\bar{\gamma}$. Thus, we find a unique $\gamma \in \Gamma$ such that V_{γ} intersect K at a point.

By definition, all facets of $\bar{K}_W(\lambda)$ are disjoint and cover $\bar{K}_W(\lambda)$. This shows the decomposition

$$(3.7.8) \quad \bar{K}_W(\lambda) = \sqcup_{\gamma \in \Gamma} K_{\gamma}.$$

The fact that (3.7.8) gives a semi-algebraic stratification of $\bar{K}_W(\lambda)$ (i.e., they satisfy the boundary condition) can be reduced to that of $\mathcal{F}(ao)$. The fact that $\bar{K}_W(\lambda)$ becomes a semi-algebraic polyhedron with respect to (3.7.8) whose faces are normally crossing follows from (3.7.2).

Finally, we have to show the three duality properties i), ii) and iii) for $\bar{K}_W(\lambda)$ in Definition 2 in §1.3. The proofs are reduced to the duality between $\mathcal{F}(o)$ and $\mathcal{F}(ao)$ and to the local analysis discussed in the first half of this proof. In particular, for the last iii), we have to show that the intersection of the closure of a chamber C with the $\bar{K}_W(\lambda)$

$$(3.7.9) \quad \bar{C} \cap \bar{K}_W(\lambda)$$

is analytically isomorphic to the cube $[0, 1]^l$ in the following strong sense: there is a neighborhood in $V_{\mathbb{R}}$ of the set (3.7.9) and a real analytic isomorphism of the neighborhood to an open subset of \mathbb{R}^l such that the set (3.7.9) is mapped homeomorphically onto the cube $[0, 1]^l$. The fact that the set (3.7.9) is homeomorphic to a cube follows from the fact that the restriction of π_W on the set (3.7.9) is a homeomorphism onto the polyhedron $\bar{J}_W(\lambda)$. On the other hand, we have shown in Assertion 3.3 that faces are normally crossing at any point of the boundary of the set (3.7.9). These two show the required result.

These complete the proof of Theorem B.


Note. The set (3.7.9) is given by two systems of l inequalities on $V_{\mathbb{R}}$.

$$(3.7.10) \quad (\cap_{\alpha \in \Pi} \{f_{\alpha} \geq 0\}) \cap (\cap_{i=1,2} \cap_{\alpha \in \Pi_i} \{(-1)^i \phi_{\alpha} \leq \lambda\}).$$

3.8. Examples of type A_3 .

Fig. 2. The linearization maps of type A_3

The shaded area in the upper and lower right figures are the central region $E_{A_3}^{[\varepsilon]}$ and the $\Gamma(A_3)$ -cone $E_{\Gamma(A_3)}$, respectively. The tube domain $(\pi_\tau^{[\varepsilon]})^{-1}(E_{A_3}^{[\varepsilon]})$ in $S_{A_3, \mathbb{R}}^{[\varepsilon]}$ is illustrated in the upper left figure as the domain sandwiched by the covers of an open booklet. The tube domain $\pi_\Pi^{-1}(E_{\Gamma(A_3)})$ in \widehat{V}_Π is illustrated in the lower left figure as the domain sandwiched by the straight covers of an open booklet.



4. FUNDAMENTAL GROUP OF $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$

A presentation of the fundamental group of the regular orbit space $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ in terms of the Coxeter matrix M_W is given by E. Brieskorn [Br1]. A group G with that presentation, that is: generator system a_α ($\alpha \in \Pi$) and the braid relations: $\underbrace{a_\alpha a_\beta \dots}_{m_{\alpha\beta}\text{-letters}} = \underbrace{a_\beta a_\alpha \dots}_{m_{\alpha\beta}\text{-letters}}$ for $\alpha, \beta \in \Pi$

as for the fundamental relations, is called an Artin group [BS]. The generators of $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}})$ are described in [Br1] in terms of adjacent chambers in $V_{\mathbb{R}}$ (see also a work by P. Deligne [D1], where the Artin group is given in terms of the galleries of chambers). Then, several authors including Brieskorn, Deligne and the author asked to describe the generator system of the Artin group in terms of a geometry on $S_{W,\mathbb{C}}$.

As a consequence of Theorems A, B and C, we give two different answers to this question. The first one is to use the 1-skeleton of $J_W(\lambda)^{\{\varepsilon\}}$ and is described in 4.1. Identification with Brieskorn's generator system is given in 4.2. The second one is to use τ -orbits as the pencil of Zariski-van Kampen method. It is described in §4.3 and is identified in §4.4 with the one given in §4.1. The generator systems, we have described, depend on the choice of $\varepsilon \in \{\pm 1\}$ since the base points belong to the different central component $\mathcal{C}^{\{\varepsilon\}}$. Their relation is given in 4.5.

In this section, we sometimes identify a path and its homotopy class.


4.1. **1-skeleton of the polyhedron** $J_W^{\{\varepsilon\}}(\lambda)$.

With a use of the 1-skeleton of the polyhedron $J_W^{\{\varepsilon\}}(\lambda)$ (Theorem A), we construct a generator system of $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}})$. Using Theorem B, they are identified in §4.2 with the one studied by Brieskorn [Br].

Let $ao^{\{\varepsilon\}}(\lambda)$ be the vertex of $J_W^{\{\varepsilon\}}(\lambda)$ which is antipodal to the origin (belonging to AO^ε). Due to Assertion §1.1 5, for each $\alpha \in \Pi$, there exists a unique edge $[ao^{\{\varepsilon\}}(\lambda), p_\alpha]$ of $J_W^{\{\varepsilon\}}(\lambda)$ which starts from $ao^{\{\varepsilon\}}(\lambda)$ and terminates at a point p_α on the α th face of $\bar{\mathcal{C}}^{\{\varepsilon\}}$ (cf. Fig. 4).

Since the edge $[ao^{\{\varepsilon\}}(\lambda), p_\alpha]$ intersects the discriminant $D_{W,\mathbb{R}}^{[\varepsilon]}$ transversally at p_α (Theorem A), a complexification of $[ao^{\{\varepsilon\}}(\lambda), p_\alpha]$ (a complex open curve in $S_{W,\mathbb{C}}$ which contains $[ao^{\{\varepsilon\}}(\lambda), p_\alpha]$) intersects the discriminant locus $D_{W,\mathbb{C}}$ transversally at p_α . In the complexification, let us consider a closed path γ_α based at $ao^{\{\varepsilon\}}(\lambda)$ and turning once around the discriminant locus at p_α counter-clockwise as in Fig. 3.

Fig. 3. The generator γ_α on an edge of $J_W^{\{\varepsilon\}}(\lambda)$ (cf. Fig. 4).



Theorem 4.1. *The system of the homotopy classes of γ_α ($\alpha \in \Pi$) in $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^{\{\varepsilon\}}(\lambda))$ coincides with the generator system $\{g_\alpha\}_{\alpha \in \Pi}$ given by Brieskorn [Br1, Zusatz]. Therefore, $\pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^{\{\varepsilon\}}(\lambda))$ is an Artin group with respect to the generator system γ_α ($\alpha \in \Pi$).*



4.2. Proof of Theorem 4.1.

Let $\tilde{a}o^\varepsilon(\lambda)$ be the vertex of $K_W^\varepsilon(\lambda)$ in the chamber C^ε which projects to $ao^\varepsilon(\lambda)$. For $\alpha \in \Pi$, let $\alpha \cdot \tilde{a}o^\varepsilon(\lambda)$ be the image of $\tilde{a}o^\varepsilon(\lambda)$ by the reflection α , and $[\tilde{a}o^\varepsilon(\lambda), \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$ be the 1-edge of the polyhedron $K_W^\varepsilon(\lambda)$ connecting two vertices $\tilde{a}o^\varepsilon(\lambda)$ and $\alpha \cdot \tilde{a}o^\varepsilon(\lambda)$, which intersects $H_{\alpha,\mathbb{R}}^\varepsilon$ transversally at an inverse image \tilde{p}_α of the point p_α . Then, π_W projects $[\tilde{a}o^\varepsilon(\lambda), \tilde{p}_\alpha]$ and $[\tilde{p}_\alpha, \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$ onto the edge $[p_\alpha, ao^\varepsilon(\lambda)]$.

The inverse image of the path γ_α (see Fig. 3), which starts at $\tilde{a}o^\varepsilon(\lambda)$, is a path in the complexification of $[\tilde{a}o^\varepsilon(\lambda), \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$ connecting $\tilde{a}o^\varepsilon(\lambda)$ and $\alpha \cdot \tilde{a}o^\varepsilon(\lambda)$ described as follows: start at $\tilde{a}o^\varepsilon(\lambda)$ and move along $[\tilde{a}o^\varepsilon(\lambda), \tilde{p}_\alpha]$ close to \tilde{p}_α . Then, just before reaching \tilde{p}_α turn along a half circle centered at \tilde{p}_α in the counter-clockwise direction (inside a complexification of $[\tilde{a}o^\varepsilon(\lambda), \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$, in which $[\tilde{a}o^\varepsilon(\lambda), \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$ crosses the discriminant locus at the point \tilde{p}_α) and then to come back to a point $[\tilde{p}_\alpha, \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$. Then, again move along $[\tilde{p}_\alpha, \alpha \cdot \tilde{a}o^\varepsilon(\lambda)]$ till the point $\alpha \cdot \tilde{a}o^\varepsilon(\lambda)$. In fact, this path is homotopic to the path g_α described by Brieskorn [Br1,Zusatz].

Note. Let us give briefly explain how the braid relations follow immediately from the description of γ_α . For any pair $\alpha, \beta \in \Pi$, consider the 2-dimensional facet, denoted by $[ao^\varepsilon(\lambda), p_\alpha, p_\beta]$, of $\overline{J_W^{\{\varepsilon\}}}(\lambda)$ containing the edges $[ao^\varepsilon(\lambda), p_\alpha]$ and $[ao^\varepsilon(\lambda), p_\beta]$ (Fig. 4). The $[ao^\varepsilon(\lambda), p_\alpha, p_\beta]$ is a parallelogram transversal to the 2-codimensional stratum of the discriminant locus of label $m_{\alpha,\beta}$. The inverse image of the parallelogram in $K_W^\varepsilon(\lambda)$ is a union of $2m_{\alpha,\beta}$ -gons, whose boundary are $m_{\alpha,\beta}$ -alternating sequence of inverse images of $[ao^\varepsilon(\lambda), p_\alpha]$ and $[ao^\varepsilon(\lambda), p_\beta]$. One translates a $2m_{\alpha,\beta}$ -gon $K_{\alpha\beta}$ to $\tilde{K}_{\alpha\beta}$ in a complex direction in $V_\mathbb{C}$ such that i) $\tilde{K}_{\alpha\beta}$ does not meet with reflection hyperplanes, ii) the boundary of $\tilde{K}_{\alpha\beta}$ is homotopic to an alternating sequence of inverse images γ_α and of γ_β . This implies the homotopy relation: $\gamma_\alpha\gamma_\beta \dots = \gamma_\beta\gamma_\alpha \dots$ ($m_{\alpha,\beta}$ -terms).

Fig.4. The 2-Facet $[ao^\varepsilon(\lambda), p_\alpha, p_\beta]$ **Fig.5.** Pencil close to $\overline{\alpha\beta}$ -edge



4.3. Zariski-van Kampen generator system.


We identify the generator system $\{\gamma_\alpha\}_{\alpha \in \Pi}$ in §4.1 with the well-known Zariski-van Kampen generator system. This is achieved since the τ -direction is transversal to the discriminant divisor, and hence the τ -orbits in S_W play a good role as the Zariski pencil.

Choose a base point $*^\varepsilon$ in the central component $\mathcal{C}^{\{\varepsilon\}}$ in $S_{W, \mathbb{R}}^{[\varepsilon]}$. The real line $\tau^{[\varepsilon]}(\mathbb{R})(*^\varepsilon)$ and the real discriminant locus $D_{W, \mathbb{R}}^{[\varepsilon]}$ intersect by l -points (counted with multiplicity). We order them as

$$(4.3.1) \quad t_1 \leq \dots \leq t_{l_1} < *^\varepsilon < t_{l_1+1} \leq \dots \leq t_l,$$

where $u \leq v$ (resp. $u < v$) means $v \in \tau^{[\varepsilon]}(\mathbb{R}_{\geq 0})u$ (resp. $v \in \tau^{[\varepsilon]}(\mathbb{R}_{> 0})u$). For a generic $*^\varepsilon$ (precisely, if $\omega_{W,2}(*^\varepsilon) \neq 0$), the points t_1, \dots, t_l are distinct. Inside the complexification $\tau(\mathbb{C})(*^\varepsilon)$ of the line, we choose l -closed paths $\delta_1, \dots, \delta_l$ based at $*^\varepsilon$ and turning once around the points t_1, \dots, t_l counter-clockwise as in the Fig. 6.

Fig. 6. The Zariski-van Kampen generators on a τ -orbit (cf. Fig.5.).



It is well-known that they generate the fundamental group of the complement of the discriminant locus and that their fundamental relations are determined by the Zariski-van Kampen method.

We compare the two generator systems introduced in §4.1 and in the present subsection. Let ao^ε and $*^\varepsilon$ be the base points chosen in §4.1 and §4.3. The paths in $\mathcal{C}^{\{\varepsilon\}}$ connecting ao^ε and $*^\varepsilon$ consist of a single homotopy class, denoted by $[ao^\varepsilon, *^\varepsilon]$, since $\mathcal{C}^{\{\varepsilon\}}$ is simply connected.

Theorem 4.2. *The conjugation by $[ao^\varepsilon, *^\varepsilon]$ induces the bijection of the generator systems:*

$$\{\gamma_\alpha\}_{\alpha \in \Pi} \simeq \{\delta_i\}_{1 \leq i \leq l},$$

where the bijection $\{1, \dots, l\} \simeq \Pi$ of the index sets is given by the map $c_W: i \leftrightarrow \alpha \Leftrightarrow c_W(t_i) \in H_\alpha$. The homotopy classes $\delta_1, \dots, \delta_{l_1}$ mutually commute, and so do the homotopy classes $\delta_{l_1+1}, \dots, \delta_l$.

4.4. Proof of Theorem 4.2.

Let $*^\varepsilon \in \mathcal{C}^{\{\varepsilon\}}$ be the base point as above. Consider the real τ -orbit $\tau^{[\varepsilon]}(\mathbb{R})(*^\varepsilon)$. Due to the homeomorphism (3.5.5), we identify the tube domain $(\pi_\tau^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}})$ with $(\pi_\Pi)^{-1}(E_{\Gamma(W)})$. For $\alpha \in \Pi_1$, the half line $\tau^{[\varepsilon]}(\sigma\varepsilon^{[h/2]}\mathbb{R}_{> 0}) \cdot *^\varepsilon$ intersects the hyperplane $H_\alpha = \{\phi_\alpha = 0\}$ at a point, which we write t_α . For $\alpha \in \Pi_2$, the other half line $\tau^{[\varepsilon]}(-\sigma\varepsilon^{[h/2]}\mathbb{R}_{> 0}) \cdot *^\varepsilon$ intersects the hyperplane $H_\alpha = \{\phi_\alpha = 0\}$ at a point, which we write t_α .

If $\pi_\tau(*^\varepsilon) \notin B_{W,2,\mathbb{R}}^{[\varepsilon]}$, then all t_α 's are distinct. We choose the paths as in Fig. 6. Let us index them by the set Π : the path turning around the point t_α shall be called δ_α .

We first show that

i) the homotopy classes of δ_α for $\alpha \in \Pi_1$ mutually commute and so do the homotopy classes of δ_α for $\alpha \in \Pi_2$,

ii) for two choices of base points $*_1$ and $*_2 \in \mathcal{C}^{\{\varepsilon\}}$, the conjugation by a path connecting $*_1$ and $*_2$ in $\mathcal{C}^{\{\varepsilon\}}$ induces one to one correspondences between the generators of the same index.

They follow from the descriptions of the discriminant locus and the central component in Theorem C, (5) and (6) as follows. Move the line $\tau^{[\varepsilon]}(\mathbb{R})(*)$ by moving $*$ in $\mathcal{C}^{\{\varepsilon\}}$ and trace the l -points $\{t_\alpha\}_{\alpha \in \Pi} = \tau^{[\varepsilon]}(\mathbb{R})(*) \cap D_{W,\mathbb{R}}^{[\varepsilon]}$ in the line. The fact that $E_W^{\{\varepsilon\}}$ does not intersect higher bifurcation set $B_{W,\geq 3}$ but only with ordinary $B_{W,2,\mathbb{R}}^{[\varepsilon]}$ implies that one obtains only some commutative relations among generators. As far as $*$ moves inside $\mathcal{C}^{\{\varepsilon\}}$, the set of points $\{t_\alpha \mid \alpha \in \Pi_1\}$ and the set of the points $\{t_\alpha \mid \alpha \in \Pi_2\}$ are separated by the base point $*$ (Theorem C (6)). Theorem C (5) claims that if α and β belong to the same Π_1 or Π_2 , then the hyperplanes H_α and H_β are normal crossing in the tube domain $(\pi_{\tau,\mathbb{R}}^{[\varepsilon]})^{-1}(\pi_{\tau,\mathbb{R}}^{[\varepsilon]}(\mathcal{C}^{\{\varepsilon\}})) \subset (\pi_{\tau,\mathbb{R}}^{[\varepsilon]})^{-1}(E_W^{\{\varepsilon\}})$. This proves i) and ii).

Next, we show that

iii) the conjugation by a path connecting $ao^\varepsilon(\lambda)$ and $*^\varepsilon$ in $\mathcal{C}^{\{\varepsilon\}}$ induces a correspondence of the homotopy class of γ_α to that of δ_α for $\alpha \in \Pi$.

We prove this by ii) as follows.

For each $\alpha \in \Pi$, we can choose a base point $*_\alpha$ such that the line $\tau^{[\varepsilon]}(\mathbb{R}) \cdot *_\alpha$ and the discriminant locus $D_{W,\mathbb{R}}^{[\varepsilon]}$ intersect at the point p_α introduced in §4.1. Let $[ao^{\{\varepsilon\}}(\lambda), *_\alpha]$ be a path in $\mathcal{C}^{\{\varepsilon\}}$ connecting the two vertices and let $[*_\alpha, p_\alpha]$ be the interval in the line $\tau^{[\varepsilon]}(\mathbb{R}) \cdot *_\alpha$. Then, the path $[ao^{\{\varepsilon\}}(\lambda), p_\alpha]$ (the edge of $J_W^{\{\varepsilon\}}(\lambda^{[\varepsilon]})$ connecting the vertices $ao^{\{\varepsilon\}}(\lambda)$ and p_α) is homotopic to the composition of paths $[ao^{\{\varepsilon\}}(\lambda), *_\alpha][*_\alpha, p_\alpha]$ in $\mathcal{C}^{\{\varepsilon\}}$. This means that the conjugation by $[ao^{\{\varepsilon\}}(\lambda), *_\alpha]$ induces the correspondence between the homotopy classes of γ_α and that of δ_α . This fact together with ii) implies iii).

Note. That the generator system $\{\delta_\alpha\}_{\alpha \in \Pi}$ satisfies the braid relation can be shown by the standard Zariski-van Kampen method.


4.5. Comparison of generator systems for $\varepsilon \in \{\pm 1\}$.

Our identification of the fundamental group of $S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}$ with the Artin group (either by the use of $\{\gamma_\alpha\}_{\alpha \in \Pi}$ or of $\{\delta_\alpha\}_{\alpha \in \Pi}$) depends on the choice of the base point locus. Actually, depending on $\varepsilon \in \{\pm 1\}$, the base point is chosen in the central component $\mathcal{C}^{\{\varepsilon\}}$. For a path

γ connecting $\mathcal{C}^{\{+\}}$ and $\mathcal{C}^{\{-}}$, an isomorphism of the two fundamental groups is given by $\text{Ad}_{[\gamma]}$, where $[\gamma]$ is the homotopy class of the path and by Ad we mean the conjugation action on homotopy classes.

Here we choose the simplest path connecting the two components $\mathcal{C}^{\{+1\}}$ and $\mathcal{C}^{\{-1\}}$. Namely, let the base points ao^+ (resp. ao^-) lie on the positive (resp. negative) half v.o. axis AO^ε . Consider the complexification of the v.o. axis. Inside the complex v.o. axis deleted by the origin, let γ_+ (resp. γ_-) be the path connecting ao^+ to ao^- (resp. ao^- to ao^+) by turning half around the origin counter-clockwise.

Fig. 7 The complexification of the vertex orbit axis AO



By the use of them, we have the isomorphisms:

$$\begin{aligned} \pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^+) &\xrightarrow{\text{Ad}_{[\gamma_+]}} \pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^-) \\ \pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^+) &\xleftarrow{\text{Ad}_{[\gamma_-]}} \pi_1(S_{W,\mathbb{C}} \setminus D_{W,\mathbb{C}}, ao^-) \end{aligned}$$

In order to state the following Assertion, we recall the *fundamental element* [BS]. Consider the monoid generated by the letters a_α ($\alpha \in \Pi$) satisfying the braid relations as the defining relations. The fundamental element Δ is the shortest element in the monoid which is divisible (from both sides) by any of the generators a_α . Such Δ exists uniquely in the monoid. Since the monoid is embedded in the Artin group, we identify Δ with its image.

Assertion 4.3. *The homotopy classes $[\gamma_+][\gamma_-]$ and $[\gamma_-][\gamma_+]$ are the fundamental element Δ in each of the fundamental group based at ao^+ and ao^- regarded as an Artin group with respect to the generator systems $\{\gamma_\alpha\}_{\alpha \in \Pi}$.*

This fact follows from i) the length of $[\gamma_+][\gamma_-]$ as an element of the monoid is given by $l([\gamma_+][\gamma_-]) = \deg(\Delta_W)/\deg(R) = hl/2 = l(\Delta)$, and ii) a description of the monodromy of $[\gamma_+][\gamma_-]$ on $b_W : T_{W,\mathbb{C}} \dashrightarrow V_{\Pi,\mathbb{C}}$, see [S4, §8,9).

Note. 1. Δ^2 belongs to the center of the Artin group for any type of W . However, Δ does not belong to the center if W is of type A_n for $n \geq 2$, D_{2k+1} for $k \geq 2$, E_6 and $I_2(2q+1)$ for $q \geq 1$.

2. The fundamental element Δ projects to the longest element of the Coxeter group W .

APPENDIX. DIHEDRAL GROUP OF TYPE $I_2(h)$

For the dihedral group $W(I_2(h))$ ($h \geq 3$), we describe the data 1 – 9.

1. The action of $W(I_2(h))$ on U (recall Notation §2.5 i) – vi):
 $W(I_2(h))$ is generated by the reflections α_1 and α_2 on $U := \mathbb{R} \oplus \mathbb{R}i$ given by $\alpha_1(z) := \bar{z}$ and $\alpha_2(z) := \omega^2 \bar{z}$, where $z := x + yi \in U$ and $\omega := \exp(\pi i/h)$, which satisfy the fundamental relation $(\alpha_1 \alpha_2)^h = 1$.
2. The set of reflections of $W(I_2(h))$ is given by

$$R(W(I_2(h))) = \{\alpha_k := \alpha_1(\alpha_1 \alpha_2)^{k-1} \mid k = 1, \dots, h\}.$$

3. The normalizer $N(W(I_2(h)))$ in $\text{GL}(U)$ is equal to $W(I_2(2h))$ and $N(W(I_2(h)))/W(I_2(h)) = \{[1], [\beta]\}$ where β is the reflection: $\beta(z) = \omega \bar{z}$. One has $[-1] = [1]$ for even h , and $[-1] = [\beta]$ for odd h .
4. The twisted real vertex orbit planes are given by

$$S_{W(I_2(h)), \mathbb{R}}^{[+1]} = \text{Spec}(\mathbb{R}[R, S])_{\mathbb{R}} \text{ and } S_{W(I_2(h)), \mathbb{R}}^{[\beta]} = \text{Spec}(\mathbb{R}[R, S^{[\beta]}])_{\mathbb{R}},$$

where $R = R^{[\beta]} = x^2 + y^2$, $S = S^{[1]} = \text{Re}((x + iy)^h)$ and $S^{[\beta]} = S/\sqrt{-1}$.

5. For $[u] \in N(W)/W$, the twisted real discriminant locus $D_{W(I_2(h)), \mathbb{R}}^{[u]}$ in $S_{W(I_2(h)), \mathbb{R}}^{[+1]}$ is defined by the equation:

$$\Delta_{W(I_2(h))} = R^h - S^2 = R^h + (S^{[\beta]})^2 = \varepsilon^h((\varepsilon R)^h - (S^{[\varepsilon]})^2) = 0.$$

This implies that $D_{W(I_2(h)), \mathbb{R}}^{[\beta]} = \{0\}$ for even h , but $D_{W(I_2(h)), \mathbb{R}}^{[\varepsilon]} \neq \{0\}$ for any ε . Therefore, in 7. and 8., we consider only cases for $[\varepsilon]$.

6. The $\tau^{[u]}$ -action on the plane: $(R, S^{[u]}) \mapsto (R, S^{[u]} + \lambda^{[u]})$ for $\lambda^{[u]} \in \mathbb{R}$,
7. The equation for the inverse image $(\pi_{W, \mathbb{R}}^{[\varepsilon]})^{-1}(D_{W(I_2(h)), \mathbb{R}}^{[\varepsilon]} + \lambda^{[\varepsilon]})$ by the polar coordinates $x + iy = \sqrt{\varepsilon} r \exp(i\theta)$ on U^ε is given by

$$\begin{aligned} (\tau^{[\varepsilon]})^*(\lambda^{[\varepsilon]})\Delta_W &= R^h - \varepsilon^h(S^{[\varepsilon]} + \lambda^{[\varepsilon]})^2 \\ &= \varepsilon^h(r^{2h} \sin^2(h\theta) - 2\varepsilon^{h(h-1)/2} r^h \lambda^{[\varepsilon]} \cos(h\theta) - (\lambda^{[\varepsilon]})^2) \\ &= \varepsilon^h(r^h(1 - \varepsilon^{h(h-1)/2} \cos(h\theta)) - \lambda^{[\varepsilon]})(r^h(1 + \varepsilon^{h(h-1)/2} \cos(h\theta)) + \lambda^{[\varepsilon]}) \end{aligned}$$


8. The dual polyhedron is described by the polar coordinates as

$$\begin{aligned} \bar{K}_W^\varepsilon(\lambda^{[\varepsilon]}) &= \{z \in U^\varepsilon \mid \tau^{[\varepsilon]}(\lambda^{[\varepsilon]})\Delta(z) \leq 0\} \cap \{\tau^{[\varepsilon]}(-\lambda^{[\varepsilon]})\Delta(z) \leq 0\} \\ &= \{\sqrt{\varepsilon} r \exp(i\theta) \in U^\varepsilon \mid r^h \leq \lambda^{[\varepsilon]}/(1 - \varepsilon^{h(h-1)/2} \cos(h\theta)) \\ &\quad r^h \leq \lambda^{[\varepsilon]}/(1 + \varepsilon^{h(h-1)/2} \cos(h\theta))\}. \end{aligned}$$

9. For $h=3$ and 4, for $\varepsilon=1$ or -1 and for $\lambda^{[\varepsilon]}=1$, we draw the figures:
 - i) the real discriminant locus $D_{W, \mathbb{R}}^{[\varepsilon]}$ and the $\lambda^{[\varepsilon]}$ -shifted real discriminant locus: $D_{W, \mathbb{R}}^{[\varepsilon]} \pm \lambda^{[\varepsilon]} := \tau^{[\varepsilon]}(\pm \lambda^{[\varepsilon]})(D_{W, \mathbb{R}}^{[u]})$ in $S_{W(I_2(h)), \mathbb{R}}^{[\varepsilon]}$.
 - ii) the inverse images $(\pi_{W, \mathbb{R}}^{[\varepsilon]})^{-1}(D_{W, \mathbb{R}}^{[\varepsilon]})$ and $(\pi_{W, \mathbb{R}}^{[\varepsilon]})^{-1}(D_{W, \mathbb{R}}^{[\varepsilon]} \pm \lambda^{[\varepsilon]})$ in U^ε .
 - iii) the parallelograms $J_W^{[\varepsilon]}(\lambda^{[\varepsilon]})$ and the polyhedra $K_W^\varepsilon(\lambda^{[\varepsilon]})$ (shaded).
 - iv) the two twisted real forms $S_{W(I_2(h)), \mathbb{R}}^{[1]}$ and $S_{W(I_2(h)), \mathbb{R}}^{[\beta]}$ which are embedded in the real 3-space $S_{W(I_2(h)), \mathbb{C}} \cap \{\text{Im}(R) = 0\}$.

1. **h is odd and coset [1]:** In this case, $[\beta] = [-1] \notin [W(I_2(h))]$.


Fig.8. Polyhedra $J_{A_2}^{\{+1\}}$ and $K_{A_2}^{+1}$ for $\lambda^{[+1]} = 1$



3 lines in $V_{A_2}^+$ are the reflection hyperplanes for W_{A_2} .


2. **h is odd and coset $[\beta] = [-1]$:**

Fig.9. Polyhedra $J_{A_2}^{\{-1\}}$ and $K_{A_2}^{-1}$ for $\lambda^{[-1]} = 1$




3 lines in $V_{A_2}^-$ are the reflection hyperplanes for W_{A_2} .

Fig.10. Positions of $S_{A_2, \mathbb{R}}^{[+1]}$ and $S_{A_2, \mathbb{R}}^{[-1]}$ inside $S_{A_2, \mathbb{C}} \cap \{\text{Im}(R) = 0\}$.



3. **h is even and coset $[1] = [-1]$:** In this case, $-1 \in W(I_2(h))$.


Fig.11. Polyhedra $J_{B_2}^{\{\pm 1\}}(\lambda)$ and $K_{B_2}^{\pm 1}(\lambda)$ for $\lambda = 1$.



4. **h is even and coset $[\beta]$:** In this case, $[\beta] \neq [-1] \in [W(I_2(h))]$.

The discriminant locus $D_{W(I_2(h)), \mathbb{R}}^{[\beta]}$ in the real form $S_{W(I_2(h)), \mathbb{R}}^{[\beta]}$ consists only of the origin $\{o\}$. Therefore, we omit the figure for this case.

Fig.12. Positions of $S_{B_2, \mathbb{R}}^{[\pm 1]}$ and $S_{B_2, \mathbb{R}}^{[\beta]}$ inside $S_{B_2, \mathbb{C}} \cap \{\text{Im}(R) = 0\}$.



Remark 15. We compare the multiplicities of two equations of the bifurcation set in the vertex orbit curve $T_{W(I_2(h))} := \text{Spec}(\mathbb{R}[R])$.

i) The discriminant δ of the map $R = z\bar{z} = x^2 + y^2 : U \rightarrow \mathbb{R}$ is equal to R itself. (The free resolution of $\mathbb{R}[x, y]/(\partial_x R, \partial_y R)$ as an $\mathbb{R}[R]$ -module is given by $0 \rightarrow \mathbb{R}[R] \xrightarrow{R} \mathbb{R}[R] \rightarrow \mathbb{R}[x, y]/(\partial_x R, \partial_y R) \rightarrow 0$.)

ii) The discriminant ω of the quadratic polynomial $\Delta_{W(I_2(h))} = R^h - S^2$ in S is equal to $4 \cdot R^h$.

Comparing i) and ii), we obtain a relation:

$$(*) \quad \omega = 4 \cdot \delta^h.$$

REFERENCES

- [Ar1] Arnol'd, Vlasimir I.: Springer numbers and morsification spaces, *J. Alg. Geom.* **1** (1992), 197-214.
- [Ar2] Arnol'd, Vlasimir I.: The calculus of snakes and combinatorics of Bernoulli, Euler and Springer numbers of Coxeter groups, *Russian Math. Surveys* **47**:1 (1992), 1-51
- [A] Artin, Emil: Theorie der Zöpfe, *Abhandlungen aus dem Mathematischen Seminar der Hamburgischen Universität*, 4.Band, (1926), 47-72.
- [B] Bourbaki, N.: *Elements de mathematique*, Fasc. XXXIV, *Groupes et algebres de Lie*, Chs. 4-6, Hermann, Paris 1968.
- [Br1] Brieskorn, Egbert: Die Fundamental Gruppe des Raumes der Regulären Orbits einer endlichen komplexen Spiegelungsgruppe, *Inventiones Math.* **12** (1971), 57-61.
- [Br2] Brieskorn, Egbert: Sur les groupes de tresses. *Sém. Bourbaki*, (1971).
- [BS] Brieskorn, Egbert & Saito, Kyoji: Artin Gruppen und Coxeter Gruppen, *Inventiones Math.* **17** (1972), 245-271.
- [Ch] Chevalley, Claude: Invariants of finite groups generated by reflections, *Amer. J. Math.* **77** (1955), 778-782.
- [C] Coleman, A. John: The Betti numbers of the simple Lie groups, *Canad. J. Math.* **10**, (1958), 349-356.
- [D1] Deligne, Pierre: Les immeubles des groupes de tresses généralisé, *Inventiones Math.* **17** (1972), 273-302.
- [D2] Deligne, Pierre: Action du groupe des tresses sur la catégorie, *Invent. math.* **128** (1997) 159-175.
- [Hi] Hilbert, David: *Math. Annalen* **36** (1890), 473-534.
- [Hu] Hung, Nguyen H.V.: The modulo 2 cohomology algebras of symmetric groups, *Japan. J. Math.*, Vol 13 (1987), 169-208.
- [K] Kostant, Bernard: The principal three-dimensional subgroup and the Betti numbers of a complex simple group, *Amer. J. Math.*, **81** (1959), 973-1032.
- [Lo1] Looijenga, Eduard: The complement of the bifurcation variety of a simple singularity, *Invent. math.* **23** (1974), 105-116.
- [Lo2] Looijenga, Eduard: The Discriminant of a real simple singularity, *Compositio Mathematica*, Vol.37, Fasc.1 (1978), 51-62.

- [Ly] Lyashko, O.V.: The geometry of the bifucation diagrams, *Usp. Math. Nauk* **34**, 205-206 (1979); *Russian Math. Surveys* **34** (1979), 209-210.
- [M] Huin Mui: Duality on the infinite symmetric products, *Acta Math. Vietnam* **5** (1980), 100-149.
- [N] Nakamura, Tokushi: On cohomology operations, *Japan. J. Math.* **33** (1963), 93-145.
- [P-S] Procesi, Claudio & Schwarz, Gerald: Inequqlities defining orbit spaces, *Invent. math.* **81** (1985), 539-554.
- [S1] Saito, Kyoji: On a linear structureof the quotient variety by a finite reflexion group, *Publ. RIMS, Kyoto Univ.* **29** (1993), 535-579.
- [S2] Saito, Kyoji: Period mapping associated to a primitive form, *Publ. RIMS, Kyoto Univ.*, **19** (1983), 1231-1264.
- [S3] Saito, Kyoji: Uniformization of the orbifold of a finite reflection group, *Proceedings of MPI Syposium.*
- [S4] Saito, Kyoji: Polyhedra dual to the Weyl chamber decomposition, in preparation.
- [Sa] Salvetti, Mario: Topology of the complement of real hyperplanes in \mathbb{C}^n , *Invent. math.* **83** (1987), 603-618.
- [So] Solomon, Louis: A decomposition of the group algebra of a finite Coxeter group, *J. Alg.*, **9** (1968), 220-239.
- [Sp1] Springer, Tony A.: Remarks on a combinatorial problem, *Nieuw Arch. Wisk.*(1)**19**(1971),30-36.
- [Sp2] Springer, Tony A.: Reductive Groups, *Proc. of Symposia in Pure Math.* Vol. **33** (1979), Part 1, pp.3-27.
- [Te] Terao, Hiroaki: The Bifucation set and logarithmic vector fields, *Math. Ann* **263** (1983), 313-321.
- [T1] Thom, René: L'équivalence d'une fonction différentiable et d'un polynome, *Topology* **3** (1965), 297-307.
- [T2] Thom, René: *Stabilité Structurelle et Morphogénèse*, W.A. Benjamin, Inc., Advanced Book Program, Reading, Massachusetts 01867