

RIMS-1882

**Topics in the Anabelian Geometry of Mixed-characteristic
Local Fields**

By

Yuichiro HOSHI

Oct 2017



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

TOPICS IN THE ANABELIAN GEOMETRY OF MIXED-CHARACTERISTIC LOCAL FIELDS

YUICHIRO HOSHI

OCTOBER 2017

ABSTRACT. — In the present paper, we study the *anabelian geometry of mixed-characteristic local fields* by an *algorithmic approach*. We begin by discussing some generalities on *log-shells* of mixed-characteristic local fields. One main topic of this discussion is the *difference between the log-shell and the ring of integers*. This discussion concerning log-shells allows one to establish *mono-anabelian reconstruction algorithms* for constructing some objects related to the p -adic valuations. Next, we consider *open* homomorphisms between profinite groups of *MLF-type*. This consideration leads us to a *bi-anabelian result* for absolutely unramified mixed-characteristic local fields. Next, we establish some *mono-anabelian reconstruction algorithms* related to each of *absolutely abelian* mixed-characteristic local fields, mixed-characteristic local fields of *degree one*, and *Galois-specifiable* mixed-characteristic local fields. For instance, we give a *mono-anabelian reconstruction algorithm* for constructing the *Norm map* with respect to the finite extension determined by the uniquely determined minimal mixed-characteristic local subfield. Finally, we apply various results of the present paper to prove some facts concerning outer automorphisms of the absolute Galois groups of mixed-characteristic local fields that arise from *field automorphisms* of the mixed-characteristic local fields.

CONTENTS

INTRODUCTION	2
§0. NOTATIONS AND CONVENTIONS	8
§1. GENERALITIES ON LOG-SHELLS	10
§2. RECONSTRUCTION ALGORITHMS RELATED TO VALUATIONS	18
§3. OPEN HOMOMORPHISMS BETWEEN PROFINITE GROUPS OF MLF-TYPE	25
§4. RECONSTRUCTION ALGORITHMS RELATED TO ABSOLUTELY ABELIAN MLF'S	30
§5. RECONSTRUCTION ALGORITHMS RELATED TO MLF'S OF DEGREE ONE	40
§6. RECONSTRUCTION ALGORITHMS RELATED TO GALOIS-SPECIFIABLE MLF'S	46
§7. ON OUTER AUTOMORPHISMS ARISING FROM FIELD AUTOMORPHISMS I	53
§8. ON OUTER AUTOMORPHISMS ARISING FROM FIELD AUTOMORPHISMS II	59
REFERENCES	68

2010 MATHEMATICS SUBJECT CLASSIFICATION. — 11S20.

KEY WORDS AND PHRASES. — anabelian geometry, mono-anabelian geometry, mono-anabelian reconstruction algorithm, MLF, group of MLF-type, log-shell, absolutely abelian MLF, Galois-specifiable MLF.

This research was supported by JSPS KAKENHI Grant Number 15K04780.

INTRODUCTION

In the present paper, we study the *anabelian geometry of mixed-characteristic local fields*. More specifically, we continue our study [cf. [8], [2], [3]] of the *mono-anabelian geometry* [cf., e.g., [8], Introduction; [8], Remark 1.9.8; [3], Introduction] of mixed-characteristic local fields.

One central object of the study in the present paper is a *mixed-characteristic local field*, i.e., an *MLF*. We shall refer to a finite extension of \mathbb{Q}_p , for some prime number p , as an *MLF* [cf. [3], Definition 1.1]. If k is an *MLF*, then we shall write

- $\mathcal{O}_k \subseteq k$ for the ring of integers of k ,
- $\mathfrak{m}_k \subseteq \mathcal{O}_k$ for the maximal ideal of \mathcal{O}_k ,
- $\underline{k} \stackrel{\text{def}}{=} \mathcal{O}_k/\mathfrak{m}_k$ for the residue field of \mathcal{O}_k ,
- $p_k \stackrel{\text{def}}{=} \text{char}(\underline{k})$ for the residue characteristic of k ,
- $d_k \stackrel{\text{def}}{=} \dim_{\mathbb{Q}_{p_k}}(k_+)$, $f_k \stackrel{\text{def}}{=} \dim_{\mathbb{F}_{p_k}}(\underline{k}_+)$ [cf. the discussion entitled “Rings” in §0],
- $e_k \stackrel{\text{def}}{=} \#(k^\times/(\mathcal{O}_k^\times \cdot p_k^\mathbb{Z}))$ for the absolute ramification index of k ,
- $k^{(d=1)} \subseteq k$ for the [uniquely determined] minimal MLF contained in k ,
- $\epsilon_k \stackrel{\text{def}}{=} 1$ (respectively, $\stackrel{\text{def}}{=} 2$) if $p_k \neq 2$ (respectively, $p_k = 2$),
- a_k for the largest nonnegative integer such that k contains a $p_k^{a_k}$ -th root of unity, and
- $\text{ord}_k: k \setminus \{0\} \rightarrow \mathbb{Z}$ for the [uniquely determined] p_k -adic valuation normalized so that ord_k is surjective

[cf. the notational conventions introduced at the beginning of §1]. Moreover, for a positive integer n , we use the notation “ ζ_n ” to denote a primitive n -th root of unity.

Another central object of the study in the present paper is a [profinite — cf. [3], Proposition 3.3, (i)] group *of MLF-type*. We shall say that a group is *of MLF-type* if the group is isomorphic, as an abstract group, to the absolute Galois group of an MLF [cf. [3], Definition 3.1]. If G is a group *of MLF-type*, then, by applying various *mono-anabelian reconstruction algorithms* [cf., e.g., [8], Introduction; [8], Remark 1.9.8] of [3], §3, to G , we obtain

- a prime number $p(G)$,
- positive integers $d(G)$, $f(G)$, and $e(G)$,
- topological modules $k^\times(G)$ and $k_+(G)$, and
- a monoid $k_\times(G)$

which “*correspond*” to

- the prime number p_k ,
- the positive integers d_k , f_k , and e_k ,
- the topological modules k^\times and k_+ [cf. the discussion entitled “Rings” in §0], and
- the monoid k_\times [cf. the discussion entitled “Fields” in §0],

respectively [cf. [3], Summary 3.15]. Moreover, by applying the *mono-anabelian reconstruction algorithms* of Definition 2.4, (i), (ii), of the present paper to G , we obtain

- nonnegative integers $\epsilon(G)$ and $a(G)$

which “correspond” to

- the nonnegative integers ϵ_k and a_k ,

respectively [cf. Proposition 2.5, (i), of the present paper].

In §1, we discuss some generalities on *log-shells* of MLF’s. If k is an MLF, then we shall refer to the compact open topological submodule

$$\mathcal{I}_k \stackrel{\text{def}}{=} \frac{1}{2p_k} \cdot \log_k(\mathcal{O}_k^\times) \subseteq k_+$$

— where we write $\log_k: \mathcal{O}_k^\times \rightarrow k_+$ for the p_k -adic logarithm — of the topological module k_+ as the *log-shell* of k [cf. [8], Definition 5.4, (iii)]. As is well-known [cf., e.g., [3], Lemma 1.2, (vi)], the log-shell *contains* the compact open topological submodule $(\mathcal{O}_k)_+ \subseteq k_+$ of k_+ :

$$(\mathcal{O}_k)_+ \subseteq \mathcal{I}_k.$$

One main topic of the study of §1 is the *difference* between $(\mathcal{O}_k)_+$ and \mathcal{I}_k . In §1, we prove, for instance, the following result [cf. Proposition 1.5; Lemma 1.8, (i); Proposition 1.10, (i)].

THEOREM A. — *Let k be an MLF. Then the following hold:*

- (i) *The quotient*

$$\mathcal{I}_k / (\mathcal{O}_k)_+$$

is isomorphic, as an abstract module, to the module defined by

$$\prod_{\nu=1}^{\infty} (\mathbb{Z}_+ / p_k^\nu \mathbb{Z}_+)^{\oplus b_k(\nu) - \delta(\nu, a_k)}$$

— where we write

$$b_k(\nu) \stackrel{\text{def}}{=} \left(\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu-1}} \right\rfloor - 2 \cdot \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^\nu} \right\rfloor + \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu+1}} \right\rfloor \right) \cdot f_k$$

and $\delta(i, j) \stackrel{\text{def}}{=} 1$ (respectively, $\stackrel{\text{def}}{=} 0$) if $i = j$ (respectively, $i \neq j$). In particular, the isomorphism class of $\mathcal{I}_k / (\mathcal{O}_k)_+$ **depends only** on p_k , f_k , e_k , and a_k

(ii) *It holds that the submodule $\mathcal{I}_k \subseteq k_+$ coincides with the submodule $(\mathcal{O}_k)_+ \subseteq k_+$ if and only if one of the following three conditions is satisfied:*

- *The prime number p_k is **odd**, and, moreover, the finite extension $k/k^{(d=1)}$ is **unramified**.*

- *The field k is **isomorphic** to the field \mathbb{Q}_2 .*
- *The field k is **isomorphic** to the field $\mathbb{Q}_3(\zeta_3)$.*

- (iii) *We shall define a nonnegative integer*

$$\nu_k$$

as follows:

- If either $p_k \geq 5$ or k is not isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$, then

$$\nu_k \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon_k \cdot e_k \leq p_k^\nu\}.$$

- If $p_k \leq 3$, and k is isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$, then

$$\nu_k \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon_k \cdot e_k \leq p_k^{\nu+1}\} = \min\{\nu \geq 1 \mid \epsilon_k \cdot e_k \leq p_k^\nu\} - 1.$$

Then the nonnegative integer ν_k is the **smallest** integer such that

$$p_k^{\nu_k} \cdot \mathcal{I}_k \subseteq (\mathcal{O}_k)_+ \subseteq \mathcal{I}_k.$$

The various results of §1 may be regarded as “preparatory portions” for the establishment of *mono-anabelian reconstruction algorithms* of §2.

In §2, we establish *mono-anabelian reconstruction algorithms* for constructing, from a group G of *MLF-type*,

- a homomorphism of modules

$$\text{ord}_{\boxtimes}(G): k^\times(G) \longrightarrow \mathbb{Z}_+$$

[cf. Definition 2.2] which “corresponds” [cf. Proposition 2.3] to the p_k -adic valuation $\text{ord}_k: k \setminus \{0\} \rightarrow \mathbb{Z}$ and

- a map of sets

$$\text{ord}_{\boxplus}(G): k_+(G) \setminus \{0\} \longrightarrow \mathbb{Z}$$

[cf. Definition 2.6, (ii)] which “corresponds” [cf. Proposition 2.7, (ii)] to the map $\text{ord}_k^{[\mathbb{Z}]}: k \setminus \{0\} \rightarrow \mathbb{Z}$ of sets [cf. Definition 1.9, (ii)] that satisfies the following condition [cf. Proposition 1.10, (ii)]: For each $a \in k \setminus \{0\}$, it holds that

$$\text{ord}_k(a) \leq \text{ord}_k^{[\mathbb{Z}]}(a) < \text{ord}_k(a) + e_k \cdot (\nu_k + 1)$$

[cf. Theorem A, (iii)], i.e., a sort of “ p_k -adic valuation with an indeterminacy” [cf. Remark 1.10.1; also Remark 2.11.1].

Moreover, we also establish *mono-anabelian reconstruction algorithms* for constructing, from a group G of *MLF-type* such that $\epsilon(G) \cdot e(G) = f(G) + a(G)$ [cf. also Remark 2.11.2], topological submodules

$$\mathfrak{m}^n(G) \subseteq \mathcal{O}_+(G) \subseteq k_+(G)$$

[cf. Definition 2.9, (i), (ii)] — where n is a nonnegative integer — of $k_+(G)$ which “correspond” [cf. Proposition 2.10] to the topological submodules $\mathfrak{m}_k^n \subseteq (\mathcal{O}_k)_+ \subseteq k_+$ of k_+ , respectively.

In §3, we consider *open* homomorphisms between profinite groups of *MLF-type*. One main application of the results of §3 is as follows [cf. Theorem 3.6, Corollary 3.7].

THEOREM B. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a profinite group of **MLF-type**. Let

$$\alpha: G_\circ \longrightarrow G_\bullet$$

be an **open** homomorphism. Then the following hold:

(i) Suppose that $d(G_\circ) \leq d(G_\bullet)$ [which is the case if, for instance, $d(G_\circ) = 1$]. Then α is an **isomorphism**.

(ii) Suppose that $e(G_\circ) \leq e(G_\bullet)$ [which is the case if, for instance, $e(G_\circ) = 1$]. Then α is **injective**.

Theorem B leads us to the following *bi-anabelian* [cf., e.g., [8], Introduction; [8], Remark 1.9.8; [3], Introduction] result [cf. Corollary 3.8].

THEOREM C. — For each $\square \in \{\circ, \bullet\}$, let k_\square be an MLF and \bar{k}_\square an algebraic closure of k_\square ; write $G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$. Suppose that $e_{k_\circ} = 1$. Then it holds that the field k_\circ is **isomorphic** to the field k_\bullet if and only if there exists a **surjection** $G_\circ \twoheadrightarrow G_\bullet$.

In §4, we discuss some *mono-anabelian reconstruction algorithms* related to *absolutely abelian* MLF's. We shall say that an MLF k is *absolutely abelian* if the finite extension $k/k^{(d=1)}$ is Galois, and the Galois group is abelian [cf. Definition 4.2, (ii)]. In §4, we establish, for instance, a *mono-anabelian reconstruction algorithm* for constructing, from a group G of *MLF-type*, a homomorphism of topological modules

$$\text{Nm}_{\text{abs}}(G)$$

[cf. Definition 4.7, (iii)] which “corresponds” [cf. Proposition 4.9, (i)] to the *Norm map* $\text{Nm}_{k/k^{(d=1)}}: k^\times \rightarrow (k^{(d=1)})^\times$ with respect to the finite extension $k/k^{(d=1)}$. This homomorphism $\text{Nm}_{\text{abs}}(G)$ allows one to define the notion of *MLF-Galois label* of G , i.e., the triple consisting of the prime number $p(G)$, the positive integer $d(G)$, and the image of the homomorphism $\text{Nm}_{\text{abs}}(G)$ [cf. Definition 4.10]. By applying the main theorems of [4] and [13], we obtain the following result [cf. Theorem 4.11].

THEOREM D. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a group of **MLF-type**. Suppose that $\{(p(G_\circ), a(G_\circ)), (p(G_\bullet), a(G_\bullet))\} \not\subseteq \{(2, 1)\}$. Then it holds that the group G_\circ is **isomorphic** to the group G_\bullet if and only if the *MLF-Galois label* of G_\circ **coincides** with the *MLF-Galois label* of G_\bullet .

Moreover, in §4, we also obtain the following *bi-anabelian* result [cf. Corollary 4.14].

THEOREM E. — For each $\square \in \{\circ, \bullet\}$, let k_\square be an MLF and \bar{k}_\square an algebraic closure of k_\square ; write $G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$. Suppose that there exists a **surjection** $G_\circ \twoheadrightarrow G_\bullet$ [which thus implies that $p_{k_\circ} = p_{k_\bullet}$ — cf. Proposition 3.4, (iii)] **compatible** with the respective p_{k_\circ} -adic, i.e., p_{k_\bullet} -adic, cyclotomic characters [which is the case if, for instance, the surjection $G_\circ \twoheadrightarrow G_\bullet$ is an **isomorphism** — cf. [3], Proposition 4.2, (iv)]. Then the following hold:

(i) The [uniquely determined] **maximal absolutely abelian MLF** contained in k_\circ is **isomorphic** to the [uniquely determined] **maximal absolutely abelian MLF** contained in k_\bullet .

(ii) Suppose that k_\circ is **absolutely abelian**. Then the field k_\circ is **isomorphic** to the field k_\bullet .

Here, observe that Theorem E, (i), may be regarded as a *refinement* of the main theorem of [6] [cf. Remark 4.14.1].

In §5, we discuss some *mono-abelian reconstruction algorithms* related to MLF's of *degree one*, i.e., such that the integer " $d_{(-)}$ " is *equal to one*. For instance, we establish a *mono-abelian reconstruction algorithm* for constructing, from a group G of *MLF-type* such that $d(G) = 1$ [cf. Remark 5.10.1], a *structure of topological field* on $k_{\times}(G)$ [cf. Definition 5.2] which "*corresponds*" [cf. Theorem 5.4, (i)] to the *topological field structure* of k , i.e., on k_{\times} .

In §6, we discuss *Galois-specifiable* MLF's. We shall say that an MLF k is *Galois-specifiable* if k is Galois over $k^{(d=1)}$, and, moreover, the following condition is satisfied: If L is an MLF such that the absolute Galois group of k is isomorphic to the absolute Galois group of L , then the field k is isomorphic to the field L [cf. Definition 6.1]. We prove the following result [cf. Theorem 5.9, (ii); Remark 5.9.1; Theorem 6.4; Remark 6.4.1].

THEOREM F. — *Let k be an MLF. Consider the following five conditions:*

- (1) *The MLF k is **absolutely abelian** [cf. Definition 4.2, (ii)].*
- (2) *The MLF k is **Galois-specifiable** [cf. Definition 6.1].*
- (3) *The MLF k is **absolutely strictly radical** [cf. Definition 5.6, (iii)].*
- (4) *The MLF k is **absolutely characteristic** [cf. Definition 5.7].*
- (5) *The MLF k is **absolutely Galois** [cf. Definition 4.2, (i)].*

Then the following hold:

- (i) *The implications*

$$(3)$$

$$\Downarrow$$

$$(1) \implies (2) \implies (4) \implies (5)$$

hold.

- (ii) *Suppose that $(p_k, a_k) \neq (2, 1)$. Then the equivalence*

$$(1) \iff (2)$$

holds.

- (iii) *There exists an MLF that **violates** the implication $(4) \implies (2)$ (respectively, $(4) \implies (3)$; $(5) \implies (4)$).*

Moreover, in the present paper, we observe that the condition for an MLF to be *absolutely abelian* and the condition for an MLF to be *Galois-specifiable* may be considered to be "*group-theoretic*" [cf. Remark 4.15.1, (i); Remark 6.14.1], but each of the condition for an MLF to be *absolutely strictly radical*, the condition for an MLF to be *absolutely characteristic*, and the condition for an MLF to be *absolutely Galois* should be considered to be "*not group-theoretic*" [cf. ; Remark 5.9.2, (i); Remark 5.9.2, (ii); Remark 4.15.1, (ii)].

Let k be an MLF and \bar{k} an algebraic closure of k . Write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$. Then let us recall that we have a natural *injection* $\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$ [cf., e.g., [3], Proposition 2.1]. By means of this injection, let us regard $\text{Aut}(k)$ as a subgroup of $\text{Out}(G_k)$:

$$\text{Aut}(k) \subseteq \text{Out}(G_k).$$

In §6, we also establish a *mono-abelian reconstruction algorithm* for constructing, from a group G of MLF-type that satisfies a certain condition [cf. Definition 6.9, (i)] “*corresponding*” [cf. Theorem 6.11] to the condition for an MLF to be *Galois-specifiable*, a collection

$$\text{Orb}_{\text{sqg}}(G)$$

[cf. Definition 6.9, (ii)] of subgroups of $\text{Out}(G)$ which “*corresponds*” [cf. Theorem 6.13, (ii)] to the $\text{Out}(G_k)$ -orbit, i.e., by conjugation, of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$.

In §7 and §8, we discuss outer automorphisms of the absolute Galois groups of MLF’s that arise from *field automorphisms* of the MLF’s. For instance, we prove the following result [cf. Theorem 7.2, (i); Theorem 7.5; Corollary 8.7].

THEOREM G. — *Let k be an MLF and \bar{k} an algebraic closure of k . Write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$. Then the following hold:*

(i) *Suppose that the MLF k is **absolutely characteristic**, and that p_k is **odd**. Then the subgroup*

$$\text{Aut}(k) \subseteq \text{Out}(G_k)$$

*is **not normally terminal**.*

(ii) *Write $k^{(\text{ab})} \subseteq k$ for the [uniquely determined] maximal **absolutely abelian MLF** contained in k . Suppose that a **maximal intermediate field** of $k/k^{(\text{ab})}$ **tamely ramified** over $k^{(\text{ab})}$ does **not coincide** with $k^{(d=1)}$ [which is the case if, for instance, $k^{(\text{ab})} \neq k^{(d=1)}$], and that $(p_k, a_k) \neq (2, 1)$. Let n be a nonnegative integer such that $[k : k^{(\text{ab})}] \in p_k^n \mathbb{Z}$ and A an abelian p_k -group that satisfies the following two conditions:*

(1) *It holds that $\sharp A = p_k^n$.*

(2) *The finite abelian group A is **generated** by at most $(d_k/p_k^n) - 1$ elements.*

*Then there exists a subgroup of $\text{Out}(G_k)$ **isomorphic** to A .*

(iii) *Suppose that p_k is **odd**, and that*

$$k = \mathbb{Q}_{p_k}(\zeta_{p_k}, p_k^{1/p_k}).$$

Then the subgroup

$$\text{Aut}(k) \subseteq \text{Out}(G_k)$$

*is **neither normally terminal nor normal**.*

One motivation of studying Theorem G is as follows [cf. Remark 7.5.2]: Let k be an MLF and \bar{k} an algebraic closure of k . Write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$. Then, as is well-known [cf., e.g., the discussion given at the final portion of [12], Chapter VII, §5], in general, the natural *injection*

$$\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$$

is *not surjective*. Under this state of affairs, one may consider the following problem:

Problem: Is there a certain “suitable” characterization of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$?

[Here, let us observe that

the mono-anabelian reconstruction algorithm of “ $\text{Orb}_{\text{sqg}}(G)$ ” in the discussion preceding Theorem G may be regarded as a certain *affirmative solution* to this problem, i.e., in the case where the MLF k is *Galois-specifiable*.]

From the point of view of this problem, let us observe

the [easily verified] *finiteness* of the group $\text{Aut}(k)$.

In particular, as one of possible solutions to the above problem, one may discuss the following question:

(*_{fin}) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ the *uniquely determined maximal finite* subgroup of $\text{Out}(G_k)$? Put another way, is every element of $\text{Out}(G_k)$ of *finite order* contained in the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$?

Now let us observe that it is immediate that an *affirmative answer* to this question (*_{fin}) implies an *affirmative answer* to the following question (*_{char}), hence also an *affirmative answer* to the following question (*_{nor}):

(*_{char}) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ *characteristic*?

(*_{nor}) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ *normal*?

Then one may easily find that

- Theorem G, (i), is related to the question (*_{nor}),
- Theorem G, (ii) [cf. also the example in Remark 7.5.1], yields a *negative answer* to the question (*_{fin}), and
- Theorem G, (iii), yields a *negative answer* to the question (*_{nor}), hence also *negative answers* to the questions (*_{fin}) and (*_{char}).

This is one motivation of studying Theorem G.

Finally, in Remark 8.7.1, we recall some of the discussions of §8 from the point of view of the notion of “link” [cf. [9], §2.7, (i)].

0. NOTATIONS AND CONVENTIONS

NUMBERS. — If $a \in \mathbb{Q}$ is a rational number, then we shall write $\lfloor a \rfloor \in \mathbb{Z}$ for the largest integer such that $\lfloor a \rfloor \leq a$.

SETS. — If S is a finite set, then we shall write $\sharp S$ for the *cardinality* of S . If G is a group, and T is a set equipped with an action of G , then we shall write $T^G \subseteq T$ for the subset of G -invariants of T .

MONOIDS. — In the present paper, every “monoid” is assumed to be *commutative*. Let M be a [multiplicative] monoid. We shall write $M^\times \subseteq M$ for the abelian group of

invertible elements of M . We shall write M^{gp} for the *groupification* of M [i.e., the abelian group given by the set of equivalence classes with respect to the relation \sim on $M \times M$ defined by, for $(a_1, b_1), (a_2, b_2) \in M \times M$, $(a_1, b_1) \sim (a_2, b_2)$ if there exists an element $c \in M$ of M such that $ca_1b_2 = ca_2b_1$]. We shall write M^{pf} for the *perfection* of M [i.e., the monoid obtained by forming the injective limit of the injective system of monoids

$$\dots \longrightarrow M \longrightarrow M \longrightarrow \dots$$

given by assigning to each positive integer n a copy of M , which we denote by I_n , and to each two positive integers n, m such that n divides m the homomorphism $I_n = M \rightarrow I_m = M$ given by multiplication by m/n]. We shall write $M^{\otimes} \stackrel{\text{def}}{=} M \cup \{*_M\}$; we regard M^{\otimes} as a *monoid* [that contains M as a submonoid] by setting $*_M \cdot *_M \stackrel{\text{def}}{=} *_M$ and $a \cdot *_M \stackrel{\text{def}}{=} *_M$ for every $a \in M$.

MODULES. — Let M be a module. If n is a positive integer, then we shall write $M[n] \subseteq M$ for the submodule obtained by forming the kernel of the endomorphism of M given by multiplication by n . We shall write $M_{\text{tor}} \stackrel{\text{def}}{=} \bigcup_{n \geq 1} M[n] \subseteq M$ for the submodule of torsion elements of M and

$$M^{\wedge} \stackrel{\text{def}}{=} \varprojlim_n M/(n \cdot M)$$

— where the projective limit is taken over the positive integers n . [So if M is finitely generated, then M^{\wedge} coincides with the profinite completion of M .]

GROUPS. — Let G be a group and $H \subseteq G$ a subgroup of G . We shall write $Z_G(H) \subseteq G$ for the *centralizer* of H in G [i.e., the subgroup consisting of $g \in G$ such that $gh = hg$ for every $h \in H$] and $N_G(H) \subseteq G$ for the *normalizer* of H in G [i.e., the subgroup consisting of $g \in G$ such that $gH = Hg$]. We shall say that H is *normally terminal* in G if $N_G(H) = H$, or, alternatively, $N_G(H) \subseteq H$.

TOPOLOGICAL GROUPS. — If G is a topological group, then we shall write G^{ab} for the *abelianization* of G [i.e., the quotient of G by the closure of the commutator subgroup of G], $G^{\text{ab-tor}} \stackrel{\text{def}}{=} (G^{\text{ab}})_{\text{tor}} \subseteq G^{\text{ab}}$, and $G^{\text{ab/tor}}$ for the quotient of G^{ab} by the closure of $G^{\text{ab-tor}} \subseteq G^{\text{ab}}$. If H is a profinite group, and p is a prime number, then we shall write $H^{(p)}$ for the *maximal pro- p quotient* of H .

RINGS. — In the present paper, every “ring” is assumed to be *unital, associative, and commutative*. Let R be a ring. Then we shall write R_+ for the underlying additive module of R and $R^{\times} \subseteq R$ for the multiplicative group of units of R . If, moreover, R is a Dedekind domain, then we shall write $R^{\triangleright} \subseteq R$ for the multiplicative monoid of nonzero elements of R . [So if R is a Dedekind domain, then we have a natural inclusion $R^{\times} \subseteq R^{\triangleright}$ of monoids.]

FIELDS. — Let K be a field. Then we shall write $\mu(K) \stackrel{\text{def}}{=} (K^{\times})_{\text{tor}}$ for the group of roots of unity in K and $K_{\times} = K^{\times} \cup \{0\}$ for the underlying multiplicative monoid of K . [So we have a natural isomorphism $(K^{\times})^{\otimes} \xrightarrow{\sim} K_{\times}$ of monoids that sends $*_{K^{\times}} \mapsto 0$]. If, moreover,

K is algebraically closed and of characteristic zero, then we shall write

$$\Lambda(K) \stackrel{\text{def}}{=} \varprojlim_n \boldsymbol{\mu}(K)[n] = \varprojlim_n K^\times[n]$$

— where the projective limits are taken over the positive integers n — and refer to $\Lambda(K)$ as the *cyclotome* associated to K . Thus, the cyclotome has a natural structure of profinite, hence also topological, module and is isomorphic, as an abstract topological module, to $\widehat{\mathbb{Z}}_+$.

1. GENERALITIES ON LOG-SHELLS

In the present §1, let

$$k$$

be an *MLF* — i.e., a finite extension of \mathbb{Q}_p , for some prime number p [cf. [3], Definition 1.1] — and

$$\bar{k}$$

an algebraic closure of k . We shall write

- $\mathcal{O}_k \subseteq k$ for the ring of integers of k ,
- $\mathfrak{m}_k \subseteq \mathcal{O}_k$ for the maximal ideal of \mathcal{O}_k ,
- $\underline{k} \stackrel{\text{def}}{=} \mathcal{O}_k/\mathfrak{m}_k$ for the residue field of \mathcal{O}_k ,
- $\mathcal{O}_k^{\leq n} \stackrel{\text{def}}{=} 1 + \mathfrak{m}_k^n \subseteq \mathcal{O}_k^\times$ [where n is a positive integer] for the n -th higher unit group of \mathcal{O}_k ,
- $\mathcal{O}_k^{\times} \stackrel{\text{def}}{=} \mathcal{O}_k^{\leq 1}$ for the group of principal units of \mathcal{O}_k ,
- μ_k for the [uniquely determined] Haar measure on [the locally compact topological module] k_+ normalized so that $\mu_k((\mathcal{O}_k)_+) = 1$,
- $p_k \stackrel{\text{def}}{=} \text{char}(\underline{k})$ for the residue characteristic of k ,
- $d_k \stackrel{\text{def}}{=} \dim_{\mathbb{Q}_{p_k}}(k_+)$,
- $f_k \stackrel{\text{def}}{=} \dim_{\mathbb{F}_{p_k}}(\underline{k}_+)$,
- $e_k \stackrel{\text{def}}{=} \sharp(k^\times / (\mathcal{O}_k^\times \cdot p_k^{\mathbb{Z}}))$ for the absolute ramification index of k ,
- $\log_k: \mathcal{O}_k^\times \rightarrow k_+$ for the p_k -adic logarithm,
- $\mathcal{I}_k \stackrel{\text{def}}{=} (2p_k)^{-1} \cdot \log_k(\mathcal{O}_k^\times) \subseteq k_+$ for the log-shell of k ,
- $\mathcal{O}_{\bar{k}} \subseteq \bar{k}$ for the ring of integers of \bar{k} ,
- $\underline{\bar{k}}$ for the residue field of $\mathcal{O}_{\bar{k}}$,
- $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$,
- $I_k \subseteq G_k$ for the inertia subgroup of G_k ,
- $P_k \subseteq I_k$ for the wild inertia subgroup of G_k , and
- $\text{Frob}_k \in \text{Gal}(\bar{k}/\underline{\bar{k}}) \stackrel{\sim}{\leftarrow} G_k/I_k$ for the [$\sharp k$ -th power] Frobenius element

[cf. the notational conventions introduced in the discussions following [3], Definition 1.1, and [3], Lemma 1.3]. We shall write, moreover,

- $k^{(d=1)} \subseteq k$ for the [uniquely determined] minimal MLF contained in k ,
- $e_k^{[\mu]} = \lfloor e_k / (p_k - 1) \rfloor$,
- $\epsilon_k \stackrel{\text{def}}{=} 1$ (respectively, $\stackrel{\text{def}}{=} 2$) if $p_k \neq 2$ (respectively, $p_k = 2$) [cf. [3], Lemma 1.3, (iii)],
- a_k for the largest nonnegative integer such that k contains a $p_k^{a_k}$ -th root of unity [i.e. the “ a ” in [3], Lemma 1.2, (i)],
- $a_k^{[\delta]} \stackrel{\text{def}}{=} 0$ (respectively, $\stackrel{\text{def}}{=} 1$) if $a_k = 0$ (respectively, $a_k \neq 0$),
- $\mathcal{I}_k^{(n)} \stackrel{\text{def}}{=} (2p_k)^{-1} \cdot \log_k(\mathcal{O}_k^{\prec n}) \subseteq \mathcal{I}_k$ [where n is a positive integer], and
- $\text{ord}_k: k \setminus \{0\} \rightarrow \mathbb{Z}$ for the [uniquely determined] p_k -adic valuation normalized so that ord_k is surjective.

Finally, for each positive integer n , let

$$\zeta_n \in \bar{k}$$

be a primitive n -th root of unity.

In the present §1, we discuss some generalities on *log-shells* of MLF’s.

PROPOSITION 1.1. — *The following hold:*

- (i) *It holds that $\mathcal{I}_k^{(1)} = \mathcal{I}_k$.*
- (ii) *It holds that $\mu_k(\mathcal{I}_k) = p_k^{\epsilon_k \cdot d_k - f_k - a_k}$.*
- (iii) *Let n be an integer such that $n > e_k^{[\mu]}$. Then it holds that $\mathcal{I}_k^{(n)} = \mathfrak{m}_k^{n - \epsilon_k \cdot e_k}$.*
- (iv) *If $a_k^{[\delta]} = 1$, then it holds that $(f_k, e_k) = (1, p_k^{a_k - 1} \cdot (p_k - 1))$ if and only if k is isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$.*
- (v) *It holds that $p_k^{a_k - 1} \cdot (p_k - 1) \leq e_k$. If, moreover, $a_k^{[\delta]} = 1$, then it holds that $e_k \in p_k^{a_k - 1} \cdot (p_k - 1) \cdot \mathbb{Z}$.*

PROOF. — Assertion (i) follows from [3], Lemma 1.2, (i), (ii), (v). Assertion (ii) is the content of [3], Lemma 1.3, (iii). Assertion (iii) follows from [11], Chapter II, Proposition 5.5. Finally, since $(f_{\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})}, e_{\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})}) = (1, p_k^{a_k - 1} \cdot (p_k - 1))$ if $a_k^{[\delta]} = 1$ [cf. [11], Chapter II, Proposition 7.13, (i)], assertions (iv), (v) follow immediately from the [easily verified] fact that k always contains an MLF isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$. This completes the proof of Proposition 1.1. \square

LEMMA 1.2. — *Let $a \in k \setminus \{0\}$ be an element of $k \setminus \{0\}$. Then the integer $\text{ord}_k(a) \in \mathbb{Z}$ coincides with the uniquely determined integer n such that $\text{Frob}_k^n \in G_k/I_k$ coincides with the image of $a \in k \setminus \{0\}$ by the composite of the injective homomorphism $\text{rec}_k: k^\times \hookrightarrow G_k^{\text{ab}}$ of [3], Lemma 1.7, and the natural surjection $G_k^{\text{ab}} \twoheadrightarrow G_k/I_k$ [cf. [3], Lemma 1.5, (i)].*

PROOF. — This assertion follows immediately from [3], Lemma 1.7, (1). \square

LEMMA 1.3. — *The following hold:*

(i) *Suppose that $a_k^{[\delta]} = 1$. Let ν be an integer such that $1 \leq \nu \leq a_k$. Then it holds that $\zeta_{p_k^\nu} \in \mathcal{O}_k^{\prec e_k^{[\mu]}/p_k^{\nu-1}}$ [cf. Proposition 1.1, (v)] but $\zeta_{p_k^\nu} \notin \mathcal{O}_k^{\prec (e_k^{[\mu]}/p_k^{\nu-1})+1}$.*

(ii) *Let n be a positive integer. Then the modules $\mathcal{O}_k^{\prec n}/\mathcal{O}_k^{\prec n+1}$, $\mathcal{I}_k^{(n)}/\mathcal{I}_k^{(n+1)}$ are **annihilated** by p_k . In particular, these modules have respective natural structures of \mathbb{F}_{p_k} -vector spaces. Moreover, the \mathbb{F}_{p_k} -vector space $\mathcal{O}_k^{\prec n}/\mathcal{O}_k^{\prec n+1}$ is of **dimension** f_k .*

(iii) *Let n be a positive integer. Then the p_k -adic logarithm $\log_k: \mathcal{O}_k^\times \rightarrow k_+$ determines a **surjection** of \mathbb{F}_{p_k} -vector spaces [cf. (ii)]*

$$\mathcal{O}_k^{\prec n}/\mathcal{O}_k^{\prec n+1} \twoheadrightarrow \mathcal{I}_k^{(n)}/\mathcal{I}_k^{(n+1)}.$$

(iv) *In the situation of (iii), if the integer n is of the form “ $e_k^{[\mu]}/p_k^{\nu-1}$ ” for some integer ν such that $1 \leq \nu \leq a_k$, then the **kernel** of the surjection of (iii) is generated by the image of $\zeta_{p_k^\nu} \in \mathcal{O}_k^{\prec e_k^{[\mu]}/p_k^{\nu-1}}$ [cf. (i)] [hence also of dimension one over \mathbb{F}_{p_k}]. If the integer n is **not of the form** “ $e_k^{[\mu]}/p_k^{\nu-1}$ ” for any integer ν such that $1 \leq \nu \leq a_k$, then the surjection of (iii) is an **isomorphism**.*

PROOF. — Assertion (i) follows immediately from Proposition 1.1, (iv), together with [11], Chapter II, Proposition 7.13, (iv). Assertions (ii), (iii) follow from [11], Chapter II, Proposition 3.10, together with the definition of “ $\mathcal{I}_k^{(n)}$ ”. Assertion (iv) follows immediately from assertion (i), together with [3], Lemma 1.2, (ii), (v). This completes the proof of Lemma 1.3. \square

DEFINITION 1.4.

(i) For each positive integer ν , we shall write

$$b_k(\nu) \stackrel{\text{def}}{=} \left(\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu-1}} \right\rfloor - 2 \cdot \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^\nu} \right\rfloor + \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu+1}} \right\rfloor \right) \cdot f_k.$$

Moreover, we shall write

$$b_k(0) \stackrel{\text{def}}{=} \infty.$$

(ii) We shall write

$$\mathbb{I}_k \stackrel{\text{def}}{=} \prod_{\nu=1}^{\infty} (\mathbb{Z}_+/p_k^\nu \mathbb{Z}_+)^{\oplus b_k(\nu) - \delta(\nu, a_k)}$$

— where we write $\delta(i, j) \stackrel{\text{def}}{=} 1$ (respectively, $\stackrel{\text{def}}{=} 0$) if $i = j$ (respectively, $i \neq j$).

REMARK 1.4.1. — One verifies easily that the isomorphism class of the module \mathbb{I}_k of Definition 1.4, (ii), *depends only* on p_k , f_k , e_k , and a_k .

PROPOSITION 1.5. — *The module $\mathcal{I}_k/(\mathcal{O}_k)_+$ [cf. [3], Lemma 1.2, (vi)] is **isomorphic**, as an abstract module, to the module \mathbb{I}_k . In particular, the isomorphism class of $\mathcal{I}_k/(\mathcal{O}_k)_+$ **depends only** on p_k , f_k , e_k , and a_k [cf. Remark 1.4.1].*

PROOF. — If $(\epsilon_k, e_k) = (1, 1)$, then Proposition 1.5 follows from Proposition 1.1, (ii), (v). Thus, we may assume without loss of generality that $(\epsilon_k, e_k) \neq (1, 1)$. If $a_k^{[\delta]} = 0$, then Proposition 1.5 follows immediately from [10], Theorem 2 [i.e., in the case where we take the “ (N, t) ” of [10], Theorem 2, to be $(\epsilon_k \cdot e_k - 1, 0)$], together with Proposition 1.1, (iii); Lemma 1.3, (iv). If $a_k^{[\delta]} = 1$, then Proposition 1.5 follows immediately from [10], Theorem 3 [i.e., in the case where we take the “ N ” of [10], Theorem 3, to be $\epsilon_k \cdot e_k - 1$], together with Proposition 1.1, (iii); Lemma 1.3, (i), (iv). This completes the proof of Proposition 1.5. \square

REMARK 1.5.1. — One may give an *alternative proof* of Proposition 1.1, (ii), by applying Proposition 1.5. Indeed, it follows from conditions (1) and (2) of [3], Lemma 1.3, (i), that $\mu_k(\mathcal{I}_k) = \sharp(\mathcal{I}_k/(\mathcal{O}_k)_+)$. On the other hand, it follows from Proposition 1.5 that

$$\begin{aligned} \log_{p_k} \left(\sharp(\mathcal{I}_k/(\mathcal{O}_k)_+) \right) &= \log_{p_k} (\sharp \mathbb{I}_k) \\ &= \sum_{\nu=1}^{\infty} \left(\nu \cdot (b_k(\nu) - \delta(\nu, a_k)) \right) = \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^0} \right\rfloor \cdot f_k - a_k = \epsilon_k \cdot d_k - f_k - a_k. \end{aligned}$$

Thus, Proposition 1.1, (ii), holds.

LEMMA 1.6. — *The following hold:*

(i) *The \mathbb{F}_{p_k} -vector space $(\mathcal{I}_k/(\mathcal{O}_k)_+) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$ is of dimension*

$$\epsilon_k \cdot d_k - f_k - a_k^{[\delta]} - \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k} \right\rfloor \cdot f_k.$$

(ii) *If $p_k = 2$, then the \mathbb{F}_{p_k} -vector space $(\mathcal{I}_k/(\mathcal{O}_k)_+) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$ is of dimension $d_k - 1$.*

(iii) *The \mathbb{F}_{p_k} -vector space $(\mathcal{I}_k/(\mathcal{O}_k)_+) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$ is of dimension $< d_k$.*

PROOF. — First, we verify assertion (i). It follows from Proposition 1.5, together with the definition of \mathbb{I}_k , that the dimension under consideration is given by

$$\begin{aligned} \sum_{\nu=1}^{\infty} (b_k(\nu) - \delta(\nu, a_k)) &= \left(\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^0} \right\rfloor - \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^1} \right\rfloor \right) \cdot f_k - a_k^{[\delta]} \\ &= \epsilon_k \cdot d_k - f_k - a_k^{[\delta]} - \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k} \right\rfloor \cdot f_k. \end{aligned}$$

This completes the proof of assertion (i). Assertion (ii) follows from assertion (i), together with the [easily verified] fact that if $p_k = 2$, then $(\epsilon_k, a_k^{[\delta]}) = (2, 1)$.

Finally, we verify assertion (iii). If p_k is *odd*, then since $\epsilon_k = 1$, $f_k \geq 1$, $e_k \geq 1$, and $a_k^{[\delta]} \geq 0$, assertion (iii) follows from assertion (i). If $p_k = 2$, then assertion (iii) follows from assertion (ii). This completes the proof of assertion (iii), hence also of Lemma 1.6. \square

COROLLARY 1.7. — *It holds that*

$$(\mathcal{O}_k)_+ \not\subseteq \frac{1}{2} \cdot \log_k(\mathcal{O}_k^\times).$$

PROOF. — Since \mathcal{I}_k is given by $(2p_k)^{-1} \cdot \log_k(\mathcal{O}_k^\times)$, it follows immediately from [3], Lemma 1.2, (vi), that it holds that $(\mathcal{O}_k)_+$ is contained in $2^{-1} \cdot \log_k(\mathcal{O}_k^\times)$ if and only if $\dim_{\mathbb{F}_{p_k}}((\mathcal{I}_k/(\mathcal{O}_k)_+) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k})$ is equal to $\dim_{\mathbb{F}_{p_k}}(\mathcal{I}_k \otimes_{\mathbb{Z}} \mathbb{F}_{p_k})$, i.e., d_k . Thus, Corollary 1.7 follows from Lemma 1.6, (iii). This completes the proof of Corollary 1.7. \square

LEMMA 1.8. — *The following hold:*

(i) *The following four conditions are equivalent:*

(1) *The submodule $\mathcal{I}_k \subseteq k_+$ coincides with the submodule $(\mathcal{O}_k)_+ \subseteq k_+$.*

(2) *There exists $a(n)$ [necessarily nonpositive — cf. [3], Lemma 1.2, (vi)] integer ν such that the submodule $\mathcal{I}_k \subseteq k_+$ coincides with the submodule $p_k^\nu \cdot (\mathcal{O}_k)_+ \subseteq k_+$.*

(3) *It holds that $\epsilon_k \cdot d_k = f_k + a_k$.*

(4) *One of the following three conditions is satisfied:*

(a) *It holds that $(\epsilon_k, e_k) = (1, 1)$ [i.e., that the prime number p_k is **odd**, and, moreover, $e_k = 1$].*

(b) *It holds that $(p_k, f_k, e_k) = (2, 1, 1)$ [i.e., that k is isomorphic to \mathbb{Q}_2].*

(c) *It holds that $(p_k, f_k, e_k, a_k) = (3, 1, 2, 1)$ [i.e., that k is isomorphic to $\mathbb{Q}_3(\zeta_3)$ — cf. Proposition 1.1, (iv)].*

(ii) *Suppose that either (a) or (b) in (i) is satisfied. Then, for each nonnegative integer ν , it holds that $p_k^\nu \cdot \mathcal{I}_k = \mathfrak{m}_k^\nu$.*

(iii) *Suppose that (c) in (i) is satisfied. Then, for each nonnegative integer ν , it holds that $p_k^\nu \cdot \mathcal{I}_k = \mathfrak{m}_k^{2\nu}$, $p_k^{\nu-1} \cdot \mathfrak{m}_k^3 = \mathfrak{m}_k^{2\nu+1}$.*

(iv) *Suppose that (c) in (i) is satisfied. Write $K \stackrel{\text{def}}{=} k(\zeta_9) \subseteq \bar{k}$. Then the image of the composite*

$$\mathcal{O}_K^\times \hookrightarrow \mathcal{O}_K^\times \xrightarrow{\text{Nm}_{K/k}} \mathcal{O}_k^\times \xrightarrow{\log_k} k_+$$

— where we write $\text{Nm}_{K/k}$ for the Norm map with respect to the finite extension K/k — coincides with $\mathfrak{m}_k^3 \subseteq k_+$.

PROOF. — First, we verify assertion (i). The implication (1) \Rightarrow (2) is immediate. Moreover, the equivalence (1) \Leftrightarrow (3) follows from Proposition 1.1, (ii), and [3], Lemma 1.2, (vi). One also verifies immediately the implication (4) \Rightarrow (3) by straightforward calculations [cf. also Proposition 1.1, (v)].

Next, we verify the implication (2) \Rightarrow (1). Suppose that condition (2) is satisfied. Then since $(\mathcal{O}_k)_+$ is a free \mathbb{Z}_{p_k} -module of rank d_k , we conclude that the module $\mathcal{I}_k/(\mathcal{O}_k)_+$ is a free $\mathbb{Z}/p_k^{-\nu}\mathbb{Z}$ -module of rank d_k . In particular, if $\nu \neq 0$, then the \mathbb{F}_{p_k} -vector space $(\mathcal{I}_k/(\mathcal{O}_k)_+) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$ is of dimension d_k . Thus, it follows from Lemma 1.6, (iii), that $\nu = 0$, as desired. This completes the proof of the implication (2) \Rightarrow (1).

Finally, we verify the implication (3) \Rightarrow (4). Suppose that condition (3) is satisfied. Then since $p_k^{a_k-1} \cdot (p_k - 1) \leq e_k$ [cf. Proposition 1.1, (v)], we obtain that

$$\epsilon_k \cdot f_k \cdot p_k^{a_k-1} \cdot (p_k - 1) \leq \epsilon_k \cdot d_k = f_k + a_k.$$

Now suppose that p_k is *odd*, i.e., ≥ 3 . Then we obtain that

$$(3^{a_k-1} \cdot (p_k - 1) - 1) \cdot f_k \leq a_k.$$

Thus, one verifies easily that either $(p_k, f_k, a_k) = (3, 1, 1)$ or $a_k = 0$. Now observe that it follows from condition (3) that $(p_k, f_k, a_k) = (3, 1, 1)$ (respectively, $a_k = 0$) implies that $(p_k, f_k, e_k, a_k) = (3, 1, 2, 1)$ (respectively, $e_k = 1$), as desired. This completes the proof of the implication (3) \Rightarrow (4) in the case where p_k is *odd*.

Next, suppose that $p_k = 2$. Then, by the above inequality $\epsilon_k \cdot f_k \cdot p_k^{a_k-1} \cdot (p_k - 1) \leq f_k + a_k$, we obtain that

$$(2^{a_k} - 1) \cdot f_k \leq a_k,$$

which thus implies that $a_k = 1$. In particular, it follows from condition (3) that $2d_k = f_k + 1$, i.e., $f_k \cdot (2e_k - 1) = 1$. Thus, we conclude that $(f_k, e_k) = (1, 1)$, as desired. This completes the proof of the implication (3) \Rightarrow (4), hence also of assertion (i).

Assertions (ii), (iii) follow from the implication (4) \Rightarrow (1) of assertion (i). Finally, we verify assertion (iv). Let us first observe that one verifies easily that the integer “ t ” discussed in [14], Chapter V, §3, for the finite Galois extension K/k [that is *totally ramified* and *of degree 3*] is *equal* to 2. Moreover, it follows from Proposition 1.1, (iv), that $f_K = 1$.

Now since “ t ” is equal to 2, it follows from the second equality of [14], Chapter V, §3, Corollary 3, that $\text{Nm}_{K/k}(\mathcal{O}_K^\times)$ *contains* $\mathcal{O}_k^{\times 3}$, which thus implies [cf. [11], Chapter II, Proposition 5.5] that

$$\mathfrak{m}_k^3 \subseteq \log_k(\text{Nm}_{K/k}(\mathcal{O}_K^\times)).$$

Next, observe that since $f_K = 1$, one verifies immediately from Lemma 1.3, (i), (ii), that \mathcal{O}_K^\times is *generated* by $\mathcal{O}_K^{\times 2} \subseteq \mathcal{O}_K^\times$ and $\zeta_9 \in \mathcal{O}_K^\times$. Thus, it follows from [3], Lemma 1.2, (v), that

$$\log_k(\text{Nm}_{K/k}(\mathcal{O}_K^\times)) = \log_k(\text{Nm}_{K/k}(\mathcal{O}_K^{\times 2})).$$

Next, observe that since “ t ” is equal to 2, and $f_K = 1$, it follows immediately from [14], Chapter V, §3, Proposition 5, (iii), together with Lemma 1.3, (ii), that $\text{Nm}_{K/k}(\mathcal{O}_K^{\times 2})$ is *contained* in $\mathcal{O}_k^{\times 3}$, which thus implies [cf. [11], Chapter II, Proposition 5.5] that

$$\log_k(\text{Nm}_{K/k}(\mathcal{O}_K^{\times 2})) \subseteq \mathfrak{m}_k^3.$$

Thus, we conclude that $\mathfrak{m}_k^3 = \log_k(\text{Nm}_{K/k}(\mathcal{O}_K^\times))$, as desired. This completes the proof of assertion (iv), hence also of Lemma 1.8. \square

DEFINITION 1.9.

(i) We shall write

$$\nu_k$$

for the nonnegative integer defined as follows [cf. also Remark 1.9.1 below]:

(1) Suppose that either $(\epsilon_k, e_k) = (1, 1)$ or $(p_k, f_k, e_k, a_k) \in \{(2, 1, 1, 1), (3, 1, 2, 1)\}$. Then

$$\nu_k \stackrel{\text{def}}{=} 0.$$

(2) Suppose that the condition in (1) is not satisfied [which thus implies that $\epsilon_k \cdot e_k - 1 \neq 0$], and that either $p_k \geq 5$ or $k \not\cong \mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$. Then

$$\nu_k \stackrel{\text{def}}{=} \max \left\{ \nu \geq 0 \mid \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu-1}} \right\rfloor \neq 0 \right\}.$$

(3) Suppose that the condition in (1) is not satisfied [which thus implies that $\epsilon_k \cdot e_k - 1 \neq 0$], that $p_k \leq 3$, and that $k \cong \mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$ [which thus implies that $a_k^{[\delta]} = 1$]. Then

$$\nu_k \stackrel{\text{def}}{=} a_k - 1,$$

or, alternatively [cf. the proof of Proposition 1.10, (i), below],

$$\nu_k \stackrel{\text{def}}{=} \max \left\{ \nu \geq 0 \mid \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu-1}} \right\rfloor \neq 0 \right\} - 1.$$

(ii) We shall write

$$\text{ord}_k^{[\mathbb{Z}]}: k \setminus \{0\} \longrightarrow \mathbb{Z}$$

for the map of sets defined by

$$\text{ord}_k^{[\mathbb{Z}]}(a) \stackrel{\text{def}}{=} -e_k \cdot \min\{\nu \in \mathbb{Z} \mid p_k^\nu \cdot a \in \mathcal{I}_k\} + e_k - 1.$$

REMARK 1.9.1. — One verifies easily that the nonnegative integer ν_k of Definition 1.9, (i), may be defined as follows:

(a) If either $p_k \geq 5$ or k is not isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$, then

$$\nu_k \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon_k \cdot e_k \leq p_k^\nu\}.$$

(b) If $p_k \leq 3$, and k is isomorphic to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$, then

$$\nu_k \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon_k \cdot e_k \leq p_k^{\nu+1}\} = \min\{\nu \geq 1 \mid \epsilon_k \cdot e_k \leq p_k^\nu\} - 1.$$

PROPOSITION 1.10. — *The following hold:*

(i) *The nonnegative integer ν_k is the **smallest** integer such that*

$$p_k^{\nu_k} \cdot \mathcal{I}_k \subseteq (\mathcal{O}_k)_+ \subseteq \mathcal{I}_k.$$

(ii) *For each $a \in k \setminus \{0\}$, it holds that*

$$\text{ord}_k(a) \leq \text{ord}_k^{[\mathbb{Z}]}(a) < \text{ord}_k(a) + e_k \cdot (\nu_k + 1).$$

PROOF. — First, we verify assertion (i). Assertion (i) in the case where the condition in (1) of Definition 1.9, (i), is satisfied follows from the implication (4) \Rightarrow (1) of Lemma 1.8, (i). Thus, we may assume without loss of generality that the condition in (1) of Definition 1.9, (i), is not satisfied. [In particular, it holds that $\epsilon_k \cdot e_k - 1 \neq 0$.]

Write

$$\nu_{\mathcal{I}}$$

for the *smallest* integer such that $p_k^{\nu_{\mathcal{I}}} \cdot \mathcal{I}_k \subseteq (\mathcal{O}_k)_+ \subseteq \mathcal{I}_k$ and

$$\nu_b \stackrel{\text{def}}{=} \max\{\nu \geq 0 \mid b_k(\nu) \neq 0\}.$$

Then it is immediate from Proposition 1.5 that

$$\nu_{\mathcal{I}} = \max\{\nu \geq 0 \mid b_k(\nu) - \delta(\nu, a_k) \neq 0\}.$$

In particular, we obtain the following two assertions:

(a) If $b_k(\nu_b) \neq \delta(\nu_b, a_k)$, then it holds that $\nu_{\mathcal{I}} = \nu_b$.

(b) If $b_k(\nu_b) = \delta(\nu_b, a_k)$ [or, alternative, $\nu_b = a_k \geq 1$ and $b_k(\nu_b) = 1$], and $b_k(\nu_b - 1) \neq 0$, then it holds that $\nu_{\mathcal{I}} = \nu_b - 1$.

Moreover, let us observe that it follows immediately from the definition of $b_k(\nu)$ that

$$\nu_b = \max\left\{\nu \geq 0 \mid \left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{\nu-1}} \right\rfloor \neq 0\right\}.$$

Now we verify assertion (i) in the case where the condition in (2) of Definition 1.9, (i), is satisfied. Suppose that the condition in (2) of Definition 1.9, (i), is satisfied. Assume, moreover, that $b_k(\nu_b) = \delta(\nu_b, a_k)$ [which thus implies — cf. the above assertion (b) — that $\nu_b = a_k \geq 1$ and $b_k(\nu_b) = 1$]. Then one verifies immediately that

$$\nu_b = a_k \geq 1, \quad f_k = 1, \quad p_k^{\nu_b-1} \leq \epsilon_k \cdot e_k - 1 < 2 \cdot p_k^{\nu_b-1}.$$

In particular, since $p_k^{a_k-1} \cdot (p_k - 1) \leq e_k$ [cf. Proposition 1.1, (v)], we obtain that

$$\epsilon_k \cdot p_k^{a_k-1} \cdot (p_k - 1) - 1 < 2 \cdot p_k^{a_k-1},$$

which thus implies that

$$\epsilon_k \cdot (p_k - 1) - p_k^{1-a_k} < 2.$$

Thus, since $a_k \geq 1$, we obtain that $p_k \leq 3$.

Next, let us observe that since $a_k \geq 1$, $f_k = 1$, and $p_k \leq 3$, it follows immediately from the condition in (2) of Definition 1.9, (i), together with Proposition 1.1, (iv), (v), that

$$2 \cdot p_k^{a_k-1} \cdot (p_k - 1) \leq e_k.$$

In particular, since $\epsilon_k \cdot e_k - 1 < 2 \cdot p_k^{\nu_b-1}$, we obtain that

$$2 \cdot \epsilon_k \cdot p_k^{a_k-1} \cdot (p_k - 1) - 1 < 2 \cdot p_k^{a_k-1},$$

which thus implies that

$$2 \cdot \epsilon_k \cdot (p_k - 1) - p_k^{1-a_k} < 2.$$

Thus, since $a_k \geq 1$, we obtain a *contradiction*. In particular, we obtain that $b_k(\nu_b) \neq \delta(\nu_b, a_k)$, which thus implies [cf. the above assertion (a)] assertion (i) in the case where the condition in (2) of Definition 1.9, (i), is satisfied. This completes the proof of assertion (i) in the case where the condition in (2) of Definition 1.9, (i), is satisfied.

Finally, we verify assertion (i) in the case where the condition in (3) of Definition 1.9, (i), is satisfied. Suppose that the condition in (3) of Definition 1.9, (i), is satisfied. Then since k is *isomorphic* to $\mathbb{Q}_{p_k}(\zeta_{p_k^{a_k}})$, and $a_k^{[\delta]} = 1$, it follows from Proposition 1.1, (iv), that $e_k = p_k^{a_k-1} \cdot (p_k - 1)$. In particular, since $p_k \leq 3$, we obtain that

$$\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{a_k}} \right\rfloor = \left\lfloor \frac{\epsilon_k \cdot p_k^{a_k-1} \cdot (p_k - 1) - 1}{p_k^{a_k}} \right\rfloor = \left\lfloor \epsilon_k - \frac{\epsilon_k}{p_k} - \frac{1}{p_k^{a_k}} \right\rfloor = 0,$$

$$\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{a_k - 1}} \right\rfloor = \left\lfloor \frac{\epsilon_k \cdot p_k^{a_k - 1} \cdot (p_k - 1) - 1}{p_k^{a_k - 1}} \right\rfloor = \left\lfloor \epsilon_k \cdot p_k - \epsilon_k - \frac{1}{p_k^{a_k - 1}} \right\rfloor = 1,$$

$$\left\lfloor \frac{\epsilon_k \cdot e_k - 1}{p_k^{a_k - 2}} \right\rfloor = \left\lfloor \frac{\epsilon_k \cdot p_k^{a_k - 1} \cdot (p_k - 1) - 1}{p_k^{a_k - 2}} \right\rfloor = \left\lfloor \epsilon_k \cdot p_k^2 - \epsilon_k \cdot p_k - \frac{1}{p_k^{a_k - 2}} \right\rfloor \geq 3.$$

Thus, since $f_k = 1$ [cf. Proposition 1.1, (iv)], we conclude that

$$\nu_b = a_k \geq 1, \quad b_k(\nu_b) = \delta(\nu_b, a_k), \quad b_k(\nu_b - 1) \neq 0.$$

In particular, assertion (i) in the case where the condition in (3) of Definition 1.9, (i), is satisfied follows from the above assertion (b). This completes the proof of assertion (i) in the case where the condition in (3) of Definition 1.9, (i), is satisfied, hence also of assertion (i).

Next, we verify assertion (ii). Write $N \stackrel{\text{def}}{=} -\min\{\nu \in \mathbb{Z} \mid p_k^\nu \cdot a \in \mathcal{I}_k\}$. Then it follows from the definition of N that $p_k^{-N} \cdot a \in \mathcal{I}_k$ but $p_k^{-N-1} \cdot a \notin \mathcal{I}_k$. Thus, it follows from assertion (i) that $p_k^{\nu_k - N} \cdot a \in p_k^{\nu_k} \cdot \mathcal{I}_k \subseteq (\mathcal{O}_k)_+$ but $p_k^{-N-1} \cdot a \notin (\mathcal{O}_k)_+$. In particular, we obtain that $\text{ord}_k(p_k^{\nu_k - N} \cdot a) \geq 0$ and $\text{ord}_k(p_k^{-N-1} \cdot a) < 0$, which thus implies that

$$e_k \cdot (N - \nu_k) \leq \text{ord}_k(a) < e_k \cdot (N + 1).$$

Thus, it follows from the definition of $\text{ord}_k^{[Z]}(a)$ that

$$\text{ord}_k^{[Z]}(a) - e_k + 1 - e_k \cdot \nu_k \leq \text{ord}_k(a) \leq \text{ord}_k^{[Z]}(a).$$

This completes the proof of assertion (ii), hence also of Proposition 1.10. \square

REMARK 1.10.1. — By Proposition 1.10, (ii), one may regard the map $\text{ord}_k^{[Z]} : k \setminus \{0\} \rightarrow \mathbb{Z}$ of Definition 1.9, (ii), as a sort of “ p_k -adic valuation with an indeterminacy”.

2. RECONSTRUCTION ALGORITHMS RELATED TO VALUATIONS

In the present §2, we maintain the notational conventions introduced at the beginning of §1. In particular, we have been given an *MLF*

$$k.$$

Moreover, let

$$G$$

be a [*profinite* — cf. [3], Proposition 3.3, (i)] group of *MLF-type* [cf. [3], Definition 3.1]. Thus, by applying the various *group-theoretic reconstruction algorithms* [cf. [8], Remark 1.9.8] of [3], §3, and [3], §4, to the group G of *MLF-type*, we obtain

- a prime number $p(G)$,
- positive integers $d(G)$, $f(G)$, and $e(G)$,
- subgroups $P(G) \subseteq I(G) \subseteq G$ of G ,
- an element $\text{Frob}(G) \in G/I(G)$ of $G/I(G)$,
- topological monoids $\mathcal{O}^<(G) \subseteq \mathcal{O}^\times(G) \subseteq \mathcal{O}^>(G) \subseteq k^\times(G) \xrightarrow{\text{rec}(G)} G^{\text{ab}}$,

- monoids $k^\times(G) \subseteq k_\times(G)$ and $k_\times(G)$,
- topological modules $\mathcal{I}(G) \subseteq k_+(G)$,
- a measure $\mu(G)$ on $k_+(G)$,
- G -monoids $\overline{\mathcal{O}}^\times(G) \subseteq \overline{\mathcal{O}}^\triangleright(G) \subseteq \overline{k}^\times(G) \subseteq \overline{k}_\times(G)$ and $\overline{k}^\times(G) \subseteq \overline{k}_\times(G)$,
- a G -module $\overline{k}_+(G)$,
- a G -module $\boldsymbol{\mu}(G)$, and
- a topological G -module $\Lambda(G)$

[cf. [3], Summary 3.15; [3], Summary 4.3].

In the present §2, we establish *group-theoretic reconstruction algorithms* for constructing, from the group G of *MLF-type*, a homomorphism of modules

$$\text{ord}_{\boxtimes}(G): k^\times(G) \longrightarrow \mathbb{Z}_+$$

which “*corresponds*” to the p_k -adic valuation $\text{ord}_k: k \setminus \{0\} \rightarrow \mathbb{Z}$ [cf. Definition 2.2, Proposition 2.3 below] and a map of sets

$$\text{ord}_{\boxplus}(G): k_+(G) \setminus \{0\} \longrightarrow \mathbb{Z}$$

which “*corresponds*” to the map $\text{ord}_k^{[\mathbb{Z}]}: k \setminus \{0\} \rightarrow \mathbb{Z}$ of sets of Definition 1.9, (ii) [cf. Definition 2.6, (ii); Proposition 2.7, (ii), below], i.e., a sort of “ p_k -adic valuation with an indeterminacy” [cf. Remark 1.10.1]. Moreover, we also establish *group-theoretic reconstruction algorithms* for constructing, from a group of *MLF-type* that satisfies an additional condition, topological submodules

$$\text{“}\mathfrak{m}^n(-) \subseteq \mathcal{O}_+(-) \subseteq k_+(-)\text{”}$$

— where n is a nonnegative integer — of “ $k_+(-)$ ” which “*correspond*” to the topological submodules $\mathfrak{m}_k^n \subseteq (\mathcal{O}_k)_+ \subseteq k_+$ of k_+ , respectively [cf. Definition 2.9, (i), (ii); Proposition 2.10 below].

LEMMA 2.1. — *The module $k^\times(G)/\mathcal{O}^\times(G)$ is torsion-free and generated by $\text{Frob}(G) \in k^\times(G)/\mathcal{O}^\times(G)$.*

PROOF. — This assertion follows — in light of [3], Proposition 3.6; [3], Proposition 3.9; [3], Proposition 3.11, (i) — from [3], Lemma 1.5, (i) and [3], Lemma 1.7, (1). \square

DEFINITION 2.2. — We shall write

$$\text{ord}_{\boxtimes}(G): k^\times(G) \longrightarrow \mathbb{Z}$$

for the map defined as follows [cf. [2], Theorem 1.4, (7)]: For each $a \in k^\times(G)$, write $\text{ord}_{\boxtimes}(G)(a) \in \mathbb{Z}$ for the uniquely determined [cf. Lemma 2.1] integer n such that the image of $a \in k^\times(G)$ in $k^\times(G)/\mathcal{O}^\times(G)$ coincides with $\text{Frob}(G)^n \in k^\times(G)/\mathcal{O}^\times(G)$.

One verifies immediately that this map is, in fact, a homomorphism $k^\times(G) \rightarrow \mathbb{Z}_+$ of modules.

PROPOSITION 2.3. — *The vertical isomorphism $k^\times \xrightarrow{\sim} k^\times(G_k)$ in the diagram of [3], Proposition 3.11, (i), fits into a **commutative diagram** of modules*

$$\begin{array}{ccc} k^\times & \xrightarrow{\text{ord}_k} & \mathbb{Z}_+ \\ \wr \downarrow & & \parallel \\ k^\times(G_k) & \xrightarrow{\text{ord}_{\boxtimes}(G_k)} & \mathbb{Z}_+. \end{array}$$

PROOF. — This assertion follows — in light of [3], Proposition 3.6; [3], Proposition 3.9; [3], Proposition 3.11, (i) — from Lemma 1.2. \square

REMARK 2.3.1. — Let us observe that one verifies immediately from Proposition 2.3 that

- the open subsets of the topological module $k^\times(G)$ ($\subseteq k_\times(G)$) and,
- for each positive integer n , the subsets of $k_\times(G)$

$$\{a \in k^\times(G) \mid \text{ord}_{\boxtimes}(G)(a) \geq n\} \cup \{*_k(G)\} \subseteq k_\times(G)$$

generate a *topology* on the underlying set of the monoid $k_\times(G)$ by means of which one may regard $k_\times(G)$ as a *topological monoid*. Moreover, one also verifies immediately from Proposition 2.3 that the isomorphism $k_\times \xrightarrow{\sim} k_\times(G_k)$ of [3], Proposition 3.11, (ii), is an *isomorphism of topological monoids*.

DEFINITION 2.4.

(i) We shall write

$$\epsilon(G) \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } p(G) \neq 2 \\ 2 & \text{if } p(G) = 2 \end{cases}$$

[cf. [3], Definition 3.13].

(ii) We shall write

$$a(G) \stackrel{\text{def}}{=} \log_{p(G)} \left(\# \left((k^\times(G)_{\text{tor}})^{(p(G))} \right) \right)$$

[cf. [3], Lemma 1.2, (i); [3], Proposition 3.11, (i)].

(iii) Let ν be a positive integer. Then we shall write

$$b(G, \nu) \stackrel{\text{def}}{=} \left(\left\lfloor \frac{\epsilon(G) \cdot e(G) - 1}{p(G)^{\nu-1}} \right\rfloor - 2 \cdot \left\lfloor \frac{\epsilon(G) \cdot e(G) - 1}{p(G)^\nu} \right\rfloor + \left\lfloor \frac{\epsilon(G) \cdot e(G) - 1}{p(G)^{\nu+1}} \right\rfloor \right) \cdot f(G).$$

(iv) We shall write

$$\mathbb{I}(G) \stackrel{\text{def}}{=} \prod_{\nu=1}^{\infty} (\mathbb{Z}_+ / p(G)^\nu \mathbb{Z}_+)^{\oplus b(G, \nu) - \delta(\nu, a(G))}$$

— where we write $\delta(i, j) \stackrel{\text{def}}{=} 1$ (respectively, $\stackrel{\text{def}}{=} 0$) if $i = j$ (respectively, $i \neq j$).

PROPOSITION 2.5. — *The following hold:*

(i) *It holds that*

$$\epsilon_k = \epsilon(G_k), \quad a_k = a(G_k).$$

(ii) *The module $\mathcal{I}_k/(\mathcal{O}_k)_+$ is **isomorphic**, as an abstract module, to the module $\mathbb{I}(G_k)$.*

PROOF. — Assertion (i) follows from [3], Proposition 3.6, and [3], Proposition 3.11, (i). Assertion (ii) follows — in light of assertion (i); [3], Proposition 3.6 — from Proposition 1.5. This completes the proof of Proposition 2.5. \square

DEFINITION 2.6.

(i) We shall write

$$\nu(G)$$

for the nonnegative integer defined as follows:

(1) If either $p(G) \geq 5$ or $(f(G), e(G)) \neq (1, p(G)^{a(G)-1} \cdot (p(G) - 1))$, then

$$\nu(G) \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon(G) \cdot e(G) \leq p(G)^\nu\}.$$

(2) If $p(G) \leq 3$ and $(f(G), e(G)) = (1, p(G)^{a(G)-1} \cdot (p(G) - 1))$, then

$$\nu(G) \stackrel{\text{def}}{=} \min\{\nu \geq 0 \mid \epsilon(G) \cdot e(G) \leq p(G)^{\nu+1}\}.$$

(ii) We shall write

$$\text{ord}_{\boxplus}(G): k_+(G) \setminus \{0\} \longrightarrow \mathbb{Z}$$

for the map of sets defined by

$$\text{ord}_{\boxplus}(G)(a) \stackrel{\text{def}}{=} -e(G) \cdot \min\{\nu \in \mathbb{Z} \mid p(G)^\nu \cdot a \in \mathcal{I}(G)\} + e(G) - 1.$$

PROPOSITION 2.7. — *The following hold:*

(i) *It holds that*

$$\nu_k = \nu(G_k).$$

(ii) *The vertical isomorphism $k_+ \xrightarrow{\sim} k_+(G_k)$ in the diagram of [3], Proposition 3.11, (iv), fits into a **commutative diagram** of sets*

$$\begin{array}{ccc} k_+ \setminus \{0\} & \xrightarrow{\text{ord}_k^{[\mathbb{Z}]}} & \mathbb{Z} \\ \wr \downarrow & & \parallel \\ k_+(G_k) \setminus \{0\} & \xrightarrow{\text{ord}_{\boxplus}(G_k)} & \mathbb{Z}. \end{array}$$

(iii) *For each $a \in k \setminus \{0\}$, it holds that*

$$\text{ord}_k(a) \leq \text{ord}_{\boxplus}(G_k)(a) < \text{ord}_k(a) + e(G_k) \cdot (\nu(G_k) + 1).$$

PROOF. — Assertion (i) follows from Proposition 2.5, (i), and [3], Proposition 3.6, together with Proposition 1.1, (iv) [cf. also Remark 1.9.1]. Assertion (ii) follows from [3], Proposition 3.6, and [3], Proposition 3.11, (iv). Assertion (iii) follows — in light of assertions (i), (ii); [3], Proposition 3.6 — from Proposition 1.10, (ii). This completes the proof of Proposition 2.7. \square

LEMMA 2.8. — *The following two conditions are equivalent:*

- (i) *It holds that $\epsilon(G) \cdot d(G) = f(G) + a(G)$.*
- (ii) *One of the following three conditions is satisfied:*
 - (1) *It holds that $(\epsilon(G), e(G)) = (1, 1)$.*
 - (2) *It holds that $(p(G), f(G), e(G)) = (2, 1, 1)$.*
 - (3) *It holds that $(p(G), f(G), e(G), a(G)) = (3, 1, 2, 1)$.*

PROOF. — This assertion follows — in light of Proposition 2.5, (i); [3], Proposition 3.6 — from the equivalence (3) \Leftrightarrow (4) of Lemma 1.8, (i). \square

DEFINITION 2.9. — Suppose that $\epsilon(G) \cdot d(G) = f(G) + a(G)$.

- (i) We shall write

$$\mathcal{O}_+(G) \stackrel{\text{def}}{=} \mathcal{I}(G) \subseteq k_+(G).$$

- (ii) Let n be a nonnegative integer. Then we shall define a topological submodule

$$\mathfrak{m}^n(G) \subseteq \mathcal{O}_+(G)$$

of $\mathcal{O}_+(G)$ as follows:

- (1) Suppose that either $(\epsilon(G), e(G)) = (1, 1)$ or $(p(G), f(G), e(G)) = (2, 1, 1)$ [cf. Lemma 2.8]. Then we shall write

$$\mathfrak{m}^n(G) \stackrel{\text{def}}{=} p(G)^n \cdot \mathcal{O}_+(G) \subseteq \mathcal{O}_+(G).$$

- (2) Suppose that $(p(G), f(G), e(G), a(G)) = (3, 1, 2, 1)$ [cf. Lemma 2.8]. If n is even, then we shall write

$$\mathfrak{m}^n(G) \stackrel{\text{def}}{=} p(G)^{n/2} \cdot \mathcal{O}_+(G).$$

If n is odd, then we shall write

$$\mathfrak{m}^n(G) \stackrel{\text{def}}{=} p(G)^{(n-3)/2} \cdot \text{Im}(\mathcal{O}^\times(H) \hookrightarrow \mathcal{O}^\times(H) \rightarrow \mathcal{O}^\times(G) \rightarrow k_+(G))$$

— where we write $H \subseteq G$ for the kernel of the natural action of G on $\boldsymbol{\mu}(G)[9]$ ($\subseteq \boldsymbol{\mu}(G)$); the first arrow “ \hookrightarrow ” is the natural inclusion; the second arrow “ \rightarrow ” is the homomorphism induced by the homomorphism $H^{\text{ab}} \rightarrow G^{\text{ab}}$ determined by the inclusion $H \hookrightarrow G$; the third arrow “ \rightarrow ” is the natural homomorphism.

PROPOSITION 2.10. — *Suppose that $\epsilon_k \cdot e_k = f_k + a_k$, or, alternatively [cf. Proposition 2.5, (i); [3], Proposition 3.6], that $\epsilon(G_k) \cdot e(G_k) = f(G_k) + a(G_k)$. Let n be a nonnegative integer. Then the vertical isomorphism $k_+ \xrightarrow{\sim} k_+(G_k)$ in the diagram of [3], Proposition 3.11, (iv), fits into a **commutative diagram** of topological modules*

$$\begin{array}{ccccc} \mathfrak{m}_k^n & \xrightarrow{\subseteq} & (\mathcal{O}_k)_+ & \xrightarrow{\subseteq} & k_+ \\ \wr \downarrow & & \wr \downarrow & & \wr \downarrow \\ \mathfrak{m}^n(G_k) & \xrightarrow{\subseteq} & \mathcal{O}_+(G_k) & \xrightarrow{\subseteq} & k_+(G_k) \end{array}$$

— where the horizontal arrows are the natural inclusions, and the vertical arrows are isomorphisms.

PROOF. — This assertion follows — in light of Proposition 2.5, (i); [3], Lemma 1.7, (2); [3], Proposition 3.6; [3], Proposition 3.11, (i), (iv) — from Lemma 1.8, (i), (ii), (iii), (iv). \square

Some of the group-theoretic reconstruction algorithms discussed in the present §2 may be summarized as follows.

SUMMARY 2.11.

(i) *There exist **group-theoretic reconstruction algorithms** [cf. [8], Remark 1.9.8] for constructing, from a group G of **MLF-type**,*

- *nonnegative integers $\epsilon(G)$, $a(G)$, and $\nu(G)$ [cf. Definition 2.4, (i), (ii); Definition 2.6, (i)],*
- *a module $\mathbb{I}(G)$ [cf. Definition 2.4, (iv)],*
- *a homomorphism $\text{ord}_{\boxtimes}(G): k^\times(G) \rightarrow \mathbb{Z}_+$ of modules [cf. Definition 2.2], and*
- *a map $\text{ord}_{\boxplus}(G): k_+(G) \setminus \{0\} \rightarrow \mathbb{Z}$ of sets [cf. Definition 2.6, (ii)]*

which “correspond” to

- *the nonnegative integers ϵ_k , a_k , and ν_k [cf. Proposition 2.5, (i); Proposition 2.7, (i)],*
- *the quotient of \mathcal{I}_k by $(\mathcal{O}_k)_+$ [cf. Proposition 2.5, (ii)],*
- *the p_k -adic valuation $\text{ord}_k: k \setminus \{0\} \rightarrow \mathbb{Z}$ [cf. Proposition 2.3], and*
- *the “ p_k -adic valuation with an indeterminacy” [cf. Remark 1.10.1] $\text{ord}_k^{[Z]}: k \setminus \{0\} \rightarrow \mathbb{Z}$ [cf. Proposition 2.7, (ii)],*

respectively.

(ii) *There exist **group-theoretic reconstruction algorithms** for constructing, from a group G of **MLF-type** such that $\epsilon(G) \cdot d(G) = f(G) + a(G)$,*

- *a topological submodule $\mathcal{O}_+(G) \subseteq k_+(G)$ of $k_+(G)$ [cf. Definition 2.9, (i)] and,*
- *for each nonnegative integer n , a topological submodule $\mathfrak{m}^n(G) \subseteq \mathcal{O}_+(G)$ of $\mathcal{O}_+(G)$ [cf. Definition 2.9, (ii)]*

which “correspond” to

- the topological submodule $(\mathcal{O}_k)_+ \subseteq k_+$ of k_+ [cf. Proposition 2.10] and,
 - for each nonnegative integer n , the topological submodule $\mathfrak{m}_k^n \subseteq (\mathcal{O}_k)_+$ of $(\mathcal{O}_k)_+$ [cf. Proposition 2.10],
- respectively.

REMARK 2.11.1. — Let us recall that, as asserted in Summary 2.11, (i), we have established [cf. Definition 2.6, (ii)] a group-theoretic reconstruction algorithm for constructing, from a group G of MLF-type, a map $\text{ord}_{\boxplus}(G): k_+(G) \setminus \{0\} \rightarrow \mathbb{Z}$ of sets which “corresponds” to the “ p_k -adic valuation with an indeterminacy” $\text{ord}_k^{[Z]}: k \setminus \{0\} \rightarrow \mathbb{Z}$ [cf. Remark 1.10.1].

Here, let us also recall that, as discussed in [3], Remark 4.3.1, (i) [cf. also [3], Remark 4.3.2], it is *impossible* to establish a group-theoretic reconstruction algorithm for constructing, from a group G of MLF-type, a topology on the module $\bar{k}_+(G)$ which “corresponds” to the p_k -adic topology on the module \bar{k}_+ . In particular, it is *impossible* to establish a group-theoretic reconstruction algorithm for constructing, from an *arbitrary* group G of MLF-type, a map $k_+(G) \setminus \{0\} \rightarrow \mathbb{Z}$ of sets which “corresponds” to the p_k -adic valuation $k \setminus \{0\} \rightarrow \mathbb{Z}$ [i.e., *without any indeterminacy*].

REMARK 2.11.2. — Let us recall that, as asserted in Summary 2.11, (ii), we have established [cf. Definition 2.9, (i)] a group-theoretic reconstruction algorithm for constructing, from a group G of MLF-type such that $\epsilon(G) \cdot d(G) = f(G) + a(G)$, a topological submodule $\mathcal{O}_+(G) \subseteq k_+(G)$ of $k_+(G)$ which “corresponds” to the topological submodule $(\mathcal{O}_k)_+ \subseteq k_+$ of k_+ .

Here, let us also recall that, as discussed in [3], Remark 4.3.1, (iii) [cf. also [2], Remark 1.4.3], it is *impossible* to establish a group-theoretic reconstruction algorithm for constructing, from an *arbitrary* group G of MLF-type, such a topological submodule of $k_+(G)$.

REMARK 2.11.3. — Let us recall that, as asserted in [3], Summary 3.15, and Summary 2.11, (i), we have established [cf. [3], Definition 3.10, (vi); Definition 2.4, (iv)] group-theoretic reconstruction algorithms for constructing, from a group G of MLF-type, modules $\mathcal{I}(G)$ and $\mathbb{I}(G)$ which “correspond” to the log-shell \mathcal{I}_k and the quotient $\mathcal{I}_k/(\mathcal{O}_k)_+$, respectively.

Here, let us also recall that, as discussed in [3], Remark 4.3.1, (iii) [cf. also [2], Remark 1.4.3], it is *impossible* to establish a group-theoretic reconstruction algorithm for constructing, from an *arbitrary* group G of MLF-type, a surjection $\mathcal{I}(G) \twoheadrightarrow \mathbb{I}(G)$ which “corresponds” to the natural surjection $\mathcal{I}_k \twoheadrightarrow \mathcal{I}_k/(\mathcal{O}_k)_+$.

3. OPEN HOMOMORPHISMS BETWEEN PROFINITE GROUPS OF MLF-TYPE

In the present §3, we maintain the notational conventions introduced at the beginnings of §1 and §2. In particular, we have been given a group *of MLF-type*

$$G.$$

In the present §3, we consider *open* homomorphisms between profinite groups *of MLF-type*. As a consequence of the results in the present §3, we prove that every open homomorphism between profinite groups of MLF-type such that the positive integer “ $e(-)$ ” [cf. the notational conventions introduced at the beginning of §2] of the domain is *equal* to the positive integer “ $e(-)$ ” of the codomain is *injective* [cf. Corollary 3.7 below].

LEMMA 3.1. — *The following hold:*

(i) *The topological module $(G^{(p(G))})^{\text{ab}/\text{tor}}$ is a **free** $\mathbb{Z}_{p(G)}$ -module of rank $d(G) + 1$. Moreover, the kernel of the natural surjection $(G^{(p(G))})^{\text{ab}} \twoheadrightarrow (G^{(p(G))})^{\text{ab}/\text{tor}}$ is **cyclic**.*

(ii) *The closed subgroup $I(G)/P(G) \subseteq G/P(G)$ of $G/P(G)$ **coincides** with the kernel of the natural surjection $G/P(G) \twoheadrightarrow (G/P(G))^{\text{ab}/\text{tor}}$.*

(iii) *It holds that*

$$f(G) = \log_{p(G)} \left(1 + \sharp \left((G/P(G))^{\text{ab}-\text{tor}} \right) \right).$$

PROOF. — Assertion (i) follows immediately — in light of [3], Proposition 3.6; the isomorphism in the final display of [3], Lemma 1.7, (1) — from [3], Lemma 1.2, (i). Assertions (ii), (iii) follow immediately — in light of [3], Proposition 3.6; [3], Proposition 3.9 — from [3], Lemma 1.5, (i), (ii), (iii). This completes the proof of Lemma 3.1. \square

DEFINITION 3.2. — Let J be a profinite group. Then we shall say that a closed subgroup $N \subseteq J$ of J is *quasi-normal* [i.e., in J] if N is normal in an open subgroup of J that contains N .

REMARK 3.2.1. — Let J be a profinite group and $N \subseteq J$ a *quasi-normal* closed subgroup of J . Then one verifies easily that, for each closed subgroup $J_1 \subseteq J$ of J , the closed subgroup $N \cap J_1 \subseteq J_1$ of J_1 is *quasi-normal*.

LEMMA 3.3. — *Let $J \subseteq G$ be a **nontrivial** closed subgroup of G . Then the following hold:*

(i) *Suppose that J is **quasi-normal** in G . Then one of the following three conditions is satisfied [cf. also Remark 3.3.1 below]:*

(1) *The image of J in $G^{(p(G))}$ is **open**.*

(2) *The maximal pro- $p(G)$ quotient $J^{(p(G))}$ is **not topologically finitely generated**.*

(3) *There is **no nontrivial** pro- $p(G)$ quotient of J .*

(ii) Suppose that J is **quasi-normal** in G . Then there exists an open subgroup of J that has a **nontrivial** pro- $p(G)$ quotient.

(iii) Suppose that the maximal pro- $p(G)$ quotient $J^{(p(G))}$ is **not procyclic**. Then there exists an open subgroup $H \subseteq G$ of G such that $J \subseteq H$, and, moreover, the image of J in $(H^{(p(G))})^{\text{ab}/\text{tor}}$ is **nontrivial** [hence also **infinite**].

(iv) Suppose that J is **quasi-normal** in G . Then the following two conditions are equivalent:

(a) There is a **nontrivial** pro- $p(G)$ quotient of J .

(b) There exists an open subgroup $H \subseteq G$ of G such that $J \subseteq H$, and, moreover, the image of J in $(H^{(p(G))})^{\text{ab}/\text{tor}}$ is **nontrivial** [hence also **infinite**].

PROOF. — First, we verify assertion (i). Let us first observe that, to verify assertion (i), it suffices to verify that if J satisfies neither condition (1) nor condition (3), then J satisfies condition (2). Suppose that J satisfies neither condition (1) nor condition (3).

To verify that J satisfies condition (2), let us observe that since J does not satisfy condition (3), there exists a normal open subgroup $N \subseteq G$ of G such that $J/(J \cap N)$ has a quotient that is a $p(G)$ -group. Thus, by considering the composite $J \hookrightarrow J \cdot N \twoheadrightarrow (J \cdot N)/N$ [that determines an isomorphism $J/(J \cap N) \xrightarrow{\sim} (J \cdot N)/N$], we conclude that the image of J in $(J \cdot N)^{(p(G))}$ is *nontrivial*. Next, since J does not satisfy condition (1), the image of J in $(J \cdot N)^{(p(G))}$ is *not open*. Thus, it follows immediately — in light of [3], Proposition 3.6 — from [7], Theorem 1.7, (ii) [cf. also Remark 3.2.1], that the image of J in $(J \cdot N)^{(p(G))}$ is *not topologically finitely generated*, which thus implies that J satisfies condition (2), as desired. This completes the proof of assertion (i).

Assertion (ii) follows immediately — in light of [3], Proposition 3.6 — from [1], Lemma 2.3. Next, we verify assertion (iii). Let us first observe that, by our assumption, there exists a normal open subgroup $N \subseteq G$ of G such that $J/(J \cap N)$ has a quotient that is a *noncyclic* $p(G)$ -group. Write $H \stackrel{\text{def}}{=} J \cdot N \subseteq G$. Now let us recall the easily verified fact that, for a given $p(G)$ -group, it holds that the $p(G)$ -group is *cyclic* if and only if the abelianization of the $p(G)$ -group is *cyclic*. Thus, by considering the composite $J \hookrightarrow H \twoheadrightarrow H/N$ [that determines an isomorphism $J/(J \cap N) \xrightarrow{\sim} H/N$], we conclude immediately that the image $\text{Im}(J) \subseteq (H^{(p(G))})^{\text{ab}}$ of J in $(H^{(p(G))})^{\text{ab}}$ is *not cyclic*. In particular, it follows immediately from Lemma 3.1, (i), that the image of $\text{Im}(J) \subseteq (H^{(p(G))})^{\text{ab}}$ in $(H^{(p(G))})^{\text{ab}/\text{tor}}$ is *nontrivial*. This completes the proof of assertion (iii).

Finally, we verify assertion (iv). The implication (b) \Rightarrow (a) is immediate. Next, we verify the implication (a) \Rightarrow (b). Suppose that the condition (a) is satisfied. If condition (1) of assertion (i) is satisfied, then the condition (b) is immediate. On the other hand, if condition (2) of assertion (i) is satisfied, then the condition (b) follows from assertion (iii). This completes the proof of assertion (iv), hence also of Lemma 3.3. \square

REMARK 3.3.1. — Let us give an example that satisfies each of the three conditions in Lemma 3.3, (i):

(i) One verifies easily that G itself satisfies condition (1) of Lemma 3.3, (i), i.e., that condition (1) of Lemma 3.3, (i), in the case where we take the “ J ” to be G is always satisfied.

(ii) Next, let us verify that condition (2) of Lemma 3.3, (i), in the case where we take the “ J ” to be the normal closed subgroup $P(G) \subseteq G$ of G is always satisfied. Indeed, this follows from [12], Proposition 7.5.1, together with [3], Proposition 3.6.

(iii) Finally, one verifies easily that condition (3) of Lemma 3.3, (i), in the case where we take the “ J ” to be the kernel of the natural surjection $G \twoheadrightarrow G^{(p(G))}$ is always satisfied [cf. also [3], Lemma 1.5, (i)].

PROPOSITION 3.4. — *For each $\square \in \{\circ, \bullet\}$, let G_\square be a profinite group of MLF-type. Let*

$$\alpha: G_\circ \longrightarrow G_\bullet$$

*be an **open** homomorphism. Then the following hold:*

(i) *The open homomorphism α fits into a **commutative diagram** of profinite groups*

$$\begin{array}{ccccc} P(G_\circ) & \xrightarrow{\subseteq} & I(G_\circ) & \xrightarrow{\subseteq} & G_\circ \\ \downarrow & & \downarrow & & \alpha \downarrow \\ P(G_\bullet) & \xrightarrow{\subseteq} & I(G_\bullet) & \xrightarrow{\subseteq} & G_\bullet \end{array}$$

— *where the horizontal arrows are the natural inclusions, and the vertical arrows are **open**. If, moreover, α is **surjective**, then the vertical arrows are **surjective**.*

(ii) *In the resulting [cf. (i)] commutative diagram of profinite groups*

$$\begin{array}{ccccc} G_\circ & \longrightarrow & G_\circ/P(G_\circ) & \longrightarrow & G_\circ/I(G_\circ) \\ \downarrow & & \downarrow & & \downarrow \\ G_\bullet & \longrightarrow & G_\bullet/P(G_\bullet) & \longrightarrow & G_\bullet/I(G_\bullet) \end{array}$$

— *where the horizontal arrows are the natural surjections — the middle and right-hand vertical arrows are **open injections**. In particular, if, moreover, α is **surjective**, then the middle and right-hand vertical arrows are **isomorphisms**.*

(iii) *It holds that*

$$p(G_\circ) = p(G_\bullet), \quad d(G_\circ) \geq d(G_\bullet), \quad f(G_\circ) \in f(G_\bullet)\mathbb{Z}, \quad e(G_\circ) \geq e(G_\bullet).$$

*If, moreover, α is **surjective**, then*

$$f(G_\circ) = f(G_\bullet).$$

(iv) *The right-hand vertical arrow of the diagram of (ii) maps $\text{Frob}(G_\circ) \in G_\circ/I(G_\circ)$ to $\text{Frob}(G_\bullet)^{f(G_\circ)/f(G_\bullet)} \in G_\bullet/I(G_\bullet)$ [cf. (iii)]. In particular, if, moreover, α is **surjective**, then the right-hand vertical arrow of the diagram of (ii) maps $\text{Frob}(G_\circ) \in G_\circ/I(G_\circ)$ to $\text{Frob}(G_\bullet) \in G_\bullet/I(G_\bullet)$ [cf. (iii)].*

PROOF. — Let us first observe that it follows immediately from [3], Proposition 3.6, and [3], Proposition 3.9, that, to verify Proposition 3.4, we may assume without loss of generality, by replacing G_\bullet by the image of α [which is *of MLF-type* — cf. the discussion following [3], Proposition 3.3], that α is *surjective*.

First, we verify assertions (i), (ii). The assertion that α restricts to a *surjection* $P(G_\circ) \twoheadrightarrow P(G_\bullet)$, as well as the assertion that the resulting homomorphism $G_\circ/P(G_\circ) \rightarrow$

$G_\bullet/P(G_\bullet)$ is an *isomorphism*, follows immediately — in light of [3], Proposition 3.6 — from [7], Proposition 3.4. In particular, the assertion that α restricts to a *surjection* $I(G_\circ) \twoheadrightarrow I(G_\bullet)$, as well as the assertion that the resulting homomorphism $G_\circ/I(G_\circ) \rightarrow G_\bullet/I(G_\bullet)$ is an *isomorphism*, follows immediately from Lemma 3.1, (ii). This completes the proofs of assertions (i), (ii).

Next, we verify assertion (iii). Let us first observe that the surjection α induces a *surjection* $G_\circ^{\text{ab/tor}}/(p(G_\bullet) \cdot G_\circ^{\text{ab/tor}}) \twoheadrightarrow G_\bullet^{\text{ab/tor}}/(p(G_\bullet) \cdot G_\bullet^{\text{ab/tor}})$. Thus, it holds that $p(G_\circ) = p(G_\bullet)$ and $d(G_\circ) \geq d(G_\bullet)$. In particular, it follows — in light of assertion (ii) — from Lemma 3.1, (iii), that $f(G_\circ) = f(G_\bullet)$, which thus implies that $e(G_\circ) \geq e(G_\bullet)$. This completes the proof of assertion (iii). Assertion (iv) follows from assertions (ii), (iii). This completes the proof of Proposition 3.4. \square

PROPOSITION 3.5. — *In the situation of Proposition 3.4, write $H_\bullet \subseteq G_\bullet$ for the image of α [which is of MLF-type — cf. the discussion following [3], Proposition 3.3]:*

$$\alpha: G_\circ \twoheadrightarrow H_\bullet \hookrightarrow G_\bullet.$$

Then the following hold:

(i) *The open homomorphism α determines a **commutative diagram** of topological monoids*

$$\begin{array}{ccccccccc} \mathcal{O}^\prec(G_\circ) & \xrightarrow{\subseteq} & \mathcal{O}^\times(G_\circ) & \xrightarrow{\subseteq} & \mathcal{O}^\triangleright(G_\circ) & \xrightarrow{\subseteq} & k^\times(G_\circ) & \xrightarrow{\text{rec}(G_\circ)} & G_\circ^{\text{ab}} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathcal{O}^\prec(H_\bullet) & \xrightarrow{\subseteq} & \mathcal{O}^\times(H_\bullet) & \xrightarrow{\subseteq} & \mathcal{O}^\triangleright(H_\bullet) & \xrightarrow{\subseteq} & k^\times(H_\bullet) & \xrightarrow{\text{rec}(H_\bullet)} & H_\bullet^{\text{ab}} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \mathcal{O}^\prec(G_\bullet) & \xrightarrow{\subseteq} & \mathcal{O}^\times(G_\bullet) & \xrightarrow{\subseteq} & \mathcal{O}^\triangleright(G_\bullet) & \xrightarrow{\subseteq} & k^\times(G_\bullet) & \xrightarrow{\text{rec}(G_\bullet)} & G_\bullet^{\text{ab}} \end{array}$$

— where the horizontal arrows are the natural inclusions, the upper vertical arrows are the **surjections** induced by α , and the lower vertical arrows are the **injections** determined by the transfer map [i.e., with respect to $H_\bullet \subseteq G_\bullet$] [cf. [3], Lemma 1.7, (3)].

(ii) *The left-hand upper and left-hand lower squares of the diagram of (i) determine **homomorphisms** of modules*

$$\underline{k}^\times(G_\circ) \xrightarrow{\sim} \underline{k}^\times(H_\bullet) \hookrightarrow \underline{k}^\times(G_\bullet)$$

— where the first arrow is an **isomorphism**, and the second arrow is **injective**.

(iii) The vertical open homomorphisms $\mathcal{O}^\times(G_\circ) \twoheadrightarrow \mathcal{O}^\times(H_\bullet) \hookrightarrow \mathcal{O}^\times(G_\bullet)$ in the diagram of (i) fit into a **commutative diagram** of topological modules

$$\begin{array}{ccccc} \mathcal{O}^\times(G_\circ) & \longrightarrow & \mathcal{I}(G_\circ) & \xrightarrow{\subseteq} & k_+(G_\circ) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{O}^\times(H_\bullet) & \longrightarrow & \mathcal{I}(H_\bullet) & \xrightarrow{\subseteq} & k_+(H_\bullet) \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{O}^\times(G_\bullet) & \longrightarrow & \mathcal{I}(G_\bullet) & \xrightarrow{\subseteq} & k_+(G_\bullet) \end{array}$$

— where the horizontal arrows are the natural homomorphisms, the upper vertical arrows are **surjective**, and the lower vertical arrows are **injective**.

PROOF. — These assertions follow immediately from Proposition 3.4, (i), (iii), (iv). \square

REMARK 3.5.1. — It follows immediately from Proposition 3.4, (iv), that the vertical surjection $k^\times(G_\circ) \twoheadrightarrow k^\times(H_\bullet)$ in the diagram of Proposition 3.5, (i) fits into a *commutative diagram* of modules

$$\begin{array}{ccc} k^\times(G_\circ) & \xrightarrow{\text{ord}_{\mathbb{Z}}(G_\circ)} & \mathbb{Z}_+ \\ \downarrow & & \parallel \\ k^\times(H_\bullet) & \xrightarrow{\text{ord}_{\mathbb{Z}}(H_\bullet)} & \mathbb{Z}_+ \end{array}$$

[cf. Definition 2.2].

THEOREM 3.6. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a profinite group of MLF-type. Let

$$\alpha: G_\circ \longrightarrow G_\bullet$$

be an **open** homomorphism. Suppose that $d(G_\circ) \leq d(G_\bullet)$ [which is the case if, for instance, $d(G_\circ) = 1$]. Then α is an **isomorphism**.

PROOF. — Since $d(G_\circ) \leq d(G_\bullet)$, by applying Proposition 3.4, (iii), to the natural surjection $G_\circ \twoheadrightarrow \alpha(G_\circ)$ and the natural inclusion $\alpha(G_\circ) \hookrightarrow G_\bullet$ [note that $\alpha(G_\circ)$ is of MLF-type — cf. the discussion following [3], Proposition 3.3], we obtain that $d(\alpha(G_\circ)) = d(G_\bullet)$. On the other hand, it follows from [3], Proposition 3.6, that this equality implies the equality $\alpha(G_\circ) = G_\bullet$, i.e., that α is *surjective*.

Now assume that α is *not injective*, i.e., that $J \stackrel{\text{def}}{=} \text{Ker}(\alpha)$ is *nontrivial*. Let us first observe that since J is contained in $P(G_\circ)$ [cf. Proposition 3.4, (ii)], the profinite group J is *pro- $p(G_\circ)$* , which thus implies that J does *not satisfy* condition (3) of Lemma 3.3, (i). Thus, it follows from Lemma 3.3, (iv), that there exists an open subgroup $H_\circ \subseteq G_\circ$ of G_\circ such that $J \subseteq H_\circ$ [i.e., $H_\circ = \alpha^{-1}(\alpha(H_\circ))$], and, moreover, the image of J in $(H_\circ^{(p(G_\circ))})^{\text{ab/tor}}$ is *nontrivial*. In particular, since $d(H_\circ) = d(G_\circ) \cdot [G_\circ : H_\circ] \leq d(G_\bullet) \cdot [G_\circ : H_\circ] = d(G_\bullet) \cdot [G_\bullet : \alpha(H_\circ)] = d(\alpha(H_\circ))$ [cf. [3], Proposition 3.6], we may assume without loss of generality, by replacing (G_\circ, G_\bullet) by $(H_\circ, \alpha(H_\circ))$, that the image of J in $(G_\circ^{(p(G_\circ))})^{\text{ab/tor}}$ is *nontrivial*. On the other hand, this implies that the surjection

$(G_{\circ}^{(p(G_{\circ}))})_{\text{ab/tor}} \rightarrow (G_{\bullet}^{(p(G_{\circ}))})_{\text{ab/tor}} = (G_{\bullet}^{(p(G_{\bullet}))})_{\text{ab/tor}}$ [cf. Proposition 3.4, (iii)] induced by α is *not injective*. Thus, it follows immediately from Lemma 3.1, (i), that $d(G_{\circ}) > d(G_{\bullet})$ — in *contradiction* to our assumption that $d(G_{\circ}) \leq d(G_{\bullet})$. This completes the proof of Theorem 3.6. \square

COROLLARY 3.7. — For each $\square \in \{\circ, \bullet\}$, let G_{\square} be a profinite group of MLF-type. Let

$$\alpha: G_{\circ} \longrightarrow G_{\bullet}$$

be an **open** homomorphism. Suppose that $e(G_{\circ}) \leq e(G_{\bullet})$ [which is the case if, for instance, $e(G_{\circ}) = 1$]. Then α is **injective**.

PROOF. — Since $e(G_{\circ}) \leq e(G_{\bullet})$, by applying Proposition 3.4, (iii), to the natural surjection $G_{\circ} \twoheadrightarrow \alpha(G_{\circ})$ and the natural inclusion $\alpha(G_{\circ}) \hookrightarrow G_{\bullet}$ [note that $\alpha(G_{\circ})$ is of MLF-type — cf. the discussion following [3], Proposition 3.3], we obtain that $e(G_{\circ}) = e(\alpha(G_{\circ}))$. Thus, to verify Corollary 3.7, we may assume without loss of generality, by replacing G_{\bullet} by $\alpha(G_{\circ})$, that α is *surjective*, and that $e(G_{\circ}) = e(G_{\bullet})$. Then since α is *surjective*, and $e(G_{\circ}) = e(G_{\bullet})$, it follows immediately from Proposition 3.4, (iii), that $d(G_{\circ}) = d(G_{\bullet})$. Thus, it follows from Theorem 3.6 that α is an *isomorphism*, as desired. This completes the proof of Corollary 3.7. \square

COROLLARY 3.8. — For each $\square \in \{\circ, \bullet\}$, let k_{\square} be an MLF and \bar{k}_{\square} an algebraic closure of k_{\square} ; write $G_{\square} \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_{\square}/k_{\square})$. Suppose that $e_{k_{\circ}} = 1$. Then the following three conditions are equivalent:

- (1) The field k_{\circ} is **isomorphic** to the field k_{\bullet} .
- (2) There exists a **surjection** $G_{\circ} \twoheadrightarrow G_{\bullet}$.
- (3) The group G_{\circ} is **isomorphic** to the group G_{\bullet} .

PROOF. — The implication (1) \Rightarrow (2) is immediate. The implication (2) \Rightarrow (3) follows — in light of [3], Proposition 3.6 — from Corollary 3.7. Finally, since [we have assumed that] $e_{k_{\circ}} = 1$, the implication (3) \Rightarrow (1) follows immediately from [3], Proposition 3.6 [cf. also [3], Lemma 1.5, (i)]. This completes the proof of Corollary 3.8. \square

4. RECONSTRUCTION ALGORITHMS RELATED TO ABSOLUTELY ABELIAN MLF'S

In the present §4, we maintain the notational conventions introduced at the beginnings of §1 and §2. In the present §4, we discuss some *group-theoretic reconstruction algorithms* [cf. [8], Remark 1.9.8] related to *absolutely abelian* [cf. Definition 4.2, (ii), below] MLF's. We establish, for instance, a group-theoretic reconstruction algorithm for constructing, from a group of MLF-type, a homomorphism which “corresponds” to the Norm map $\text{Nm}_{k/k^{(d=1)}}: k^{\times} \rightarrow (k^{(d=1)})^{\times}$ with respect to the finite extension $k/k^{(d=1)}$ [cf. Definition 4.7, (iii); Proposition 4.9, (i), below], which leads us to the notion of *MLF-Galois label* [cf. Definition 4.10, Theorem 4.11 below]. Finally, as a consequence of the group-theoretic

reconstruction algorithms, we also obtain a *refinement* of the main theorem of [6] [cf. Corollary 4.14, (i); Remark 4.14.1 below].

LEMMA 4.1. — *The following hold:*

(i) *The natural homomorphisms*

$$\mathbb{Z}_{p_k} \longrightarrow \text{End}_{\mathbb{Z}_{p_k}}(\Lambda(\bar{k})^{(p_k)}), \quad \mathbb{Q}_{p_k} \longrightarrow \text{End}_{\mathbb{Q}_{p_k}}(\Lambda(\bar{k})^{(p_k)} \otimes_{\mathbb{Z}_{p_k}} \mathbb{Q}_{p_k})$$

are isomorphisms of topological algebras. Moreover, these isomorphisms restrict to isomorphisms of topological groups

$$\mathbb{Z}_{p_k}^\times \xrightarrow{\sim} \text{Aut}_{\mathbb{Z}_{p_k}}(\Lambda(\bar{k})^{(p_k)}), \quad \mathbb{Q}_{p_k}^\times \xrightarrow{\sim} \text{Aut}_{\mathbb{Q}_{p_k}}(\Lambda(\bar{k})^{(p_k)} \otimes_{\mathbb{Z}_{p_k}} \mathbb{Q}_{p_k}),$$

respectively.

We shall write

$$\chi_{p_k\text{-cyc}}: G_k \longrightarrow \mathbb{Z}_{p_k}^\times$$

for the composite of the natural action $G_k \rightarrow \text{Aut}_{\mathbb{Z}_{p_k}}(\Lambda(\bar{k})^{(p_k)})$ and the above isomorphism $\text{Aut}_{\mathbb{Z}_{p_k}}(\Lambda(\bar{k})^{(p_k)}) \xrightarrow{\sim} \mathbb{Z}_{p_k}^\times$, i.e., the p_k -adic cyclotomic character.

(ii) *Let $\overline{\mathbb{Q}}_{p_k}$ be an algebraic closure of \mathbb{Q}_{p_k} . Then the homomorphism $G_k^{\text{ab}} \rightarrow \mathbb{Z}_{p_k}^\times$ determined by the p_k -adic cyclotomic character $\chi_{p_k\text{-cyc}}: G_k \rightarrow \mathbb{Z}_{p_k}^\times$ [cf. (i)] **coincides** with the composite*

$$G_k^{\text{ab}} \rightarrow \text{Gal}(\bar{k}/k^{(d=1)})^{\text{ab}} \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/\mathbb{Q}_{p_k})^{\text{ab}} \xleftarrow{\text{rec}_{\mathbb{Q}_{p_k}}^\wedge} (\mathbb{Q}_{p_k}^\times)^\wedge = \mathbb{Z}_{p_k}^\times \times \widehat{p_k} \rightarrow \mathbb{Z}_{p_k}^\times \xrightarrow{\sim} \mathbb{Z}_{p_k}^\times$$

— where the first arrow “ \rightarrow ” is the homomorphism induced by the natural inclusion $G_k \hookrightarrow \text{Gal}(\bar{k}/k^{(d=1)})$; the second arrow “ $\xrightarrow{\sim}$ ” is the isomorphism induced by an isomorphism $\overline{\mathbb{Q}}_{p_k} \xrightarrow{\sim} \bar{k}$ of fields [that necessarily restricts to an isomorphism $\mathbb{Q}_{p_k} \xrightarrow{\sim} k^{(d=1)}$ of fields]; the third arrow $\text{rec}_{\mathbb{Q}_{p_k}}^\wedge$ is the isomorphism in the final display of [3], Lemma 1.7, (1) [in the case where we take the “ k ” of [3], Lemma 1.7, to be \mathbb{Q}_{p_k}]; the fourth arrow “ \rightarrow ” is the first projection; the fifth arrow “ $\xrightarrow{\sim}$ ” is the isomorphism given by “ $a \mapsto a^{-1}$ ”.

(iii) *The composite of the **Norm map** $\text{Nm}_{k/k^{(d=1)}}: k^\times \rightarrow (k^{(d=1)})^\times$ with respect to the finite extension $k/k^{(d=1)}$ and the isomorphism $(k^{(d=1)})^\times \xrightarrow{\sim} \mathbb{Q}_{p_k}^\times$ induced by the [uniquely determined] isomorphism $k^{(d=1)} \xrightarrow{\sim} \mathbb{Q}_{p_k}$ of fields **coincides** with the homomorphism $k^\times \rightarrow \mathbb{Q}_{p_k}^\times$ given by*

$$k^\times \ni a \mapsto \chi_{p_k\text{-cyc}}(\text{rec}_k(a))^{-1} \cdot p_k^{f_k \cdot \text{ord}_k(a)} \in \mathbb{Q}_{p_k}^\times$$

[cf. [3], Lemma 1.7].

PROOF. — Assertion (i) follows from the [easily verified] fact that the \mathbb{Z}_{p_k} -module $\Lambda(\bar{k})^{(p_k)}$ is *free of rank one*. Assertion (ii) follows immediately from the well-known [cf., e.g., [15], Chapter III, §A.4, Corollary] fact that the p_k -adic cyclotomic character in the case where we take the “ k ” to be the MLF \mathbb{Q}_{p_k} *coincides* with the “*Lubin-Tate character* $\chi_{\sigma, \pi}^{\text{LT}}$ ” [cf. the notational convention introduced in [1], Definition 1.2, (ii)] in the case where we take the “ E ” (respectively, “ σ ”; “ π ”) of [1], Definition 1.2, (ii), to be \mathbb{Q}_{p_k} (respectively, the identity automorphism of \mathbb{Q}_{p_k} ; $p_k \in \mathcal{O}_{\mathbb{Q}_{p_k}} = \mathbb{Z}_{p_k}$). Assertion (iii) follows immediately from assertion (ii) and [3], Lemma 1.7, (1), (2). This completes the proof of Lemma 4.1. \square

DEFINITION 4.2.

- (i) We shall say that the MLF k is *absolutely Galois* if k is Galois over $k^{(d=1)}$.
- (ii) We shall say that the MLF k is *absolutely abelian* if k is absolutely Galois, and, moreover, the Galois group $\text{Gal}(k/k^{(d=1)})$ is abelian.
- (iii) We shall write $k^{(\text{ab})} \subseteq k$ for the [uniquely determined] maximal absolutely abelian MLF contained in k .
- (iv) We shall write

$$d_k^{(\text{ab})} \stackrel{\text{def}}{=} d_{k^{(\text{ab})}}, \quad e_k^{(\text{ab})} \stackrel{\text{def}}{=} e_{k^{(\text{ab})}}$$

for the “ d_k ”, “ e_k ” in the case where we take the “ k ” to be $k^{(\text{ab})}$ of (iii), respectively.

LEMMA 4.3. — *Let K be an intermediate field of the finite extension $k/k^{(\text{ab})}$. Then the following hold:*

- (i) *It holds that $K^{(\text{ab})} = k^{(\text{ab})}$.*
- (ii) *There is **no nontrivial** intermediate field of the finite extension $k/k^{(\text{ab})}$, hence also of $K/k^{(\text{ab})}$, that is **unramified** over $k^{(\text{ab})}$.*

PROOF. — Assertion (i) follows from the definition of “ $(-)^{(\text{ab})}$ ”. Assertion (ii) follows immediately from the [easily verified] fact that every intermediate field of $k/k^{(\text{ab})}$ *unramified* over $k^{(\text{ab})}$ is *absolutely abelian* [cf. [3], Lemma 1.5, (i)]. This completes the proof of Lemma 4.3. \square

LEMMA 4.4. — *The following hold:*

- (i) *It holds that*

$$d_k^{(\text{ab})} = e_k^{(\text{ab})} \cdot f_k, \quad d_k^{(\text{ab})} = [(k^{(d=1)})^\times : \text{Nm}_{k/k^{(d=1)}}(k^\times)].$$

- (ii) *The following three conditions are equivalent:*

- (1) *The MLF k is **absolutely abelian**.*
- (2) *It holds that $d_k = d_k^{(\text{ab})}$.*
- (3) *It holds that $e_k = e_k^{(\text{ab})}$.*

PROOF. — First, we verify assertion (i). The equality $d_k^{(\text{ab})} = e_k^{(\text{ab})} \cdot f_k$ follows from Lemma 4.3, (ii), together with [3], Lemma 1.2, (iii). The equality $d_k^{(\text{ab})} = [(k^{(d=1)})^\times : \text{Nm}_{k/k^{(d=1)}}(k^\times)]$ follows immediately from [3], Lemma 1.7, (1), (2). This completes the proof of assertion (i). Assertion (ii) follows immediately from assertion (i), together with [3], Lemma 1.2, (iii). This completes the proof of Lemma 4.4. \square

Recall the group of *MLF-type*

G

introduced at the beginning of §2.

DEFINITION 4.5.

(i) We shall write

$$\Lambda(G)^{(p(G))}$$

for the maximal pro- $p(G)$ quotient of the cyclotome $\Lambda(G)$ associated to G . Note that since $\Lambda(G)^{(p(G))}$ has a natural structure of free $\mathbb{Z}_{p(G)}$ -module of rank one [cf. [3], Proposition 4.2, (iv)], the perfection

$$\left(\Lambda(G)^{(p(G))}\right)^{\text{pf}}$$

of $\Lambda(G)^{(p(G))}$ has a natural structure of $\mathbb{Q}_{p(G)}$ -vector space of dimension one.

(ii) We shall write

$$\mathbb{Z}_p(G) \stackrel{\text{def}}{=} \text{End}\left(\Lambda(G)^{(p(G))}\right)$$

for the topological algebra of endomorphisms of the topological module $\Lambda(G)^{(p(G))}$.

(iii) We shall write

$$\mathbb{Q}_p(G) \stackrel{\text{def}}{=} \text{End}\left(\left(\Lambda(G)^{(p(G))}\right)^{\text{pf}}\right)$$

for the algebra of endomorphisms of the perfection $\left(\Lambda(G)^{(p(G))}\right)^{\text{pf}}$. Thus, we have a natural inclusion

$$\mathbb{Z}_p(G) \hookrightarrow \mathbb{Q}_p(G).$$

By considering the topology induced by the topology of $\mathbb{Z}_p(G)$ [cf. (ii)], we regard $\mathbb{Q}_p(G)$ as a topological algebra.

LEMMA 4.6. — *The following hold:*(i) *The natural homomorphism*

$$\mathbb{Z}_{p(G)} \longrightarrow \mathbb{Z}_p(G)$$

[i.e., obtained by the natural $\mathbb{Z}_{p(G)}$ -module structure of $\Lambda(G)^{p(G)}$] is an isomorphism of topological algebra. Moreover, this isomorphism determines an isomorphism of topological algebra

$$\mathbb{Q}_{p(G)} \xrightarrow{\sim} \mathbb{Q}_p(G).$$

(ii) *We have natural identifications*

$$\mathbb{Z}_p(G)^\times = \text{Aut}\left(\Lambda(G)^{p(G)}\right) \subseteq \mathbb{Q}_p(G)^\times = \text{Aut}\left(\left(\Lambda(G)^{p(G)}\right)^{\text{pf}}\right)$$

[cf. (i)].

PROOF. — These assertions follow immediately — in light of [3], Proposition 3.6; [3], Proposition 4.2, (iv) — from Lemma 4.1, (i). \square

DEFINITION 4.7.

(i) We shall write

$$\chi_{p\text{-cyc}}(G): G \longrightarrow \mathbb{Z}_p(G)^\times$$

for the natural action of G on $\Lambda(G)^{p(G)}$ [cf. Lemma 4.6, (ii)].

(ii) We shall write

$$p_{\in}(G) \in \mathbb{Q}_p(G)^{\times}$$

for the automorphism of $(\Lambda(G)^{p(G)})^{\text{pf}}$ given by multiplication by $p(G)$ [cf. Lemma 4.6, (ii)].

(iii) We shall write

$$\text{Nm}_{\text{abs}}(G): k^{\times}(G) \longrightarrow \mathbb{Q}_p(G)^{\times}$$

for the homomorphism of topological modules defined by

$$k^{\times}(G) \ni a \mapsto \chi_{p\text{-cyc}}(G)(\text{rec}(G)(a))^{-1} \cdot p_{\in}(G)^{f(G) \cdot \text{ord}_{\mathbb{Z}}(G)(a)} \in \mathbb{Q}_p(G)^{\times}$$

[cf. Definition 2.2].

(iv) We shall write

$$d^{(\text{ab})}(G) \stackrel{\text{def}}{=} \left[\mathbb{Q}_p(G)^{\times} : \text{Im}(\text{Nm}_{\text{abs}}(G)) \right], \quad e^{(\text{ab})}(G) \stackrel{\text{def}}{=} d^{(\text{ab})}(G)/f(G).$$

DEFINITION 4.8. — We shall say that G is of *AAMLF-type* if $d(G) = d^{(\text{ab})}(G)$. [Here, “AAMLF” is to be understood as an abbreviation for “absolutely abelian mixed-characteristic local field” — cf. Proposition 4.9, (iii), below.]

PROPOSITION 4.9. — *The following hold:*

(i) Write $\text{Nm}_{k/k^{(d=1)}}: k^{\times} \rightarrow (k^{(d=1)})^{\times}$ for the Norm map with respect to the finite extension $k/k^{(d=1)}$. Then the vertical isomorphism $k^{\times} \xrightarrow{\sim} k^{\times}(G_k)$ in the diagram of [3], Proposition 3.11, (i), fits into a **commutative diagram** of topological modules

$$\begin{array}{ccc} k^{\times} & \xrightarrow{\text{Nm}_{k/k^{(d=1)}}} & (k^{(d=1)})^{\times} \\ \wr \downarrow & & \wr \downarrow \\ k^{\times}(G_k) & \xrightarrow{\text{Nm}_{\text{abs}}(G_k)} & \mathbb{Q}_p(G_k)^{\times} \end{array}$$

— where the right-hand vertical arrow is the composite of the isomorphism $(k^{(d=1)})^{\times} \xrightarrow{\sim} \mathbb{Q}_p(G_k)^{\times}$ induced by the [uniquely determined] isomorphism $k^{(d=1)} \xrightarrow{\sim} \mathbb{Q}_p(G_k)$ of fields and the isomorphism $\mathbb{Q}_p(G_k)^{\times} = \mathbb{Q}_{p_k}^{\times} \xrightarrow{\sim} \mathbb{Q}_p(G_k)^{\times}$ [cf. [3], Proposition 3.6] determined by the isomorphism of Lemma 4.6, (i).

(ii) It holds that

$$d_k^{(\text{ab})} = d^{(\text{ab})}(G_k), \quad e_k^{(\text{ab})} = e^{(\text{ab})}(G_k).$$

(iii) It holds that the MLF k is **absolutely abelian** if and only if the group G_k is of **AAMLF-type**.

PROOF. — Assertion (i) follows — in light of Proposition 2.3; [3], Proposition 3.6; [3], Proposition 3.11, (i); [3], Proposition 4.2, (iv) — from Lemma 4.1, (iii). Assertion (ii) follows immediately from Lemma 4.4, (i), together with assertion (i) and [3], Proposition 3.6. Assertion (iii) follows from Lemma 4.4, (ii), together with assertion (ii) and [3], Proposition 3.6. This completes the proof of Proposition 4.9. \square

REMARK 4.9.1. — Let $H \subseteq G$ be an open subgroup of G . Then one verifies immediately from Proposition 4.9, (i), together with [3], Lemma 1.7, (2), that the diagram of topological modules

$$\begin{array}{ccc} k(H)^\times & \xrightarrow{\text{Nm}_{\text{abs}}(H)} & \mathbb{Q}_p(H)^\times \\ \downarrow & & \parallel \\ k(G)^\times & \xrightarrow{\text{Nm}_{\text{abs}}(G)} & \mathbb{Q}_p(G)^\times \end{array}$$

— where the left-hand vertical arrow is the homomorphism induced by the homomorphism $H^{\text{ab}} \rightarrow G^{\text{ab}}$ determined by the inclusion $H \hookrightarrow G$, and the right-hand vertical arrow is the composite of the isomorphisms $\mathbb{Q}_{p(G)}^\times \xrightarrow{\sim} \mathbb{Q}_p(G)^\times$, $\mathbb{Q}_{p(H)}^\times \xrightarrow{\sim} \mathbb{Q}_p(H)^\times$ determined by the isomorphism of Lemma 4.6, (i) [cf. also [3], Proposition 3.6] — *commutes*.

REMARK 4.9.2. — Suppose that G is of *AAMLF-type*. Then a profinite group *isomorphic* to G may be constructed as follows: Let \tilde{G} be a group of *MLF-type* such that $(p(\tilde{G}), d(\tilde{G})) = (p(G), 1)$. [Note that one verifies easily from [3], Proposition 3.6, that this condition $(p(\tilde{G}), d(\tilde{G})) = (p(G), 1)$ *completely determines* the isomorphism class of the group \tilde{G} .] Write $J \subseteq \tilde{G}^{\text{ab}}$ for the closure, i.e. in \tilde{G}^{ab} , of the inverse image ($\subseteq k^\times(\tilde{G}) \subseteq \tilde{G}^{\text{ab}}$) of $\text{Im}(\text{Nm}_{\text{abs}}(G)) \subseteq \mathbb{Q}_p(G)^\times$ by $\text{Nm}_{\text{abs}}(\tilde{G}): k^\times(\tilde{G}) \rightarrow \mathbb{Q}_p(\tilde{G})^\times$ — relative to the composite of the isomorphisms $\mathbb{Q}_{p(G)}^\times \xrightarrow{\sim} \mathbb{Q}_p(G)^\times$, $\mathbb{Q}_{p(\tilde{G})}^\times \xrightarrow{\sim} \mathbb{Q}_p(\tilde{G})^\times$ determined by the isomorphism of Lemma 4.6, (i). Then it follows immediately from Remark 4.9.1 that G is *isomorphic*, as an abstract profinite group, to the inverse image of $J \subseteq \tilde{G}^{\text{ab}}$ in \tilde{G} .

DEFINITION 4.10. — We shall refer to the collection of data

$$\left(p(G), d(G), \text{Im}((\text{Nm}_{\text{abs}})(G)) \subseteq \mathbb{Q}_p(G)^\times \xleftarrow{\sim} \mathbb{Q}_{p(G)}^\times \right)$$

[cf. Lemma 4.6, (i)] consisting of the prime number $p(G)$, the positive integer $d(G)$, and the open subgroup $\text{Im}((\text{Nm}_{\text{abs}})(G)) \subseteq \mathbb{Q}_{p(G)}^\times$ of $\mathbb{Q}_{p(G)}^\times$ as the *MLF-Galois label* of G .

THEOREM 4.11. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a group of **MLF-type**. Suppose that one of the following two conditions is satisfied:

- (1) It holds that $\{(p(G_\circ), a(G_\circ)), (p(G_\bullet), a(G_\bullet))\} \not\subseteq \{(2, 1)\}$ [cf. Definition 2.4, (ii)].
- (2) Either G_\circ or G_\bullet is of **AAMLF-type**.

Then it holds that the group G_\circ is **isomorphic** to the group G_\bullet if and only if the *MLF-Galois label* of G_\circ **coincides** with the *MLF-Galois label* of G_\bullet .

PROOF. — The *necessity* is immediate. Next, we verify the *sufficiency* in the case where condition (1) is satisfied. Suppose that condition (1) is satisfied, and that the *MLF-Galois label* of G_\circ *coincides* with the *MLF-Galois label* of G_\bullet . Then since $\text{Im}(\text{Nm}_{\text{abs}}(G_\circ)) = \text{Im}(\text{Nm}_{\text{abs}}(G_\bullet))$, one verifies immediately from Proposition 2.5, (i); Proposition 4.9, (i); [3], Lemma 1.7, (1), (2); [3], Proposition 3.6, that $(2, 1) \notin \{(p(G_\circ), a(G_\circ)), (p(G_\bullet), a(G_\bullet))\}$. Thus, since the *MLF-Galois label* of G_\circ *coincides* with the *MLF-Galois label* of G_\bullet , it

follows immediately — in light of Proposition 2.5, (i); Proposition 4.9, (i); [3], Proposition 3.6 — from the main theorems of [4] and [13], together with [3], Lemma 1.7, (1), (2), that G_\circ is *isomorphic* to G_\bullet , as desired. This completes the proof of the *sufficiency* in the case where condition (1) is satisfied

Finally, we verify the *sufficiency* in the case where condition (2) is satisfied. Suppose that G_\circ is of *AAMLF-type*, and that the MLF-Galois label of G_\circ coincides with the MLF-Galois label of G_\bullet . Then since $\text{Im}(\text{Nm}_{\text{abs}}(G_\circ)) = \text{Im}(\text{Nm}_{\text{abs}}(G_\bullet))$, we obtain that $d^{(\text{ab})}(G_\circ) = d^{(\text{ab})}(G_\bullet)$. In particular, since G_\circ is of *AAMLF-type*, the equality $d(G_\circ) = d(G_\bullet)$ implies that G_\bullet is of *AAMLF-type*. Thus, since the MLF-Galois label of G_\circ coincides with the MLF-Galois label of G_\bullet , it follows immediately from Remark 4.9.2 that G_\circ is *isomorphic* to G_\bullet , as desired. This completes the proof of the *sufficiency* in the case where condition (2) is satisfied, hence also of Theorem 4.11. \square

REMARK 4.11.1.

(i) Let us recall that the main theorem of [1] asserts that, roughly speaking, the *Hodge-Tate-ness* of p_k -adic representations of the group G_k of MLF-type is closely related to the *ring structures* of the fields $k \subseteq \bar{k}$.

(ii) Let us also recall that, as discussed in [3], Proposition 4.2, (iv), the *p_k -adic cyclotomic character* may be “reconstructed” from just the group structure of the group G_k of MLF-type.

Next, let us recall that Theorem 4.11 asserts that — under a mild assumption on “ (p_k, a_k) ” — the isomorphism class of the group G_k is *completely determined* by the *MLF-Galois label* of G_k . Now observe that the main component of the notion of MLF-Galois label is the third component, i.e., the image of the Norm map to $(k^{(d=1)})^\times$. Moreover, recall that, as discussed in Lemma 4.1, (iii), roughly speaking, the Norm map to $(k^{(d=1)})^\times$ may be essentially described by the *p_k -adic cyclotomic character*.

Thus, one may conclude that, roughly speaking, the *p_k -adic cyclotomic character* is closely related to the group structure of the group G_k of MLF-type.

(iii) It follows from the observations of (i), (ii), together with [3], Remark 4.3.3, that, in summary,

Hodge-Tate representations is closely related to *arithmetic holomorphic structures* [cf. [9], §2.7, (vii)] of MLF’s,

and, moreover,

the *cyclotomic character* [that is one of Hodge-Tate representations] is closely related to *mono-analytic structures* [cf. [9], §2.7, (vii)] of MLF’s.

Hodge-Tate representations	\iff	arithmetic holomorphic structures of MLF’s
	\implies	the main theorem of [1]
	\impliedby	immediate
cyclotomic characters	\iff	mono-analytic structures of MLF’s
	\implies	Lemma 4.1, (iii), and Theorem 4.11
	\impliedby	[3], Proposition 4.2, (iv)

DEFINITION 4.12. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a group of MLF-type. Let $\alpha: G_\circ \rightarrow G_\bullet$ be a homomorphism. Then we shall say that α is *cyclotomically compatible* if $p(G_\circ) = p(G_\bullet)$, and, moreover, the diagram

$$\begin{array}{ccc} G_\circ & \xrightarrow{\chi_{p\text{-cyc}}(G_\circ)} & \mathbb{Z}_p(G_\circ)^\times \\ \alpha \downarrow & & \wr \downarrow \\ G_\bullet & \xrightarrow{\chi_{p\text{-cyc}}(G_\bullet)} & \mathbb{Z}_p(G_\bullet)^\times \end{array}$$

— where the right-hand vertical arrow is the composite of the isomorphisms $\mathbb{Z}_{p(G_\circ)}^\times \xrightarrow{\sim} \mathbb{Z}_p(G_\circ)^\times$, $\mathbb{Z}_{p(G_\bullet)}^\times \xrightarrow{\sim} \mathbb{Z}_p(G_\bullet)^\times$ determined by the isomorphism of Lemma 4.6, (i) — commutes.

REMARK 4.12.1. — Let us recall that it follows immediately from the various definitions involved that every *open injection* between profinite groups of MLF-type induces a natural isomorphism between the *cyclotomes* “ $\Lambda(-)$ ” [cf. [3], Definition 4.1, (i), (ii), (iii)]. In particular, every *open injection* between profinite groups of MLF-type is *cyclotomically compatible*.

REMARK 4.12.2. — Let l be a prime number such that $l \neq p(G)$. Then, in the situation of Definition 4.7, (i), by considering the natural action on the maximal *pro- l* quotient of the cyclotome $\Lambda(G)$ [i.e., as opposed to the natural action on $\Lambda(G)^{p(G)}$ discussed in Definition 4.7, (i)], one may define the notion of “ *l -adic cyclotomic character*” of G [i.e., as opposed to the “ *$p(G)$ -adic cyclotomic character*” $\chi_{p\text{-cyc}}(G)$ defined in Definition 4.7, (i)].

Now let us observe that it follows immediately — in light of [3], Proposition 3.6; [3], Proposition 4.2, (iv) — from Proposition 3.4, (iii), (iv), and [3], Lemma 1.5, (i), (ii), (iii), that every *open homomorphism* between profinite groups of MLF-type is *compatible* with the respective “ *l -adic cyclotomic characters*”, i.e., a similar diagram to the diagram of Definition 4.12 *commutes*.

THEOREM 4.13. — For each $\square \in \{\circ, \bullet\}$, let G_\square be a profinite group of MLF-type. Let

$$\alpha: G_\circ \longrightarrow G_\bullet$$

be an **open homomorphism**. Then the following hold:

(i) If α is **cyclotomically compatible and surjective**, then the surjection $k^\times(G_\circ) \twoheadrightarrow k^\times(G_\bullet)$ induced by α [cf. Proposition 3.5, (i)] fits into a **commutative diagram** of topological modules

$$\begin{array}{ccc} k^\times(G_\circ) & \xrightarrow{\text{Nm}_{\text{abs}}(G_\circ)} & \mathbb{Q}_p(G_\circ)^\times \\ \downarrow & & \parallel \\ k^\times(G_\bullet) & \xrightarrow{\text{Nm}_{\text{abs}}(G_\bullet)} & \mathbb{Q}_p(G_\bullet)^\times \end{array}$$

— where the right-hand vertical arrow is the composite of the isomorphisms $\mathbb{Q}_p^\times(G_\circ) \xrightarrow{\sim} \mathbb{Q}_p(G_\bullet)^\times, \mathbb{Q}_p^\times(G_\bullet) \xrightarrow{\sim} \mathbb{Q}_p(G_\bullet)^\times$ determined by the isomorphism of Lemma 4.6, (i) [cf. also Proposition 3.4, (iii)]. Moreover, it holds that

$$d^{(\text{ab})}(G_\circ) = d^{(\text{ab})}(G_\bullet), \quad e^{(\text{ab})}(G_\circ) = e^{(\text{ab})}(G_\bullet).$$

(ii) If G_\circ is of **AAMLF-type**, then the following two conditions are equivalent:

- (1) The homomorphism α is **injective**.
- (2) The homomorphism α is **cyclotomically compatible**.

(iii) In the situation of (ii), if, moreover, (1) and (2) of (ii) are satisfied, then the group G_\bullet is of **AAMLF-type**.

PROOF. — First, we verify assertion (i). The first assertion, hence also the equality $d^{(\text{ab})}(G_\circ) = d^{(\text{ab})}(G_\bullet)$, follows immediately from Proposition 3.4, (iii), and Remark 3.5.1. Thus, the equality $e^{(\text{ab})}(G_\circ) = e^{(\text{ab})}(G_\bullet)$ follows from Proposition 3.4, (iii). This completes the proof of assertion (i).

Next, we verify assertion (ii). The implication (1) \Rightarrow (2) was already discussed in Remark 4.12.1. We verify the implication (2) \Rightarrow (1). Suppose that condition (2) is satisfied. Let us first observe that it follows from Remark 4.12.1 that, to verify the implication (2) \Rightarrow (1), we may assume without loss of generality, by replacing G_\bullet by the image of α [which is of *MLF-type* — cf. the discussion following [3], Proposition 3.3], that α is *surjective*. Thus, since [we have assumed that] $d(G_\circ) = d^{(\text{ab})}(G_\circ)$, it follows from assertion (i), together with the [easily verified] inequality $d^{(\text{ab})}(G_\bullet) \leq d(G_\bullet)$, that $d(G_\circ) \leq d(G_\bullet)$. In particular, it follows from Theorem 3.6 that α is an *isomorphism*, as desired. This completes the proof of the implication (2) \Rightarrow (1), hence also of assertion (ii).

Assertion (iii) follows — in light of Proposition 4.9, (iii) — from the [easily verified] fact that an MLF contained in an *absolutely abelian* MLF is *absolutely abelian*. This completes the proof of Theorem 4.13. \square

COROLLARY 4.14. — For each $\square \in \{\circ, \bullet\}$, let k_\square be an MLF and \bar{k}_\square an algebraic closure of k_\square ; write $G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$. Then the following hold:

(i) Suppose that there exists a **cyclotomically compatible surjection** $G_\circ \twoheadrightarrow G_\bullet$. Then the field $k_\circ^{(\text{ab})}$ is **isomorphic** to the field $k_\bullet^{(\text{ab})}$.

(ii) Suppose that k_\circ is **absolutely abelian**. Then the following three conditions are equivalent:

- (1) The field k_\circ is **isomorphic** to the field k_\bullet .
- (2) There exists a **cyclotomically compatible surjection** $G_\circ \twoheadrightarrow G_\bullet$.
- (3) The group G_\circ is **isomorphic** to the group G_\bullet .

PROOF. — Assertion (i) follows immediately — in light of Proposition 4.9, (i) — from Theorem 4.13, (i), and [3], Lemma 1.7, (1), (2). Next, we verify assertion (ii). The implication (1) \Rightarrow (2) is immediate. The implication (2) \Rightarrow (3) follows — in light of

Proposition 4.9, (iii) — from Theorem 4.13, (ii). Finally, we verify the implication (3) \Rightarrow (1). Suppose that condition (3) is satisfied. Then it follows from Proposition 4.9, (iii), that k_\bullet is *absolutely abelian*. Thus, the implication (3) \Rightarrow (1) follows from assertion (i). This completes the proof of the implication (3) \Rightarrow (1), hence also of assertion (ii). \square

REMARK 4.14.1. — The main theorem of [6] is equivalent to Corollary 4.14, (i), in the case where the surjection “ $G_\circ \rightarrow G_\bullet$ ” is an *isomorphism*. Now let us recall that it is immediate that every *isomorphism* between groups of MLF-type is *cyclotomically compatible*. Thus, Corollary 4.14, (i), may be regarded as a *refinement* of the main theorem of [6].

Some of the group-theoretic reconstruction algorithms discussed in the present §4 may be summarized as follows.

SUMMARY 4.15.

(i) *There exist group-theoretic reconstruction algorithms [cf. [8], Remark 1.9.8] for constructing, from a group G of MLF-type,*

- *topological rings $\mathbb{Z}_p(G) \subseteq \mathbb{Q}_p(G)$ [cf. Definition 4.5, (ii), (iii)],*
- *a homomorphism $\text{Nm}_{\text{abs}}(G): k^\times(G) \rightarrow \mathbb{Q}_p(G)^\times$ of topological modules [cf. Definition 4.7, (iii)], and*
- *integers $d^{(\text{ab})}(G)$ and $e^{(\text{ab})}(G)$ [cf. Definition 4.7, (iv)]*

which “correspond” to

- *the topological rings $\mathbb{Z}_{p_k} \subseteq \mathbb{Q}_{p_k}$ [cf. Lemma 4.6, (i)],*
- *the Norm map $\text{Nm}_{k/k^{(d=1)}}: k^\times \rightarrow (k^{(d=1)})^\times$ with respect to the finite extension $k/k^{(d=1)}$ [cf. Proposition 4.9, (i)], and*
- *the integers $d_k^{(\text{ab})}$ and $e_k^{(\text{ab})}$ [cf. Proposition 4.9, (ii)],*

respectively.

(ii) *There exists a group-theoretic condition for a group of MLF-type [cf. Definition 4.8] which “corresponds” to the condition for an MLF to be absolutely abelian [cf. Proposition 4.9, (iii)].*

REMARK 4.15.1.

(i) By Summary 4.15, (ii), one may conclude that

the condition for an MLF to be *absolutely abelian* may be considered to be “*group-theoretic*”.

(ii) On the other hand, it follows from example (1) given in [4], §2 [i.e., “ L_1 ” and “ L_3 ” in (1) of [4], §2], that there exist an *absolutely Galois* MLF k_\circ and an MLF k_\bullet that is *not absolutely Galois* such that the absolute Galois group of k_\circ is *isomorphic*, as an abstract profinite group, to the absolute Galois group of k_\bullet . By this fact, one may conclude that

the condition for an MLF to be *absolutely Galois* should be considered to be “*not group-theoretic*”.

5. RECONSTRUCTION ALGORITHMS RELATED TO MLF’S OF DEGREE ONE

In the present §5, we maintain the notational conventions introduced at the beginnings of §1 and §2. In particular, we have been given a group *of MLF-type*

$$G.$$

In the present §5, suppose that

$$d(G) = 1.$$

In the present §5, we establish some *group-theoretic reconstruction algorithms* [cf. [8], Remark 1.9.8] related to MLF’s *of degree one*, i.e., such that the integer “ $d_{(-)}$ ” [cf. the notational conventions introduced at the beginning of §1] is *equal to one*. As a consequence of the group-theoretic reconstruction algorithms, we also prove [cf. Theorem 5.9, (ii), below] that every *absolutely strictly radical* [cf. Definition 5.6, (iii), below] MLF is *absolutely characteristic* [cf. Definition 5.7 below].

LEMMA 5.1. — *The homomorphism*

$$\mathrm{Nm}_{\mathrm{abs}}(G): k^\times(G) \longrightarrow \mathbb{Q}_p(G)^\times$$

[cf. Definition 4.7, (iii)] *is an isomorphism of topological modules.*

PROOF. — Since [we have assumed that] $d(G) = 1$, this assertion follows from Proposition 4.9, (i). \square

DEFINITION 5.2. — Consider the isomorphism $k_\times(G) \xrightarrow{\sim} \mathbb{Q}_p(G)_\times$ of topological monoids [cf. Remark 2.3.1] determined by the isomorphism $k^\times(G) \xrightarrow{\sim} \mathbb{Q}_p(G)^\times$ of Lemma 5.1 [cf. the discussion entitled “Fields” in §0]. Then, by means of the topological field structure of $\mathbb{Q}_p(G)$, i.e., on $\mathbb{Q}_p(G)_\times$, together with this isomorphism, one may define a *structure of topological field* on $k_\times(G)$. We shall write

$$k(G)$$

for the *resulting topological field*. Thus, we have a tautological isomorphism of topological fields

$$k(G) \xrightarrow{\sim} \mathbb{Q}_p(G)$$

and natural identifications

$$k(G)_\times = k_\times(G), \quad k(G)^\times = k^\times(G).$$

REMARK 5.2.1. — One verifies immediately that the topological field $k(G)$ is *isomorphic*, as an abstract topological field, to the topological field $\mathbb{Q}_p(G)$ [cf. also Lemma 4.6, (i)].

DEFINITION 5.3. — Let $H \curvearrowright M$ be an $\text{MLF}^\triangleright$ -pair [cf. [3], Definition 5.3] such that $d(H) = 1$ [cf. [3], Remark 5.4,1]. Thus, the Kummer poly-isomorphism $\kappa(H \curvearrowright M): (H \curvearrowright M) \xrightarrow{\sim} (H \curvearrowright M^\triangleright(H) = \overline{\mathcal{O}}^\triangleright(H))$ [cf. [3], Definition 5.8; [3], Definition 7.4] associated to $H \curvearrowright M$ consists of a single isomorphism [cf. [3], Definition 5.5] [i.e., as opposite to just a poly-isomorphism]. Then, by means of the topological field structure of $k(H)$, i.e., on $k_\times(H)$, of Definition 5.2, together with the isomorphism $((M^{\text{gp}})^H)^\otimes \xrightarrow{\sim} (\overline{k}^\times(H)^H)^\otimes = k_\times(H)$ [cf. [3], Proposition 5.7, (i)] induced by the Kummer poly-isomorphism $\kappa(H \curvearrowright M)$ [consisting of a single isomorphism], one may define a *structure of topological field* on $((M^{\text{gp}})^H)^\otimes$. We shall write

$$k(H \curvearrowright M)$$

for the *resulting topological field*. Thus, we have tautological isomorphisms of topological fields

$$k(H \curvearrowright M) \xrightarrow{\sim} k(H) \xrightarrow{\sim} \mathbb{Q}_p(H)$$

and natural identifications

$$k(H \curvearrowright M)_\times = ((M^{\text{gp}})^H)^\otimes, \quad k(H \curvearrowright M)^\times = (M^{\text{gp}})^H.$$

REMARK 5.3.1. — One verifies immediately that, in the situation of Definition 5.3, the topological field $k(H \curvearrowright M)$ is *isomorphic*, as an abstract topological field, to the topological field $\mathbb{Q}_p(H)$ [cf. also Lemma 4.6, (i)].

REMARK 5.3.2. — Let us recall the “*étale-like*” $\text{MLF}^\triangleright$ -pair $G \curvearrowright \overline{\mathcal{O}}^\triangleright(G)$ [cf. [3], Definition 5.8]. Then one verifies immediately from the various definitions involved that we have a natural identification

$$k(G) = k(G \curvearrowright \overline{\mathcal{O}}^\triangleright(G)).$$

Recall the MLF

$$k$$

introduced at the beginning of §1.

THEOREM 5.4. — *Suppose that $d_k = 1$, which thus implies that $d(G_k) = 1$ [cf. [3], Proposition 3.6]. Then the following hold:*

(i) *The homomorphism*

$$\text{rec}_k: k^\times \hookrightarrow G_k^{\text{ab}}$$

of [3], Lemma 1.7, determines an isomorphism of topological fields

$$k \xrightarrow{\sim} k(G_k).$$

(ii) *By applying the reconstruction algorithm of Definition 5.3 to the model $\text{MLF}^\triangleright$ -pair $G_k \curvearrowright \mathcal{O}_k^\triangleright$ [cf. [3], Definition 5.2], we obtain a topological field $k(G_k \curvearrowright \mathcal{O}_k^\triangleright)$ whose underlying set may be identified with the underlying set of k . Then the topological field structure of k on the underlying set of k **coincides**, relative to this identification, with the topological field structure of $k(G_k \curvearrowright \mathcal{O}_k^\triangleright)$ on the underlying set of k .*

PROOF. — These assertions follow immediately from Proposition 4.9, (i). \square

COROLLARY 5.5. — *The image of the natural homomorphism $\text{Aut}(G) \rightarrow \text{Aut}(G^{\text{ab}})$ is trivial.*

PROOF. — Let α be an automorphism of G . Now let us observe that since the subset $k^\times(G) \subseteq G^{\text{ab}}$ of G^{ab} is *dense* [cf. [3], Lemma 1.7, (1); [3], Proposition 3.11, (i)], to verify Corollary 5.5, it suffices to verify that the automorphism $k^\times(\alpha)$ of $k^\times(G)$ induced by α is the *identity automorphism*. On the other hand, since $k^\times(\alpha)$ extends to an *automorphism of the topological field $k(G)$* , and the topological field $k(G)$ is *isomorphic*, as an abstract topological field, to the topological field \mathbb{Q}_{p_k} [cf. Remark 5.2.1; [3], Proposition 3.6], we conclude that $k^\times(\alpha)$ is the *identity automorphism*, as desired. This completes the proof of Corollary 5.5. \square

DEFINITION 5.6.

(i) We shall refer to a collection of data

$$(n; m; r_1, \dots, r_m; a_1, \dots, a_m)$$

consisting of

- positive integers n, m, r_1, \dots, r_m such that $n \in \bigcap_{i=1}^m r_i \mathbb{Z}$ and
- elements $a_1, \dots, a_m \in k^\times$ of k^\times

as a *strictly radical data* for k .

(ii) Let $K \subseteq \bar{k}$ be a finite extension of k . Then we shall say that the finite extension K/k is *strictly radical* if there exists a strictly radical data $(n; m; r_1, \dots, r_m; a_1, \dots, a_m)$ for k such that

$$K = k(\zeta_n, a_1^{1/r_1}, \dots, a_m^{1/r_m}) \subseteq \bar{k}.$$

(iii) We shall say that the MLF k is *absolutely strictly radical* if the finite extension $k/k^{(d=1)}$ is strictly radical.

REMARK 5.6.1. — One verifies easily that a *strictly radical* extension is *Galois*. In particular, an *absolutely strictly radical* MLF is *absolutely Galois* [cf. Definition 4.2, (i)].

DEFINITION 5.7. — We shall say that the MLF k is *absolutely characteristic* if the open subgroup $G_k \subseteq \text{Gal}(\bar{k}/k^{(d=1)})$ of $\text{Gal}(\bar{k}/k^{(d=1)})$ is characteristic [cf. Remark 5.7.1 below].

REMARK 5.7.1. — One verifies immediately that the issue of whether or not the MLF k satisfies the condition that the open subgroup $G_k \subseteq \text{Gal}(\bar{k}/k^{(d=1)})$ of $\text{Gal}(\bar{k}/k^{(d=1)})$ is characteristic [cf. Definition 5.7] does *not depend* on the choice of \bar{k} , i.e., *depends only* on k .

REMARK 5.7.2.

(i) Let us recall that since G is *topologically finitely generated* [cf., e.g., [3], Lemma 1.4, (i)], one verifies easily that the topology of the profinite group G admits a basis of *characteristic* open subgroups.

(ii) It follows from (i) that there exists a finite extension $K \subseteq \bar{k}$ of k such that the MLF K is *absolutely characteristic*.

DEFINITION 5.8.

(i) Let $H \curvearrowright M$ be an $\text{MLF}^{\triangleright}$ -pair such that $d(H) = 1$ and $(n; m; r_1, \dots, r_m; a_1, \dots, a_m)$ a strictly radical data for the MLF $k(H \curvearrowright M)$ of Definition 5.3 [cf. also Remark 5.3.1]. Then we shall write

$$(H \curvearrowright M)(n; m; r_1, \dots, r_m; a_1, \dots, a_m) \subseteq H$$

for the uniquely determined maximal subgroup of H which acts trivially on the subset of M^{gp}

$$\{ a \in M^{\text{gp}} \mid a^n = 1 \text{ or } a^{r_i} = a_i \text{ for some } i \in \{1, \dots, m\} \}.$$

(ii) Let $(n; m; r_1, \dots, r_m; a_1, \dots, a_m)$ be a strictly radical data for the MLF $k(G)$ of Definition 5.2 [cf. also Remark 5.2.1]. Then we shall write

$$G(n; m; r_1, \dots, r_m; a_1, \dots, a_m) \stackrel{\text{def}}{=} (G \curvearrowright \overline{\mathcal{O}}^{\triangleright}(G))(n; m; r_1, \dots, r_m; a_1, \dots, a_m) \subseteq G$$

[cf. Remark 5.3.2].

THEOREM 5.9. — *The following hold:*

(i) *Suppose that $d_k = 1$, which thus implies that $d(G_k) = 1$ [cf. [3], Proposition 3.6]. Let $(n; m; r_1, \dots, r_m; a_1, \dots, a_m)$ be a **strictly radical data** for $k \xrightarrow{\sim} k(G_k)$ [cf. Theorem 5.4, (i)]. Then it holds that*

$$G_k(n; m; r_1, \dots, r_m; a_1, \dots, a_m) = \text{Gal}(\bar{k}/k(\zeta_n, a_1^{1/r_1}, \dots, a_m^{1/r_m}))$$

— *i.e., as subgroups of G_k .*

(ii) *Every **absolutely strictly radical MLF** is **absolutely characteristic** [cf. also Remark 5.9.1 below].*

PROOF. — Assertion (i) follows immediately from the various definitions involved. Assertion (ii) follows immediately from assertion (i) and Corollary 5.5. This completes the proof of Theorem 5.9. \square

REMARK 5.9.1. — Note that there exists an *absolutely characteristic* MLF that is *not absolutely strictly radical*. Indeed, let us observe that one verifies immediately from *Kummer theory* that if k is *absolutely strictly radical*, then the Galois group $\text{Gal}(k/k^{(d=1)})$ [cf. Remark 5.6.1] has a structure of *extension of an abelian group by an abelian group*. Thus, it follows from Remark 5.7.2, (i), that if every *absolutely characteristic* MLF is *absolutely strictly radical*, then we conclude that the absolute Galois group $\text{Gal}(\bar{k}/k^{(d=1)})$

has a structure of *extension of an abelian group by an abelian group* — in contradiction to some well-known group-theoretic properties [cf., e.g., [12], Theorem 7.5.12] of the group $\text{Gal}(\bar{k}/k^{(d=1)})$.

REMARK 5.9.2.

(i) It follows from example (1) given in [4], §2 [i.e., “ L_1 ” and “ L_3 ” in (1) of [4], §2], that there exist an *absolutely strictly radical* MLF k_\circ and an MLF k_\bullet that is *not absolutely strictly radical* [cf. Remark 5.6.1] such that the absolute Galois group of k_\circ is *isomorphic*, as an abstract profinite group, to the absolute Galois group of k_\bullet . By this fact, one may conclude that

the condition for an MLF to be *absolutely strictly radical* should be considered to be “*not group-theoretic*”.

(ii) It follows from example (1) given in [4], §2 [i.e., “ L_1 ” and “ L_3 ” in (1) of [4], §2], that there exist an *absolutely characteristic* MLF k_\circ [cf. Theorem 5.9, (ii)] and an MLF k_\bullet that is *not absolutely characteristic* such that the absolute Galois group of k_\circ is *isomorphic*, as an abstract profinite group, to the absolute Galois group of k_\bullet . By this fact, one may conclude that

the condition for an MLF to be *absolutely characteristic* should be considered to be “*not group-theoretic*”.

REMARK 5.9.3.

(i) Let us first observe the following “*tautological*” assertion in the anabelian geometry of *absolutely characteristic* MLF’s:

For each $\square \in \{\circ, \bullet\}$, let k_\square be an *absolutely characteristic* MLF and \bar{k}_\square an algebraic closure of k_\square ; write $G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$. Then the following two conditions are equivalent:

- (1) The field k_\circ is *isomorphic* to the field k_\bullet .
- (2) There exists an *isomorphism* $G_\circ \xrightarrow{\sim} G_\bullet$ compatible with the respective natural outer actions of $\text{Gal}(k_\circ/k_\circ^{(d=1)})$, $\text{Gal}(k_\bullet/k_\bullet^{(d=1)})$ [i.e., by conjugation] relative to some isomorphism $\text{Gal}(k_\circ/k_\circ^{(d=1)}) \xrightarrow{\sim} \text{Gal}(k_\bullet/k_\bullet^{(d=1)})$.

[This assertion follows immediately from the definition of the notion of *absolutely characteristic* MLF — cf. also [3], Proposition 3.6.]

(ii) It follows from Theorem 5.9, (ii), that one may apply the “*tautological*” assertion of (i) to *absolutely strictly radical* MLF’s.

(iii) Finally, let us observe that it follows from example (1) given in [4], §2 [i.e., “ L_1 ” and “ L_2 ” in (1) of [4], §2], that there exist *absolutely strictly radical* MLF’s k_\circ , k_\bullet such that the field k_\circ is *not isomorphic* to the field k_\bullet , but the absolute Galois group of k_\circ is *isomorphic*, as an abstract profinite group, to the absolute Galois group of k_\bullet . Thus, we conclude from Theorem 5.9, (ii), that, in the “*tautological*” assertion of (i), one *cannot* replace condition (2) by the following condition:

(2') There exists an *isomorphism* $G_o \xrightarrow{\sim} G_\bullet$.

REMARK 5.9.4. — Let us recall the following three well-known facts in *anabelian geometry*:

(1) One verifies easily that an immediate consequence of the *Neukirch-Uchida* theorem [cf. the main theorem of [16]] is that every *normal* open subgroup of the absolute Galois group of the field of rational numbers is *characteristic*.

(2) It follows immediately from [7], Corollary 3.7, that it holds that the natural injection $\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$ [cf., e.g., [3], Proposition 2.1] is *bijective* if and only if each member of the filtration on G_k given by the higher ramification groups in the upper numbering is *characteristic*.

(3) Suppose that $d_k = 1$. Then it follows immediately from the equivalence of (2) [cf. also the argument in Remark 6.4.1, (ii), below] that the natural injection $\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$ [cf., e.g., [3], Proposition 2.1] is *bijective* if and only if every *normal* open subgroup of G_k is *characteristic*. [Note that this equivalence also follows from [5], Theorem A.]

By these facts, one may find the *importance* of discussing the issue of whether or not a given closed subgroup of the absolute Galois group of a field is *characteristic* in the study of *anabelian geometry*. This observation is one of motivations of studying Theorem 5.9, (ii).

Some of the group-theoretic reconstruction algorithms discussed in the present §5 may be summarized as follows.

SUMMARY 5.10. — *There exist group-theoretic reconstruction algorithms* [cf. [8], Remark 1.9.8] *for constructing, from a group G of MLF-type such that $d(G) = 1$,*

- *a structure of topological field on $k_\times(G)$ [cf. Definition 5.2] and*
- *subgroups of G [cf. Definition 5.8, (ii)]*

which “correspond” to

- *the topological field structure of k on k_\times [cf. Theorem 5.4, (i)] and*
- *the open subgroups of G_k corresponding to the absolutely strictly radical MLF's contained in \bar{k} [cf. Theorem 5.9, (i)],*

respectively.

REMARK 5.10.1. — Let us recall that, as asserted in Summary 5.10, we have established [cf. Definition 5.2] a group-theoretic reconstruction algorithm for constructing, from a group G of MLF-type such that $d(G) = 1$, a *structure of topological field* on $k_\times(G)$ which “*corresponds*” to the *topological field structure* of k , i.e., on k_\times .

Here, let us also recall that, as discussed in [2], Remark 1.4.1, (ii), it is *impossible* to establish a group-theoretic reconstruction algorithm for constructing, from an *arbitrary* group G of MLF-type, such a *structure of topological field* on $k_\times(G)$.

6. RECONSTRUCTION ALGORITHMS RELATED TO GALOIS-SPECIFIABLE MLF'S

In the present §6, we maintain the notational conventions introduced at the beginnings of §1 and §2. In the present §6, we consider *Galois-specifiable* [cf. Definition 6.1 below] MLF's. Moreover, we also establish some *group-theoretic reconstruction algorithms* [cf. [8], Remark 1.9.8] related to *Galois-specifiable* MLF's. For instance, we establish a *group-theoretic reconstruction algorithm* for constructing, from a group of *MLF-type* that satisfies a certain condition, a collection of subgroups of the outer automorphism group of the group of MLF-type which “corresponds” to the $\text{Out}(G_k)$ -orbit, i.e., by conjugation, of the subgroup of $\text{Out}(G_k)$ determined by the field automorphisms of k [cf. Definition 6.9, (ii); Theorem 6.13, (ii), below].

DEFINITION 6.1. — We shall say that the MLF k is *Galois-specifiable* if the MLF k is absolutely Galois [cf. Definition 4.2, (i)], and, moreover, the following condition is satisfied: If L is an MLF such that there exist an algebraic closure \bar{L} of L and an isomorphism $G_k \xrightarrow{\sim} \text{Gal}(\bar{L}/L)$ of groups, then the field k is isomorphic, as an abstract field, to the field L [cf. Remark 6.1.1 below].

REMARK 6.1.1. — One verifies immediately that the issue of whether or not the MLF k satisfies the condition in Definition 6.1 does *not depend* on the choice of \bar{k} , i.e., *depends only* on k .

REMARK 6.1.2. — Suppose that $(p_k, a_k) \neq (2, 1)$. Then it follows immediately — in light of Proposition 2.5, (i); Proposition 4.9, (i); [3], Proposition 3.6 — from Theorem 4.11, together with [3], Lemma 1.7, (1), (2), that it holds that the MLF k is *Galois-specifiable* if and only if the following condition is satisfied: If $K \subseteq \bar{k}$ is a finite extension of $k^{(d=1)}$ such that $d_k = d_K$, and, moreover, $k^{(\text{ab})} = K^{(\text{ab})}$ [cf. Definition 4.2, (iii)], i.e., as subfields of \bar{k} , then $k = K$, i.e., as subfields of \bar{k} .

PROPOSITION 6.2. — *Suppose that the MLF k is Galois-specifiable. Let K be an intermediate field of the finite extension $k/k^{(\text{ab})}$ tamely ramified over $k^{(\text{ab})}$. Then the MLF K is Galois-specifiable.*

PROOF. — Let L be an MLF, \bar{L} an algebraic closure of L , and $\alpha: G_K \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/K) \xrightarrow{\sim} G_L \stackrel{\text{def}}{=} \text{Gal}(\bar{L}/L)$ an isomorphism of groups. Thus, it follows from [3], Proposition 3.6, that

$$(1) \quad (p_K, e_K) = (p_L, e_L).$$

Write $L_k \subseteq \bar{L}$ for the finite extension of L which corresponds to the open subgroup $\alpha(G_k) \subseteq G_L$ of G_L . [So we have a natural identification $G_{L_k} \stackrel{\text{def}}{=} \text{Gal}(\bar{L}/L_k) = \alpha(G_k)$.] Then since [we have assumed that] k is *Galois-specifiable*, we conclude that

$$(2) \quad \text{the field } k \text{ is isomorphic to the field } L_k.$$

Next, observe that, by applying Corollary 4.14, (i), to the isomorphism α , we obtain that

(3) the field $K^{(\text{ab})}$, i.e., $k^{(\text{ab})}$ [cf. Lemma 4.3, (i)], is *isomorphic* to the field $L^{(\text{ab})}$.

Now I *claim* the following assertion:

(*): Suppose that K is the [uniquely determined] *maximal* MLF contained in k *tamely ramified* over $k^{(\text{ab})}$. Then the field K is *isomorphic* to the field L .

To this end, suppose that K is the [uniquely determined] *maximal* MLF contained in k *tamely ramified* over $k^{(\text{ab})}$. In particular, it follows from Lemma 4.3, (ii), that the intermediate field K of $k/k^{(\text{ab})}$ is the *uniquely determined* intermediate field of $k/k^{(\text{ab})}$ such that e_k/e_K is a *power of* p_K , and, moreover, $e_K/e_{k^{(\text{ab})}}$ is *prime to* p_K . Thus, it follows immediately from (1), (2), and (3) that the intermediate field L of $L_k/L^{(\text{ab})}$ is the *uniquely determined* intermediate field of $L_k/L^{(\text{ab})}$ such that e_{L_k}/e_L is a *power of* p_L , and, moreover, $e_L/e_{L^{(\text{ab})}}$ is *prime to* p_L . In particular, by (2) and (3), we conclude that the field K is *isomorphic* to the field L , as desired. This completes the proof of the above *claim* (*).

Next, observe that it follows immediately from the above *claim* (*) that, to complete the verification of Proposition 6.2, we may assume without loss of generality, by replacing k by the [uniquely determined] *maximal* MLF contained in k *tamely ramified* over $k^{(\text{ab})}$, that k is *tamely ramified* over $k^{(\text{ab})}$. Then one verifies immediately from Lemma 4.3, (ii), and [3], Lemma 1.5, (ii), that the intermediate field K (respectively, L) of $k/k^{(\text{ab})}$ (respectively, $L_k/L^{(\text{ab})}$) is the *uniquely determined* intermediate field M of $k/k^{(\text{ab})}$ (respectively, $L_k/L^{(\text{ab})}$) such that $e_M = e_K$ (respectively, $e_M = e_L$). Thus, by (1), (2), and (3), we conclude that the field K is *isomorphic* to the field L , as desired. This completes the proof of Proposition 6.2. \square

LEMMA 6.3. — *Suppose that k is Galois over $k^{(\text{ab})}$. Let $K \subseteq \bar{k}$ be a finite unramified [necessarily Galois] extension of $k^{(\text{ab})}$. Note that it follows immediately from Lemma 4.3, (ii), that we have a natural isomorphism $\text{Gal}(k \cdot K/k^{(\text{ab})}) \xrightarrow{\sim} \text{Gal}(k/k^{(\text{ab})}) \times \text{Gal}(K/k^{(\text{ab})})$. Let $\phi: \text{Gal}(K/k^{(\text{ab})}) \rightarrow \text{Gal}(k/k^{(\text{ab})})$ be a homomorphism of groups. Write L for the intermediate field of the finite Galois extension $k \cdot K/k^{(\text{ab})}$ which corresponds, relative to the above natural isomorphism $\text{Gal}(k \cdot K/k^{(\text{ab})}) \xrightarrow{\sim} \text{Gal}(k/k^{(\text{ab})}) \times \text{Gal}(K/k^{(\text{ab})})$, to the graph ($\subseteq \text{Gal}(k/k^{(\text{ab})}) \times \text{Gal}(K/k^{(\text{ab})})$) of the homomorphism ϕ . Then the equalities $d_L = d_k$, $L^{(\text{ab})} = k^{(\text{ab})}$ hold.*

PROOF. — The equality $d_L = d_k$ follows from the fact that the graph ($\subseteq \text{Gal}(k/k^{(\text{ab})}) \times \text{Gal}(K/k^{(\text{ab})})$) of the homomorphism ϕ is *isomorphic*, as an abstract group, to the group $\text{Gal}(K/k^{(\text{ab})})$.

To verify the equality $L^{(\text{ab})} = k^{(\text{ab})}$, assume that $L^{(\text{ab})} \neq k^{(\text{ab})}$, i.e., that the extension $L^{(\text{ab})}/k^{(\text{ab})}$ is *not of degree one*. Then since the intermediate field L corresponds to the graph of ϕ , one verifies immediately from the natural isomorphism $\text{Gal}(k \cdot K/k^{(\text{ab})}) \xrightarrow{\sim} \text{Gal}(k/k^{(\text{ab})}) \times \text{Gal}(K/k^{(\text{ab})})$ that the extension $L^{(\text{ab})} \cdot K/K$ is *not of degree one*, which thus implies that the extension $k \cap (L^{(\text{ab})} \cdot K)/k^{(\text{ab})}$ is *not of degree one*. On the other hand, one verifies easily [cf. the proof of Lemma 4.3, (ii)] that the MLF $L^{(\text{ab})} \cdot K$, hence also the MLF $k \cap (L^{(\text{ab})} \cdot K)$, is *absolutely abelian* [cf. Definition 4.2, (ii)]. Thus, we obtain a *contradiction* [cf. Lemma 4.3, (i)]. This completes the proof of the equality $L^{(\text{ab})} = k^{(\text{ab})}$, hence also of Lemma 6.3. \square

THEOREM 6.4. — Consider the following four conditions:

- (1) The MLF k is **absolutely abelian** [cf. Definition 4.2, (ii)].
- (2) The MLF k is **Galois-specifiable**.
- (3) The MLF k is **absolutely characteristic** [cf. Definition 5.7].
- (4) The MLF k is **absolutely Galois** [cf. Definition 4.2, (i)].

Then the following hold:

- (i) The implications

$$(1) \implies (2) \implies (3) \implies (4)$$

hold [cf. also Remark 6.4.1 below].

- (ii) Suppose that $(p_k, a_k) \neq (2, 1)$. Then the equivalence

$$(1) \iff (2)$$

holds.

PROOF. — First, we verify assertion (i). The first implication in assertion (i) follows immediately from Corollary 4.14, (ii). The second and third implications in assertion (i) follow immediately from the various definitions involved. This completes the proof of assertion (i).

Next, we verify assertion (ii). By assertion (i), to verify assertion (ii), it suffices to verify the implication $(2) \implies (1)$. Suppose that k is *Galois-specifiable*. To verify that k is *absolutely abelian*, let $A \subseteq \text{Gal}(k/k^{(\text{ab})})$ be a *cyclic* subgroup of $\text{Gal}(k/k^{(\text{ab})})$. Then it follows from [3], Lemma 1.5, (i), that there exist a finite *unramified* [necessarily Galois] extension $K \subseteq \bar{k}$ of $k^{(\text{ab})}$ and an *isomorphism* $\phi: \text{Gal}(K/k^{(\text{ab})}) \xrightarrow{\sim} A (\subseteq \text{Gal}(k/k^{(\text{ab})}))$ of groups. Thus, it follows from Lemma 6.3 that the graph of ϕ determines an MLF $L \subseteq k \cdot K$ such that $d_L = d_k$, and, moreover, $L^{(\text{ab})} = k^{(\text{ab})}$. On the other hand, since [we have assumed that] k is *Galois-specifiable*, it follows from Remark 6.1.2 that $k = L$, i.e., that the homomorphism ϕ , hence also the subgroup A , is *trivial*. In particular, we conclude that every *cyclic* subgroup of $\text{Gal}(k/k^{(\text{ab})})$, hence also the group $\text{Gal}(k/k^{(\text{ab})})$ itself, is *trivial*, as desired. This completes the proof of assertion (ii), hence also of Theorem 6.4. \square

REMARK 6.4.1. — Suppose that we are in the situation of Theorem 6.4.

(i) In general, the implication $(3) \implies (2)$ does *not hold*. Indeed, for an *odd* prime number p_k , the MLF $\mathbb{Q}_{p_k}(\zeta_{p_k}, p_k^{1/p_k})$ is *absolutely characteristic* [cf. Theorem 5.9, (ii)] but *not Galois-specifiable* [cf. example (1) given in [4], §2].

(ii) In general, the implication $(4) \implies (3)$ does *not hold*. Indeed, let us first observe that it is immediate that the implication $(4) \implies (3)$ is equivalent to the following assertion:

Every *normal open* subgroup of $G_{k^{(d=1)}} \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k^{(d=1)})$ is *characteristic*.

Thus, since every normal closed subgroup of $G_{k^{(d=1)}}$ may be obtained as the intersection of the normal open subgroups of $G_{k^{(d=1)}}$ that contain the normal closed subgroup, if the implication $(4) \implies (3)$ holds, then we conclude that every *normal closed* subgroup of $G_{k^{(d=1)}}$ is *characteristic*. In particular, if the implication $(4) \implies (3)$ holds, then it

follows from [7], Corollary 3.7, that every outer automorphism of $G_{k^{(d=1)}}$ arises from an automorphism of the field $k^{(d=1)}$, i.e., that $\text{Out}(G_{k^{(d=1)}}) = \{1\}$. But this contradicts the conclusion of the discussion given at the final portion of [12], Chapter VII, §5, if p_k is odd.

DEFINITION 6.5. — Recall that the natural homomorphism $\text{Aut}(k) \rightarrow \text{Out}(G_k)$ is injective [cf., e.g., [3], Proposition 2.1]. By means of this injection, let us regard $\text{Aut}(k)$ as a [necessarily finite] subgroup of $\text{Out}(G_k)$:

$$\text{Aut}(k) \subseteq \text{Out}(G_k).$$

Then we shall write

$$\text{OrbAut}(k)$$

for the set of $\text{Out}(G_k)$ -conjugates of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$, i.e., the $\text{Out}(G_k)$ -orbit of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$.

Recall the group of *MLF-type*

$$G$$

introduced at the beginning of §2.

DEFINITION 6.6. — Let $\Gamma \subseteq \text{Out}(G)$ be a finite subgroup of the outer automorphism group $\text{Out}(G)$ of G .

(i) We shall write

$$G \rtimes^{\text{out}} \Gamma \subseteq \text{Aut}(G)$$

for the inverse image of $\Gamma \subseteq \text{Out}(G)$ by the natural surjection $\text{Aut}(G) \rightarrow \text{Out}(G)$. Thus, since G may be identified with $\text{Inn}(G) \subseteq \text{Aut}(G)$ by the natural isomorphism $G \xrightarrow{\sim} \text{Inn}(G)$ [cf. [3], Lemma 1.8, (i)], the group $G \rtimes^{\text{out}} \Gamma$ has a natural structure of extension of Γ by G :

$$1 \longrightarrow G \longrightarrow G \rtimes^{\text{out}} \Gamma \longrightarrow \Gamma \longrightarrow 1.$$

By means of the second arrow of this exact sequence, let us always regard G as a subgroup of $G \rtimes^{\text{out}} \Gamma$:

$$G \subseteq G \rtimes^{\text{out}} \Gamma.$$

(ii) We shall say that the finite subgroup Γ is *quasi-geometric* if the group $G \rtimes^{\text{out}} \Gamma$ is of MLF-type.

(iii) We shall say that the finite subgroup Γ is *strictly quasi-geometric* if Γ is quasi-geometric, and, moreover, the equality $d(G \rtimes^{\text{out}} \Gamma) = 1$ holds.

LEMMA 6.7. — Let $\Gamma \subseteq \text{Out}(G)$ be a **quasi-geometric** subgroup of $\text{Out}(G)$. Then the following hold:

(i) Every subgroup of Γ is a **quasi-geometric** subgroup of $\text{Out}(G)$.

(ii) The natural inclusion $k^\times(G \rtimes^{\text{out}} \Gamma) \hookrightarrow k^\times(G)$ — i.e., determined by the transfer map with respect to $G \subseteq G \rtimes^{\text{out}} \Gamma$ [cf. [3], Lemma 1.7, (3)] — and the natural action of Γ on $k^\times(G)$ determine an **isomorphism** $k^\times(G \rtimes^{\text{out}} \Gamma) \xrightarrow{\sim} k^\times(G)^\Gamma$.

(iii) It holds that $d(G) \in \sharp\Gamma\mathbb{Z}$.

(iv) It holds that Γ is **strictly quasi-geometric** if and only if $\sharp\Gamma = d(G)$.

PROOF. — Assertion (i) follows from the discussion following [3], Proposition 3.3. Assertion (ii) follows immediately from [3], Proposition 3.11, (i). Assertions (iii), (iv) follow immediately from [3], Proposition 3.6. This completes the proof of Lemma 6.7. \square

PROPOSITION 6.8. — *The subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is **quasi-geometric**. If, moreover, the MLF k is **absolutely Galois**, then the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is **strictly quasi-geometric**.*

PROOF. — This assertion follows immediately from the various definitions involved. \square

DEFINITION 6.9.

(i) We shall say that G is of *GSMLF-type* if the following two conditions are satisfied:

(1) There exists a strictly quasi-geometric subgroup of $\text{Out}(G)$.

(2) For each strictly quasi-geometric subgroup $\Gamma \subseteq \text{Out}(G)$ of $\text{Out}(G)$ and each open subgroup $H \subseteq G \rtimes^{\text{out}} \Gamma$ of $G \rtimes^{\text{out}} \Gamma$, if H is isomorphic, as an abstract group, to G , then $H = G$, i.e., as subgroups of $G \rtimes^{\text{out}} \Gamma$.

[Here, “GSMLF” is to be understood as an abbreviation for “Galois-specifiable mixed-characteristic local field” — cf. Theorem 6.11 below.]

(ii) Suppose that G is of GSMLF-type. Then we shall write

$$\text{Orb}_{\text{sqg}}(G)$$

for the set of strictly quasi-geometric subgroups of $\text{Out}(G)$.

LEMMA 6.10. — *Let $\overline{\mathbb{Q}}_{p_k}$ be an algebraic closure of $k^{(d=1)}$. Let us fix an isomorphism $\iota: \overline{k} \xrightarrow{\sim} \overline{\mathbb{Q}}_{p_k}$ of fields. Write $k_\iota \stackrel{\text{def}}{=} \iota(k) \subseteq \overline{\mathbb{Q}}_{p_k}$ for the finite extension of $k^{(d=1)}$ obtained by forming the image of $k \subseteq \overline{k}$ by ι . Let $\Gamma \subseteq \text{Out}(G_k)$ be a **strictly quasi-geometric** subgroup of the outer automorphism group $\text{Out}(G_k)$ of the group G_k of MLF-type. Then the following hold:*

(i) There exists an **isomorphism** $\alpha: G_k \rtimes^{\text{out}} \Gamma \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$ of groups.

(ii) Suppose either that the MLF k is **Galois-specifiable**, or that the group G_k is of **GSMLF-type**. Then the isomorphism α of (i) restricts to an isomorphism between the subgroup $G_k \subseteq G_k \rtimes^{\text{out}} \Gamma$ with the subgroup $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota) \subseteq \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$.

PROOF. — Assertion (i) follows immediately from the definition of the notion of *strictly quasi-geometric* subgroup [cf. also [3], Proposition 3.6].

Next, we verify assertion (ii) in the case where the MLF k is *Galois-specifiable*. Suppose that k is *Galois-specifiable*. Write $K \subseteq \overline{\mathbb{Q}}_{p_k}$ for the finite extension of $k^{(d=1)}$ which corresponds to the open subgroup $\alpha(G_k) \subseteq \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$ of $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$. [So we have a natural identification $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/K) = \alpha(G_k)$.] Thus, since k is *Galois-specifiable*, we conclude that k is *isomorphic*, as an abstract field, to K . In particular, since k is *absolutely Galois*, we conclude that $k_\iota = K$, i.e., as subfields of $\overline{\mathbb{Q}}_{p_k}$, as desired. This completes the proof of assertion (ii) in the case where the MLF k is *Galois-specifiable*.

Finally, we verify assertion (ii) in the case where the group G_k is of *GSMLF-type*. Suppose that G_k is of *GSMLF-type*. Let us first observe that it is immediate that the group G_k is *isomorphic*, as an abstract group, to the group $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota) \subseteq \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$, hence also to the group $\alpha^{-1}(\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota)) \subseteq G_k \overset{\text{out}}{\rtimes} \Gamma$. Thus, since G_k is of *GSMLF-type*, we conclude that $G_k = \alpha^{-1}(\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota))$, i.e., as subgroups of $G_k \overset{\text{out}}{\rtimes} \Gamma$, as desired. This completes the proof of assertion (ii) in the case where the group G_k is of *GSMLF-type*, hence also of Lemma 6.10. \square

THEOREM 6.11. — *It holds that the MLF k is **Galois-specifiable** if and only if the group G_k is of **GSMLF-type**.*

PROOF. — First, we verify the *necessity*. Suppose that the MLF k is *Galois-specifiable*. To verify that the group G_k is of *GSMLF-type*, let $\Gamma \subseteq \text{Out}(G_k)$ be a *strictly quasi-geometric* subgroup of $\text{Out}(G_k)$ [cf. Proposition 6.8] and $H \subseteq G_k \overset{\text{out}}{\rtimes} \Gamma$ an open subgroup of $G_k \overset{\text{out}}{\rtimes} \Gamma$ such that H is *isomorphic*, as an abstract group, to G_k . Now suppose that we are in the situation of Lemma 6.10. Thus, we have an *isomorphism* $\alpha: G_k \overset{\text{out}}{\rtimes} \Gamma \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$ of groups [cf. Lemma 6.10, (i)] that *restricts* to an isomorphism $G_k \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota)$ [cf. Lemma 6.10, (ii)]. Write $K \subseteq \overline{\mathbb{Q}}_{p_k}$ for the finite extension of $k^{(d=1)}$ which corresponds to the open subgroup $\alpha(H) \subseteq \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$ of $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$. [So we have a natural identification $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/K) = \alpha(H)$.] Thus, since H is *isomorphic* to both G_k and $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/K)$, and k is *Galois-specifiable*, we conclude that k is *isomorphic*, as an abstract field, to K . In particular, since k is *absolutely Galois*, it holds that $k_\iota = K$, i.e., as subfields of $\overline{\mathbb{Q}}_{p_k}$, which thus implies that $\alpha(G_k) = \alpha(H)$, i.e., as subgroups of $\text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$, as desired. This completes the proof of the *necessity*.

Next, we verify the *sufficiency*. Suppose that the group G_k is of *GSMLF-type*. To verify that the MLF k is *Galois-specifiable*, let L be an MLF and \overline{L} an algebraic closure of L such that G_k is *isomorphic*, as an abstract group, to $\text{Gal}(\overline{L}/L)$. Now take a *strictly quasi-geometric* subgroup $\Gamma \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$ [cf. condition (1) of Definition 6.9, (i)], and suppose that we are in the situation of Lemma 6.10. Thus, we have an *isomorphism* $\alpha: G_k \overset{\text{out}}{\rtimes} \Gamma \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k^{(d=1)})$ of groups [cf. Lemma 6.10, (i)] that *restricts* to an isomorphism $G_k \xrightarrow{\sim} \text{Gal}(\overline{\mathbb{Q}}_{p_k}/k_\iota)$ [cf. Lemma 6.10, (ii)] — which thus implies that the MLF k_ι , hence also the MLF k , is *absolutely Galois*. Then let us observe that it follows from [3], Proposition 3.6, that \overline{L} is *isomorphic*, as an abstract field, to $\overline{\mathbb{Q}}_{p_k}$. Let us identify \overline{L} with $\overline{\mathbb{Q}}_{p_k}$ by means of some fixed isomorphism of \overline{L} with $\overline{\mathbb{Q}}_{p_k}$. Then since G_k

is *isomorphic* to $\text{Gal}(\bar{L}/L) = \text{Gal}(\bar{\mathbb{Q}}_{p_k}/L)$, and G_k is of *GSMLF-type*, we conclude that $G_k = \alpha^{-1}(\text{Gal}(\bar{\mathbb{Q}}_{p_k}/L))$, i.e., as subgroups of $G_k \overset{\text{out}}{\rtimes} \Gamma$, which thus implies that $k_\iota = L$, i.e., as subfields of $\bar{\mathbb{Q}}_{p_k}$. Thus, the field k is *isomorphic* to the field L , as desired. This completes the proof of the *sufficiency*, hence also of Theorem 6.11. \square

COROLLARY 6.12. — *The following hold:*

(i) *If the group G is of **AAMLF-type** [cf. Definition 4.8], then G is of **GSMLF-type**.*

(ii) *Suppose that $(p(G), a(G)) \neq (2, 1)$ [cf. Definition 2.4, (ii)]. Then if the group G is of **GSMLF-type**, then G is of **AAMLF-type**.*

PROOF. — These assertions follow — in light of Proposition 2.5, (i), and [3], Proposition 3.6 — from Theorem 6.4, (i), (ii), together with Proposition 4.9, (iii), and Theorem 6.11. \square

THEOREM 6.13. — *Suppose that the MLF k is **Galois-specifiable**, which thus implies that the group G_k is of **GSMLF-type** [cf. Theorem 6.11]. Then the following hold:*

(i) *Let $\Gamma_1, \Gamma_2 \subseteq \text{Out}(G_k)$ be **strictly quasi-geometric** subgroups of $\text{Out}(G_k)$. Then Γ_1 is an **Out(G_k)-conjugate** of Γ_2 .*

(ii) *It holds that*

$$\text{OrbAut}(k) = \text{Orb}_{\text{sqg}}(G_k).$$

PROOF. — First, we verify assertion (i). Let $\alpha: G_k \overset{\text{out}}{\rtimes} \Gamma_1 \xrightarrow{\sim} G_k \overset{\text{out}}{\rtimes} \Gamma_2$ be an isomorphism of groups [cf. Lemma 6.10, (i)]. Then since the group G_k is of *GSMLF-type*, it is immediate that the isomorphism α *restricts* to an *automorphism* of G_k . Moreover, one verifies immediately from the various definitions involved that Γ_1 is the conjugate, by the outer automorphism of G_k determined by the resulting automorphism of G_k , of Γ_2 . This completes the proof of assertion (i).

Assertion (ii) follows from assertion (i) and Proposition 6.8. This completes the proof of Theorem 6.13. \square

REMARK 6.13.1. — Note that, in general, a similar assertion to Theorem 6.13, (i), for [the absolute Galois group of] an *absolutely characteristic* MLF does *not hold* [cf. Remark 8.6.1, (ii), below].

Some of the group-theoretic reconstruction algorithms discussed in the present §6 may be summarized as follows.

SUMMARY 6.14.

(i) *There exists a **group-theoretic condition** for a group of **MLF-type** [cf. Definition 6.9, (i)] which “**corresponds**” to the condition for an MLF to be **Galois-specifiable** [cf. Theorem 6.11].*

(ii) *There exists a **group-theoretic reconstruction algorithm** [cf. [8], Remark 1.9.8] for constructing, from a group G of **MLF-type** that satisfies the condition of (i), a collection $\text{Orb}_{\text{sqg}}(G)$ of subgroups of $\text{Out}(G)$ [cf. Definition 6.9, (ii)] which “**corresponds**” to the $\text{Out}(G_k)$ -orbit $\text{OrbAut}(k)$, i.e., by conjugation, of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ [cf. Theorem 6.13, (ii)].*

REMARK 6.14.1. — By Summary 6.14, (i), one may conclude that

the condition for an MLF to be *Galois-specifiable* may be considered to be “*group-theoretic*”.

7. ON OUTER AUTOMORPHISMS ARISING FROM FIELD AUTOMORPHISMS I

In the present §7, we maintain the notational conventions introduced at the beginnings of §1 and §2. In particular, we have a natural open injection

$$G_k \hookrightarrow G_{k^{(d=1)}} \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k^{(d=1)}).$$

In the present §7, we discuss outer automorphisms of the absolute Galois groups of MLF’s that *arise from field automorphisms of the MLF’s*. We prove that if the MLF k is *absolutely characteristic*, and that p_k is *odd*, then the subgroup of the outer automorphism group of G_k determined by the field automorphisms of k is *not normally terminal* [cf. Theorem 7.2, (i), below]. Moreover, we also prove that, under some conditions, the outer automorphism group of G_k has “*many*” *finite abelian* subgroups [cf. Theorem 7.5, Remark 7.5.1, below].

LEMMA 7.1. — *Let $H \subseteq G$ be a **characteristic** open subgroup of the group G of MLF-type. Thus, we have, by considering restrictions, a natural homomorphism*

$$\text{Aut}(G) \longrightarrow \text{Aut}(H).$$

Then the following hold:

(i) *The homomorphism $\text{Aut}(G) \rightarrow \text{Aut}(H)$ is **injective**. In particular, we also have an **injection** $\text{Aut}(G)/\text{Inn}(H) \hookrightarrow \text{Out}(H)$. Let us regard $\text{Aut}(G)$, $\text{Aut}(G)/\text{Inn}(H)$ as subgroups of $\text{Aut}(H)$, $\text{Out}(H)$ by means of these injections, respectively:*

$$\text{Aut}(G) \subseteq \text{Aut}(H), \quad \text{Aut}(G)/\text{Inn}(H) \subseteq \text{Out}(H).$$

(ii) *The natural homomorphisms $G \twoheadrightarrow \text{Inn}(G) \hookrightarrow \text{Aut}(G)$ determine an **isomorphism***

$$G/H \xrightarrow{\sim} \text{Inn}(G)/\text{Inn}(H).$$

Let us identify G/H with $\text{Inn}(G)/\text{Inn}(H)$ by means of this isomorphism:

$$G/H = \text{Inn}(G)/\text{Inn}(H) \quad (\subseteq \text{Aut}(G)/\text{Inn}(H) \subseteq \text{Out}(H)).$$

Thus, we have a natural exact sequence

$$1 \longrightarrow G/H \longrightarrow \text{Aut}(G)/\text{Inn}(H) \longrightarrow \text{Out}(G) \longrightarrow 1.$$

(iii) *It holds that*

$$N_{\text{Out}(H)}(G/H) = \text{Aut}(G)/\text{Inn}(H).$$

(iv) Recall the exact sequence

$$1 \longrightarrow G/H \longrightarrow \text{Aut}(G)/\text{Inn}(H) \longrightarrow \text{Out}(G) \longrightarrow 1$$

of (ii). Then the composite

$$\text{Aut}(G)/\text{Inn}(H) \longrightarrow \text{Aut}(G/H) \longrightarrow \text{Out}(G/H)$$

— where the first arrow is the action by conjugation via the second arrow of the above exact sequence, and the second arrow is the natural surjection — **coincides** with the composite

$$\text{Aut}(G)/\text{Inn}(H) \longrightarrow \text{Out}(G) \longrightarrow \text{Out}(G/H)$$

— where the first arrow is the third arrow of the above exact sequence, and the second arrow is the natural homomorphism.

PROOF. — First, we verify assertion (i). Let us first observe that it follows from [3], Lemma 1.8, (ii), that the action of G on H by conjugation is *faithful*. Since [it is immediate that] the resulting *injection* $G \hookrightarrow \text{Aut}(H)$ is *Aut*(G)-*equivariant*, assertion (i) holds. This completes the proof of assertion (i). Assertion (ii) follows from [3], Lemma 1.8, (i). Assertions (iii), (iv) follow immediately from the various definitions involved. This completes the proof of Lemma 7.1. \square

THEOREM 7.2. — Suppose that the MLF k is **absolutely characteristic** [cf. Definition 5.7]. Then the following hold:

(i) Suppose, moreover, that p_k is **odd**. Then the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$ is **not normally terminal**.

(ii) It holds that the MLF k is **absolutely abelian** [cf. Definition 4.2, (ii)] if and only if

$$N_{\text{Out}(G_k)}(\text{Aut}(k)) = Z_{\text{Out}(G_k)}(\text{Aut}(k)).$$

PROOF. — Since the open subgroup $G_k \subseteq G_{k(d=1)}$ of $G_{k(d=1)}$ is *characteristic*, by applying Lemma 7.1, (ii), (iii) [in the case where we take the “ $H \subseteq G$ ” of Lemma 7.1 to be $G_k \subseteq G_{k(d=1)}$], we obtain an exact sequence

$$1 \longrightarrow \text{Aut}(k) \longrightarrow N_{\text{Out}(G_k)}(\text{Aut}(k)) \longrightarrow \text{Out}(G_{k(d=1)}) \longrightarrow 1.$$

Now we verify assertion (i). Since p_k is *odd*, it follows from the discussion given at the final portion of [12], Chapter VII, §5, that $\text{Out}(G_{k(d=1)})$ is *nontrivial*. Thus, by the above exact sequence, we conclude that the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is *not normally terminal*, as desired. This completes the proof of assertion (i).

Next, we verify assertion (ii). The *sufficiency* is immediate. Let us verify the *necessity*. Suppose that the MLF k is *absolutely abelian*. Let us first observe that it is immediate that, to verify the *necessity*, it suffices to verify that the action of $N_{\text{Out}(G_k)}(\text{Aut}(k))$ on $\text{Aut}(k)$ by conjugation is *trivial*. On the other hand, since k is *absolutely abelian*, it follows immediately from Lemma 7.1, (iv), that this action factors through the natural homomorphism $\text{Out}(G_{k(d=1)}) \rightarrow \text{Out}(G_{k(d=1)}^{\text{ab}}) (= \text{Aut}(G_{k(d=1)}^{\text{ab}}))$. Thus, the desired *triviality* follows from Corollary 5.5. This completes the proof of assertion (ii), hence also of Theorem 7.2. \square

REMARK 7.2.1.

(i) Let us observe that it follows immediately from Corollary 5.5 that if $d_k = 1$, then the image of the natural homomorphism

$$\mathrm{Aut}(k) \longrightarrow \mathrm{Aut}(k_+)$$

coincides with the image of the composite

$$\mathrm{Out}(G_k) \longrightarrow \mathrm{Aut}(k_+(G_k)) \xleftarrow{\sim} \mathrm{Aut}(k_+)$$

— where the second arrow is the isomorphism obtained by conjugation by the vertical isomorphism $k_+ \xrightarrow{\sim} k_+(G_k)$ in the diagram of [3], Proposition 3.11, (iv).

(ii) On the other hand, in general, the image of the natural homomorphism

$$\mathrm{Aut}(k) \longrightarrow \mathrm{Aut}(k_+)$$

does *not coincide* with the image of the composite

$$\mathrm{Out}(G_k) \longrightarrow \mathrm{Aut}(k_+(G_k)) \xleftarrow{\sim} \mathrm{Aut}(k_+).$$

Indeed, suppose that k is *absolutely characteristic*, and that the image of the natural homomorphism $\mathrm{Out}(G_{k^{(d=1)}}) \rightarrow \mathrm{Out}(G_{k^{(d=1)}}/G_k)$ is *nontrivial*, which thus implies [cf. Lemma 7.1, (iv); also the exact sequence in the proof of Theorem 7.2] that the normal subgroup $\mathrm{Aut}(k) \subseteq N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$ is *not a direct summand*. [Note that it follows immediately from the discussion given at the final portion of [12], Chapter VII, §5, together with Remark 5.7.2, (i), that such a “ k ” exists.] Next, observe that since $\mathrm{Aut}(k)$ is *contained* in $N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$, it is immediate that, to verify the desired assertion, it suffices to verify that the image of the natural homomorphism

$$\mathrm{Aut}(k) \longrightarrow \mathrm{Aut}(k_+)$$

does *not coincide* with the image of the composite

$$N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k)) \hookrightarrow \mathrm{Out}(G_k) \rightarrow \mathrm{Aut}(k_+(G_k)) \xleftarrow{\sim} \mathrm{Aut}(k_+).$$

On the other hand, since the natural homomorphism $\mathrm{Aut}(k) \rightarrow \mathrm{Aut}(k_+)$ is *injective*, if these images *coincide*, then one verifies immediately that the normal subgroup $\mathrm{Aut}(k) \subseteq N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$ is a *direct summand* — in *contradiction* to our assumption on k .

REMARK 7.2.2. — The consideration in Remark 7.2.1, (ii), leads us to, for instance, the following assertion:

If $d_k = 2$ [which thus implies that k is *absolutely abelian*], and $p_k - 1 \notin 4\mathbb{Z}$, then the subgroup $\mathrm{Aut}(k) \subseteq N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k)) (= Z_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$ — cf. Theorem 7.2, (ii)) is a *direct summand*.

Indeed, let us first observe that it follows from elementary field theory that the action of the [unique] nontrivial element of $\mathrm{Aut}(k)$ on the \mathbb{Q}_{p_k} -vector space $k_+(G_k)$ ($\xrightarrow{\sim} k_+$ — cf. [3], Proposition 3.11, (iv)) of dimension $d_k = 2$ has two eigenvalues $1, -1$. Write $V_1, V_{-1} \subseteq k_+(G_k)$ for the eigenspaces that corresponds to the eigenvalues $1, -1$, respectively. [So we have $k_+(G_k) = V_1 \oplus V_{-1}$.] Then one verifies easily that the action of $N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$ on $k_+(G_k)$ *preserves* each of the subspaces $V_1, V_{-1} \subseteq k_+(G_k)$. Thus, we have a homomorphism

$$N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k)) \longrightarrow \mathrm{Aut}_{\mathbb{Q}_{p_k}}(V_{-1}) = \mathbb{Q}_{p_k}^\times.$$

Now let us observe that it follows from our assumption that $p_k - 1 \notin 4\mathbb{Z}$, together with [3], Lemma 1.2, (i) [cf. also Proposition 1.1, (v), if $p_k = 2$], that there exists a *surjection* $\mathbb{Q}_{p_k}^\times \twoheadrightarrow (\mathbb{Z}/2\mathbb{Z})_+$ of modules such that the composite $\{\pm 1\} \hookrightarrow \mathbb{Q}_{p_k}^\times \twoheadrightarrow (\mathbb{Z}/2\mathbb{Z})_+$ is an *isomorphism*. Thus, by considering the composite

$$\mathrm{Aut}(k) \hookrightarrow N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k)) \rightarrow \mathrm{Aut}_{\mathbb{Q}_{p_k}}(V_{-1}) = \mathbb{Q}_{p_k}^\times \twoheadrightarrow (\mathbb{Z}/2\mathbb{Z})_+,$$

we conclude that the subgroup $\mathrm{Aut}(k) \subseteq N_{\mathrm{Out}(G_k)}(\mathrm{Aut}(k))$ is a *direct summand*, as desired.

LEMMA 7.3. — *Suppose that $(p_k, a_k) \neq (2, 1)$. Write $\mathrm{Nm}_{k/k^{(d=1)}}^\wedge: (k^\times)^\wedge \rightarrow ((k^{(d=1)})^\times)^\wedge$ for the [necessarily open] homomorphism of abelian profinite groups induced by the Norm map $\mathrm{Nm}_{k/k^{(d=1)}}: k^\times \rightarrow (k^{(d=1)})^\times$. Then the following hold:*

(i) *The image of the [uniquely determined] pro- p_k -Sylow subgroup of the abelian profinite group $(k^\times)^\wedge$ by $\mathrm{Nm}_{k/k^{(d=1)}}^\wedge$ is a **free \mathbb{Z}_{p_k} -module of rank two**.*

(ii) *There exists a pro- p_k closed subgroup $M \subseteq \mathrm{Ker}(\mathrm{Nm}_{k/k^{(d=1)}}^\wedge)$ of the kernel of $\mathrm{Nm}_{k/k^{(d=1)}}^\wedge$ such that M is a **free \mathbb{Z}_{p_k} -module of rank $d_k - 1$** , and, moreover, the natural inclusion $M \hookrightarrow (k^\times)^\wedge$ is a **split injection**.*

PROOF. — First, we verify assertion (i). Write $(k^\times)^\wedge(p_k)$, $((k^{(d=1)})^\times)^\wedge(p_k)$ for the [uniquely determined] pro- p_k -Sylow subgroups of the abelian profinite groups $(k^\times)^\wedge$, $((k^{(d=1)})^\times)^\wedge$, respectively. Let us observe that it follows immediately from [3], Lemma 1.2, (i), that, to verify assertion (i), it suffices to verify that the image $\mathrm{Nm}_{k/k^{(d=1)}}^\wedge((k^\times)^\wedge(p_k)) \subseteq ((k^{(d=1)})^\times)^\wedge(p_k)$ is *torsion-free*. Thus, if p_k is *odd*, then since [one verifies easily from Proposition 1.1, (v), that] $((k^{(d=1)})^\times)^\wedge(p_k)$ is *torsion-free*, assertion (i) holds.

Suppose that $p_k = 2$. Then let us observe that since $((k^{(d=1)})^\times)_{\mathrm{tor}} = \{\pm 1\}$ [cf. Proposition 1.1, (v); [3], Lemma 1.2, (i)], it follows immediately from Lemma 4.1, (iii), that, to verify assertion (i) in the case where $p_k = 2$, it suffices to verify that the image of the inertia subgroup $I_k \subseteq G_k$ by $\chi_{p_k\text{-cyc}}: G_k \rightarrow \mathbb{Z}_{p_k}^\times$ [cf. Lemma 4.1, (i)] does *not contain* $-1 \in \mathbb{Z}_{p_k}^\times$. On the other hand, since [we have assumed that] $(p_k, a_k) \neq (2, 1)$, this follows immediately from the [easily verified] *injectivity* of the composite $\{\pm 1\} \hookrightarrow \mathbb{Z}_{p_k}^\times \twoheadrightarrow (\mathbb{Z}/4\mathbb{Z})^\times$. This completes the proof of assertion (i) in the case where $p_k = 2$, hence also of assertion (i).

Assertion (ii) follows immediately from assertion (i), together with [3], Lemma 1.2, (i). This completes the proof of Lemma 7.3. \square

LEMMA 7.4. — *Suppose that $(p_k, a_k) \neq (2, 1)$. Then there exists a Galois extension $k_\infty \subseteq \bar{k}$ of k such that $\mathrm{Gal}(k_\infty/k)$ is a **free \mathbb{Z}_{p_k} -module of rank $d_k - 1$** , and, moreover, the [uniquely determined] **maximal intermediate field of $k_\infty/k^{(d=1)}$ abelian over $k^{(d=1)}$ coincides with $k^{(\mathrm{ab})}$** [cf. Definition 4.2, (iii)].*

PROOF. — Let us first observe that one verifies immediately from [3], Lemma 1.7, (1), (2), that, to verify Lemma 7.4, it suffices to verify that there exists a *surjection* $\phi: (k^\times)^\wedge \twoheadrightarrow ((\mathbb{Z}_{p_k})_+)^{\oplus d_k - 1}$ of profinite groups such that if we write $\mathrm{Nm}_{k/k^{(d=1)}}^\wedge: (k^\times)^\wedge \rightarrow ((k^{(d=1)})^\times)^\wedge$ for the homomorphism induced by the Norm map $\mathrm{Nm}_{k/k^{(d=1)}}: k^\times \rightarrow (k^{(d=1)})^\times$,

then the equality $\mathrm{Nm}_{k/k^{(d=1)}}^{\wedge}((k^{\times})^{\wedge}) = \mathrm{Nm}_{k/k^{(d=1)}}^{\wedge}(\mathrm{Ker}(\phi))$ holds. On the other hand, this follows immediately from Lemma 7.3, (ii). This completes the proof of Lemma 7.4. \square

THEOREM 7.5. — *Suppose that a maximal intermediate field of $k/k^{(\mathrm{ab})}$ tamely ramified over $k^{(\mathrm{ab})}$ does not coincide with $k^{(d=1)} \subseteq k$ [which is the case if, for instance, $d_k^{(\mathrm{ab})} \neq 1$], and that $(p_k, a_k) \neq (2, 1)$. Let n be a nonnegative integer such that $[k : k^{(\mathrm{ab})}] \in p_k^n \mathbb{Z}$ and A an abelian p_k -group that satisfies the following two conditions:*

- (1) *It holds that $\sharp A = p_k^n$.*
- (2) *The finite abelian group A is generated by at most $(d_k/p_k^n) - 1$ elements.*

Then there exists a subgroup of $\mathrm{Out}(G_k)$ isomorphic to A .

PROOF. — Let K_1 be a maximal intermediate field of $k/k^{(\mathrm{ab})}$ tamely ramified over $k^{(\mathrm{ab})}$. Thus, it follows from Lemma 4.3, (ii), that the positive integer d_k/d_{K_1} is a power of p_k and $\geq p_k^n$. Then since [we have assumed that] $d_{K_1} \geq 2$, it follows immediately from Lemma 7.4 that there exists a finite Galois extension $K_2 \subseteq \bar{k}$ of K_1 such that $d_k = p_k^n \cdot d_{K_2}$ [i.e., that $d_k/d_{K_1} = p_k^n \cdot d_{K_2}/d_{K_1}$], and, moreover, $K_2^{(\mathrm{ab})} = K_1^{(\mathrm{ab})}$ ($= k^{(\mathrm{ab})}$ — cf. Lemma 4.3, (i)). Thus, it follows immediately — in light of condition (2) — from Lemma 7.4 that there exists a finite Galois extension $K_3 \subseteq \bar{k}$ of K_2 such that $\mathrm{Gal}(K_3/K_2)$ is isomorphic to A — which thus implies [cf. condition (1)] that $d_{K_3} = d_{K_2} \cdot \sharp A = d_{K_2} \cdot p_k^n = d_k$ — and, moreover, $K_3^{(\mathrm{ab})} = K_2^{(\mathrm{ab})}$ ($= k^{(\mathrm{ab})}$). In particular, since [we have assumed that] $(p_k, a_k) \neq (2, 1)$, it follows immediately — in light of Proposition 2.5, (i); Proposition 4.9, (i); [3], Proposition 3.6 — from Theorem 4.11, together with [3], Lemma 1.7, (1), (2), that G_k is isomorphic, as an abstract group, to $G_{K_3} \stackrel{\mathrm{def}}{=} \mathrm{Gal}(\bar{k}/K_3)$. Thus, by considering the image of the composite of the natural injection $\mathrm{Gal}(K_3/K_2) \hookrightarrow (\mathrm{Aut}(K_3) \hookrightarrow) \mathrm{Out}(G_{K_3})$ and the isomorphism $\mathrm{Out}(G_{K_3}) \xrightarrow{\sim} \mathrm{Out}(G_k)$ obtained by conjugation by some isomorphism $G_{K_3} \xrightarrow{\sim} G_k$, we obtain a subgroup of $\mathrm{Out}(G_k)$ isomorphic to A , as desired. This completes the proof of Theorem 7.5. \square

REMARK 7.5.1. — One concrete application of Theorem 7.5 is as follows: Suppose that p_k is odd. Let n be a positive integer. Suppose, moreover, that $k = \mathbb{Q}_{p_k}(\zeta_{p_k^n}, p_k^{1/p_k^n})$. Then one verifies immediately that

$$k^{(\mathrm{ab})} = \mathbb{Q}_{p_k}(\zeta_{p_k^n}), \quad d_k = p_k^{2n-1} \cdot (p_k - 1), \quad d_k^{(\mathrm{ab})} = p_k^{n-1} \cdot (p_k - 1).$$

Thus, it follows from Theorem 7.5 that, for positive integers d, r_1, \dots, r_d such that $r_1 + \dots + r_d = n$, there exists a subgroup of the group $\mathrm{Out}(G_k)$ isomorphic to the abelian p_k -group

$$\mathbb{Z}/p_k^{r_1} \mathbb{Z} \times \dots \times \mathbb{Z}/p_k^{r_d} \mathbb{Z}$$

[cf. the easily verified inequalities $d \leq n < d_k/p_k^n$].

Note that this observation in the case where $n = 2$ was already given in example (2) given in [4], §2.

REMARK 7.5.2. — One of motivations of studying Theorem 6.13, Theorem 7.2, and Theorem 7.5 is as follows:

(i) The *Neukirch-Uchida* theorem [cf. the main theorem of [16]] asserts that

(\dagger_{NF}) every outer isomorphism of profinite groups between the absolute Galois groups of number fields [i.e., finite extensions of \mathbb{Q}] arises from a *uniquely determined isomorphism between the number fields*,

which thus implies that

(\dagger_{NF}) the isomorphism class, i.e., as an abstract profinite group, of the absolute Galois group of a number field *completely determines* the isomorphism class, i.e., as an abstract field, of the number field.

On the other hand, it is well-known [cf., e.g., [3], Theorem 2.2] that *neither* the assertion (\dagger_{NF}) for *MLF's* nor the assertion (\dagger_{NF}) for *MLF's* holds. More precisely, for instance, if p is *odd*, then

(\dagger_{MLF}) there exists an outer automorphism of the absolute Galois group of \mathbb{Q}_p that is *nontrivial*, hence also does *not arise* from any automorphism of the field \mathbb{Q}_p [cf., e.g., the discussion given at the final portion of [12], Chapter VII, §5],

and, moreover,

(\dagger_{MLF}) there exist two MLF's k_\circ, k_\bullet such that the field k_\circ is *not isomorphic* to k_\bullet , but the absolute Galois group of k_\circ is *isomorphic* to the absolute Galois group of k_\bullet [cf., e.g., [17], §2, Theorem, (i)].

(ii) The assertion (\dagger_{MLF}) in (i) thus asserts that, in general [e.g., in the case where we take the “ k ” to be \mathbb{Q}_p , for some *odd* prime number p], the natural injection

$$\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$$

[cf., e.g., [3], Proposition 2.1] is *not bijective*. Under this state of affairs, one may consider the following problem:

Problem: Is there a certain “suitable” characterization of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$?

Here, let us observe that

Theorem 6.13, (ii), may be regarded as a certain *affirmative solution* to this problem, i.e., in the case where the MLF k is *Galois-specifiable* [cf. Definition 6.1].

(iii) From the point of view of the problem in (ii), let us observe

the [easily verified] *finiteness* of the group $\text{Aut}(k)$.

In particular, as one of possible solutions to the problem in (ii), one may discuss the following question:

(\ast_{fn}) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ the *uniquely determined maximal finite* subgroup of $\text{Out}(G_k)$? Put another way, is every element of $\text{Out}(G_k)$ of *finite order* contained in the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$?

Now let us observe that it is immediate that an *affirmative answer* to this question ($*_{\text{fin}}$) implies an *affirmative answer* to the following question ($*_{\text{char}}$), hence also an *affirmative answer* to the following question ($*_{\text{nor}}$):

($*_{\text{char}}$) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ *characteristic*?

($*_{\text{nor}}$) Is the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ *normal*?

an affir. sol. to ($*_{\text{fin}}$) \implies an affir. sol. to ($*_{\text{char}}$) \implies an affir. sol. to ($*_{\text{nor}}$)

(iv) Now let us observe that

Theorem 7.2 is related to the question ($*_{\text{nor}}$) in (iii),

and that

Theorem 7.5 [cf. also the example in Remark 7.5.1] yields a *negative answer* to the question ($*_{\text{fin}}$) in (iii).

(v) In §8, we will give a *negative answer* to the question ($*_{\text{nor}}$) in (iii), hence also [cf. the discussion of (iii)] *negative answers* to the questions ($*_{\text{fin}}$) and ($*_{\text{char}}$) in (iii) [cf. Corollary 8.7 below].

8. ON OUTER AUTOMORPHISMS ARISING FROM FIELD AUTOMORPHISMS II

In the present §8, we maintain the notational conventions introduced at the beginnings of §1 and §2. In particular, we have been given a group of *MLF-type*

$$G.$$

Let l be a prime number. Suppose, moreover, that

(a) $k^\times(G)[l] \neq \{1\}$,

(b) $d(G)/d^{(\text{ab})}(G) = l$ [cf. Definition 4.7, (iv)], and, moreover,

(c) $d^{(\text{ab})}(G) \notin l\mathbb{Z}$.

In the present §8, we give a *negative answer* to the question ($*_{\text{nor}}$) in Remark 7.5.2, (iii) [cf. Corollary 8.7 below].

LEMMA 8.1. — *Let $\Gamma \subseteq \text{Out}(G)$ be a **quasi-geometric** [cf. Definition 6.6, (ii)] subgroup of order l . Then the following hold:*

(i) *The group $G \rtimes^{\text{out}} \Gamma$ [cf. Definition 6.6, (i)] is of **AAMLF-type** [cf. Definition 4.8] whose **MLF-Galois label** [cf. Definition 4.10] coincides with*

$$\left(p(G), d^{(\text{ab})}(G), \text{Im}((\text{Nm}_{\text{abs}})(G)) \right)$$

[cf. Definition 4.7, (iii)].

(ii) *The isomorphism class of the group $G \rtimes^{\text{out}} \Gamma$ does **not depend** on the choice of Γ , i.e., depends only on (G, l) .*

(iii) *It holds that $e(G \rtimes^{\text{out}} \Gamma) = e(G)/l$, and that $k^\times(G \rtimes^{\text{out}} \Gamma)[l] \neq \{1\}$.*

(iv) *There exists a uniquely determined \mathbb{F}_l^\times -torsor*

$$T \subseteq k^\times(G \rtimes^{\text{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l$$

in the vector space $k^\times(G \rtimes^{\text{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l$ over \mathbb{F}_l that satisfies the following condition: Write $S \subseteq \bar{k}^\times(G)$ for the subset of $\bar{k}^\times(G)$ consisting of elements $\gamma \in \bar{k}^\times(G)$ ($\supseteq \bar{k}^\times(G)^{G \rtimes^{\text{out}} \Gamma} = k^\times(G \rtimes^{\text{out}} \Gamma)$) — cf. [3], Proposition 4.2, (i) of $\bar{k}^\times(G)$ such that the l -th power γ^l is **contained** in $k^\times(G \rtimes^{\text{out}} \Gamma)$ and, moreover, a **lifting** of an element of T , i.e., relative to the natural surjection $k^\times(G \rtimes^{\text{out}} \Gamma) \rightarrow k^\times(G \rtimes^{\text{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l$. Then the subset S is **nonempty**, and, moreover, for every element $\gamma \in S$ of S , the subgroup $G \subseteq G \rtimes^{\text{out}} \Gamma$ of $G \rtimes^{\text{out}} \Gamma$ **coincides** with the stabilizer, with respect to the natural action of $G \rtimes^{\text{out}} \Gamma$ on $\bar{k}^\times(G)$, of $\gamma \in S$.

PROOF. — First, we verify assertion (i). Let us first observe that it follows from [3], Proposition 3.6, that $p(G \rtimes^{\text{out}} \Gamma) = p(G)$ and $d(G \rtimes^{\text{out}} \Gamma) = d(G)/l = d^{(\text{ab})}(G)$ [cf. the condition (b) at the beginning of the present §8]. Thus, to verify assertion (i), it suffices to verify that $\text{Im}(\text{Nm}_{\text{abs}}(G)) = \text{Im}(\text{Nm}_{\text{abs}}(G \rtimes^{\text{out}} \Gamma))$. To this end, let us observe that it follows from Remark 4.9.1 that $\text{Im}(\text{Nm}_{\text{abs}}(G)) \subseteq \text{Im}(\text{Nm}_{\text{abs}}(G \rtimes^{\text{out}} \Gamma))$, and, moreover, $[\text{Im}(\text{Nm}_{\text{abs}}(G \rtimes^{\text{out}} \Gamma)) : \text{Im}(\text{Nm}_{\text{abs}}(G))] \in \{1, l\}$. Thus, it follows from the condition (c) at the beginning of the present §8 that $\text{Im}(\text{Nm}_{\text{abs}}(G)) = \text{Im}(\text{Nm}_{\text{abs}}(G \rtimes^{\text{out}} \Gamma))$, as desired. This completes the proof of assertion (i).

Assertion (ii) follows from assertion (i) and Theorem 4.11. Next, we verify assertion (iii). The first assertion of assertion (iii) follows immediately — in light of Proposition 4.9, (iii); [3], Proposition 3.6 — from assertion (i) and Lemma 4.3, (ii). Next, let us observe that it follows immediately from [3], Proposition 4.2, (i), that, to verify the second assertion of assertion (iii), it suffices to verify that the homomorphism $G \rtimes^{\text{out}} \Gamma \rightarrow \mathbb{F}_l^\times$ determined by the natural action of $G \rtimes^{\text{out}} \Gamma$ on $\bar{k}^\times(G)[l]$ is *trivial*. On the other hand, it follows from the condition (a) at the beginning of the present §8, together with [3], Proposition 4.2, (i), that the restriction to $G \subseteq G \rtimes^{\text{out}} \Gamma$ of the homomorphism $G \rtimes^{\text{out}} \Gamma \rightarrow \mathbb{F}_l^\times$ is *trivial*. Thus, the desired *triviality* follows from our assumption that Γ is of order l . This completes the proof of assertion (iii). Finally, since Γ is of order l , assertion (iv) follows immediately — in light of [3], Proposition 4.2, (i) — from *Kummer theory*, together with assertion (iii). This completes the proof of Lemma 8.1. \square

REMARK 8.1.1. — Suppose that p_k is *odd*. Let

$$a \in \mathbb{Q}_{p_k}^\times \setminus (\mathbb{Q}_{p_k}^\times)^{p_k}.$$

Suppose, moreover, that

$$k = \mathbb{Q}_{p_k}(\zeta_{p_k}, a^{1/p_k}).$$

Then, by the easily verified equality $(\mathbb{Q}_{p_k}^\times)^{p_k} = \mathbb{Q}_{p_k}^\times \cap (\mathbb{Q}_{p_k}(\zeta_{p_k})^\times)^{p_k}$, one verifies immediately that

$$k^{(\text{ab})} = \mathbb{Q}_{p_k}(\zeta_{p_k}), \quad d_k = p_k \cdot (p_k - 1), \quad d_k^{(\text{ab})} = p_k - 1$$

[cf. Definition 4.2, (iii), (iv)]. Thus, it follows from Proposition 4.9, (ii); [3], Proposition 3.6; [3], Proposition 3.11, (i), that the group G_k of MLF-type satisfies the three conditions (a), (b), and (c) at the beginning of the present §8 in the case where we take the prime number “ l ” to be p_k . Moreover, in this case, by Lemma 6.7, (i), and Proposition 6.8, the subgroup

$$\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k})) \subseteq (\mathrm{Aut}(k) = \mathrm{Gal}(k/\mathbb{Q}_{p_k}) \subseteq) \mathrm{Out}(G_k)$$

yields an example of a *quasi-geometric* subgroup of $\mathrm{Out}(G_k)$ of order p_k , i.e., as discussed in Lemma 8.1.

DEFINITION 8.2. — Let $\Gamma \subseteq \mathrm{Out}(G)$ be a quasi-geometric subgroup of order l .

(i) We shall write

$$T(\Gamma) \subseteq k^\times(G \rtimes^{\mathrm{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l$$

for the uniquely determined \mathbb{F}_l^\times -torsor “ T ” of Lemma 8.1, (iv).

(ii) We shall refer to an element of the subset “ S ” of Lemma 8.1, (iv), as a *Kummer generator* for Γ . Note that it follows from Lemma 8.1, (iv), that every Kummer generator for Γ is contained in $k^\times(G) (= \bar{k}^\times(G)^G \subseteq \bar{k}^\times(G))$.

(iii) We shall say that Γ is of *unit-Kummer type* if the image of the \mathbb{F}_l^\times -torsor $T(\Gamma) \subseteq k^\times(G \rtimes^{\mathrm{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l$ of (i) by the homomorphism

$$k^\times(G \rtimes^{\mathrm{out}} \Gamma) \otimes_{\mathbb{Z}} \mathbb{F}_l \longrightarrow (\mathbb{F}_l)_+$$

induced by $\mathrm{ord}_{\boxtimes}(G \rtimes^{\mathrm{out}} \Gamma)$ [cf. Definition 2.2] is $\{0\}$ [or, alternatively $\neq (\mathbb{F}_l)_+ \setminus \{0\}$].

(iv) Let $\Gamma_{\mathrm{st}} \subseteq \mathrm{Out}(G)$ be a strictly quasi-geometric [cf. Definition 6.6, (iii)] subgroup that contains Γ . Then we shall say that Γ is of *Γ_{st} -Kummer type* if there exists a Kummer generator $\gamma \in \bar{k}^\times(G)$ for Γ such that the l -th power $\gamma^l \in \bar{k}^\times(G)$ is contained in $k^\times(G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}}) (\subseteq k^\times(G \rtimes^{\mathrm{out}} \Gamma))$.

REMARK 8.2.1. — One verifies immediately from [3], Proposition 4.2, (i), together with the various definitions involved, that, in the situation of Definition 8.2, (iv), the following two conditions are equivalent:

(1) The quasi-geometric subgroup Γ is of *Γ_{st} -Kummer type*.

(2) There exists an element $a \in k(G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}})$ of the MLF $k(G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}})$ [cf. Definition 5.2, Remark 5.2.1] such that the subgroup $G \subseteq G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}}$ of $G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}}$ coincides with the intersection

$$(G \rtimes^{\mathrm{out}} \Gamma) \cap (G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}})(l; 1; l; a)$$

of the subgroup $G \rtimes^{\mathrm{out}} \Gamma \subseteq G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}}$ and the subgroup $(G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}})(l; 1; l; a) \subseteq G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}}$ of Definition 5.8, (ii), by the strictly radical data [cf. Definition 5.6, (i)] $(l; 1; l; a)$ for $k(G \rtimes^{\mathrm{out}} \Gamma_{\mathrm{st}})$.

REMARK 8.2.2. — Suppose that we are in the situation of Remark 8.1.1. Then one verifies immediately from the various definitions involved that the following hold:

(i) The $\mathbb{F}_{p_k}^\times$ -torsor

$$T\left(\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))\right) \subseteq k^\times \left(G_k \overset{\mathrm{out}}{\rtimes} \mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))\right) \otimes_{\mathbb{Z}} \mathbb{F}_{p_k} \xrightarrow{\sim} \mathbb{Q}_{p_k}(\zeta_{p_k})^\times \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$$

[cf. [3], Proposition 3.11, (i)] is given by the $\mathbb{F}_{p_k}^\times$ -torsor obtained by forming the $\mathbb{F}_{p_k}^\times$ -orbit of the image of $a \in \mathbb{Q}_{p_k}^\times \subseteq \mathbb{Q}_{p_k}(\zeta_{p_k})^\times$ in $\mathbb{Q}_{p_k}(\zeta_{p_k})^\times \otimes_{\mathbb{Z}} \mathbb{F}_{p_k}$.

(ii) The element $a^{1/p_k} \in k^\times \xrightarrow{\sim} k^\times(G_k)$ [cf. [3], Proposition 3.11, (i)] is a *Kummer generator* for the quasi-geometric subgroup $\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))$ of order p_k .

(iii) It holds that the quasi-geometric subgroup $\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))$ of order p_k is of *unit-Kummer type* if and only if $a \in \mathbb{Z}_{p_k}^\times \cdot (\mathbb{Q}_{p_k}^\times)^{p_k}$.

(iv) Let us observe that, by Proposition 6.8, the subgroup

$$\left(\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k})) \subseteq\right) \mathrm{Aut}(k) = \mathrm{Gal}(k/\mathbb{Q}_{p_k}) \subseteq \mathrm{Out}(G_k)$$

yields an example of a *strictly quasi-geometric* subgroup of $\mathrm{Out}(G_k)$ that *contains* the quasi-geometric subgroup $\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))$ of order p_k , i.e., as discussed in Definition 8.2, (iv). Moreover, in this case, since [we have assumed that] $a \in \mathbb{Q}_{p_k}^\times (= (k^\times)^{\mathrm{Aut}(k)})$, the quasi-geometric subgroup $\mathrm{Gal}(k/\mathbb{Q}_{p_k}(\zeta_{p_k}))$ of order p_k is of *Aut(k)-Kummer type*.

LEMMA 8.3. — Let $\Gamma, \Sigma \subseteq \mathrm{Out}(G)$ be **quasi-geometric** subgroups of order l . Suppose that Γ is **not of unit-Kummer type**. Let $\gamma \in k^\times(G)$ be a **Kummer generator** for Γ . Then the following hold:

(i) It holds that $\gamma \notin k^\times(G \overset{\mathrm{out}}{\rtimes} \Sigma)$.

(ii) Suppose that $\gamma^l \in k^\times(G \overset{\mathrm{out}}{\rtimes} \Sigma)$. Then γ is a **Kummer generator** for Σ .

(iii) Suppose that $\gamma^l \in k^\times(G \overset{\mathrm{out}}{\rtimes} \Sigma)$. Then the quasi-geometric subgroup Σ is **not of unit-Kummer type**.

PROOF. — First, we verify assertion (i). Let us first observe that since Γ is of order l and not of unit-Kummer type, it follows immediately from Proposition 2.3 that $\mathrm{ord}_{\boxtimes}(G)(\gamma) \notin l\mathbb{Z}$. On the other hand, since Σ is of order l , it follows immediately from Proposition 2.3 and Lemma 8.1, (iii), that $\mathrm{ord}_{\boxtimes}(G)(k^\times(G \overset{\mathrm{out}}{\rtimes} \Sigma)) = l\mathbb{Z}$. Thus, assertion (i) holds. This completes the proof of assertion (i).

Next, since $k^\times(G \overset{\mathrm{out}}{\rtimes} \Sigma)[l] \neq \{1\}$ [cf. Lemma 8.1, (iii)], and Σ is of order l , assertion (ii) follows immediately — in light of [3], Proposition 4.2, (i) — from assertion (i) and *Kummer theory*. Finally, since $\mathrm{ord}_{\boxtimes}(G)(\gamma) \notin l\mathbb{Z}$ [cf. the proof of assertion (i)], and Σ is of order l , assertion (iii) follows immediately — in light of Proposition 2.3 — from assertion (ii) and Lemma 8.1, (iii). This completes the proof of Lemma 8.3. \square

LEMMA 8.4. — Let $\Gamma_{\mathrm{st}} \subseteq \mathrm{Out}(G)$ be a **strictly quasi-geometric** subgroup of $\mathrm{Out}(G)$. Then the following hold:

(i) The group Γ_{st} has a **uniquely determined l -Sylow subgroup**. Moreover, the l -Sylow subgroup is **of order l** .

(ii) Let $\Sigma \subseteq \text{Out}(G)$ be a subgroup of $\text{Out}(G)$ such that $\Sigma \subseteq N_{\text{Out}(G)}(\Gamma_{\text{st}})$. Then it holds that

$$k^\times(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}) \subseteq k^\times(G)^\Sigma.$$

(iii) In the situation of (ii), suppose, moreover, that Σ is **quasi-geometric**. Then it holds that

$$k^\times(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}) \subseteq k^\times(G \overset{\text{out}}{\rtimes} \Sigma).$$

PROOF. — First, we verify assertion (i). Let us first observe that it follows from the conditions (b), (c) at the beginning of the present §8, together with Lemma 6.7, (iv), that each l -Sylow subgroup of Γ_{st} is *of order l* . Let $\Gamma_1, \Gamma_2 \subseteq \Gamma_{\text{st}}$ be l -Sylow subgroups of Γ_{st} [which thus implies that $\sharp\Gamma_1 = \sharp\Gamma_2 = l$]. Then since both Γ_1 and Γ_2 are *quasi-geometric* [cf. Lemma 6.7, (i)], it follows immediately from Lemma 8.1, (i), that $G \overset{\text{out}}{\rtimes} \Gamma_1 = G \overset{\text{out}}{\rtimes} \Gamma_2$, i.e., as subgroups of $G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}$. In particular, we obtain that $\Gamma_1 = \Gamma_2$, i.e., as subgroups of $\text{Out}(G)$, as desired. This completes the proof of assertion (i).

Next, since Γ_{st} is *strictly quasi-geometric* [which thus implies that $d(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}) = 1$], assertion (ii) follows immediately from Corollary 5.5. Finally, assertion (iii) follows from assertion (ii), together with Lemma 6.7, (ii). This completes the proof of Lemma 8.4. \square

THEOREM 8.5. — Let $\Gamma_{\text{st}} \subseteq \text{Out}(G)$ be a **strictly quasi-geometric** subgroup of $\text{Out}(G)$ and $\Sigma \subseteq \text{Out}(G)$ a **quasi-geometric** subgroup of order l . Write $\Gamma \subseteq \Gamma_{\text{st}}$ for the uniquely determined l -Sylow subgroup of Γ_{st} [cf. Lemma 8.4, (i)]. Suppose that the following three conditions are satisfied:

- (1) The subgroup $\Sigma \subseteq \text{Out}(G)$ **normalizes** the subgroup $\Gamma_{\text{st}} \subseteq \text{Out}(G)$.
- (2) The quasi-geometric subgroup Γ is **of Γ_{st} -Kummer type**.
- (3) The quasi-geometric subgroup Γ is **not of unit-Kummer type**.

Then the quasi-geometric subgroup Σ is **not of unit-Kummer type**.

PROOF. — It follows from Lemma 8.4, (iii), and condition (1) that $k^\times(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}) \subseteq k^\times(G \overset{\text{out}}{\rtimes} \Sigma)$. Now observe that since Γ is *of Γ_{st} -Kummer type* [cf. condition (2)], there exists a *Kummer generator* $\gamma \in k^\times(G)$ for Γ such that $\gamma^l \in k^\times(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}})$, which thus implies [cf. the above inclusion $k^\times(G \overset{\text{out}}{\rtimes} \Gamma_{\text{st}}) \subseteq k^\times(G \overset{\text{out}}{\rtimes} \Sigma)$] that $\gamma^l \in k^\times(G \overset{\text{out}}{\rtimes} \Sigma)$. Thus, since Γ is *not of unit-Kummer type* [cf. condition (3)], it follows from Lemma 8.3, (iii), that Σ is *not of unit-Kummer type*, as desired. This completes the proof of Theorem 8.5. \square

COROLLARY 8.6. — Suppose that p_k is **odd**. For each $\square \in \{\circ, \bullet\}$, let

$$a_\square \in \mathbb{Q}_{p_k}^\times \setminus (\mathbb{Q}_{p_k}^\times)^{p_k};$$

write

$$k_\square \stackrel{\text{def}}{=} \mathbb{Q}_{p_k}(\zeta_{p_k}, a_\square^{1/p_k}), \quad G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$$

— where \bar{k}_\square is an algebraic closure of k_\square . [So it follows from Theorem 4.11, together with Remark 8.1.1 — cf. also Proposition 4.9, (i); [3], Proposition 3.6 — that the group G_\circ is **isomorphic**, as an abstract group, to G_\bullet .] Write

$$\Phi: \text{Out}(G_\circ) \xrightarrow{\sim} \text{Out}(G_\bullet)$$

for the isomorphism obtained by conjugation by some fixed isomorphism $G_\circ \xrightarrow{\sim} G_\bullet$ of groups. Suppose that $a_\circ \in \mathbb{Z}_{p_k}^\times$ [e.g., $a_\circ = p_k + 1$] but $a_\bullet \notin \mathbb{Z}_{p_k}^\times$ [e.g., $a_\bullet = p_k$]. Then it holds that

$$\Phi\left(\text{Gal}(k_\circ/\mathbb{Q}_{p_k}(\zeta_{p_k}))\right) \not\subseteq N_{\text{Out}(G_\bullet)}(\text{Aut}(k_\bullet)).$$

PROOF. — This assertion follows immediately from Theorem 8.5, together with Remark 8.1.1 and Remark 8.2.2, (iii), (iv). \square

REMARK 8.6.1.

(i) Let us recall from Theorem 6.13, (i), that if k is *Galois-specifiable* [cf. Definition 6.1], then there is a *precisely one* $\text{Out}(G_k)$ -conjugacy class of *strictly quasi-geometric* subgroups of $\text{Out}(G_k)$.

(ii) Next, suppose that we are in the situation of Corollary 8.6. Then it follows immediately from Corollary 8.6 that $\Phi(\text{Aut}(k_\circ)) \neq \text{Aut}(k_\bullet)$. On the other hand, it follows from Proposition 6.8 that both $\Phi(\text{Aut}(k_\circ))$ and $\text{Aut}(k_\bullet)$ are *strictly quasi-geometric*. Thus, [since one may take the isomorphism “ $G_\circ \xrightarrow{\sim} G_\bullet$ ” of Corollary 8.6 to be an *arbitrary* isomorphism] there are *at least two* $\text{Out}(G_\bullet)$ -conjugacy classes of *strictly quasi-geometric* subgroups of $\text{Out}(G_\bullet)$. In particular — in light of Theorem 5.9, (ii) — we conclude that, in general, a similar assertion to Theorem 6.13, (i), for [the absolute Galois group of] an *absolutely characteristic* [cf. Definition 5.7] MLF does *not hold*.

COROLLARY 8.7. — *Suppose that p_k is odd, and that*

$$k = \mathbb{Q}_{p_k}(\zeta_{p_k}, p_k^{1/p_k}).$$

Then the subgroup

$$\text{Aut}(k) \subseteq \text{Out}(G_k)$$

is neither normally terminal nor normal.

PROOF. — Since k is *absolutely strictly radical* [cf. Definition 5.6, (iii)], hence also *absolutely characteristic* [cf. Theorem 5.9, (ii)], it follows from Theorem 7.2, (i), that the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$ is *not normally terminal*. Moreover, it follows immediately from Corollary 8.6 that the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ of $\text{Out}(G_k)$ is *not normal*. This completes the proof of Corollary 8.7. \square

REMARK 8.7.1. — In the present Remark 8.7.1, let us recall some of the discussions of the present §8 from the point of view of the notion of “link” [cf. [9], §2.7, (i)] as follows:

(i) Let us apply the notational conventions introduced in the statement of Corollary 8.6. In particular, the prime number p_k is *odd*. Moreover, for each $\square \in \{\circ, \bullet\}$, we are given an element

$$a_\square \in \mathbb{Q}_{p_k}^\times \setminus (\mathbb{Q}_{p_k}^\times)^{p_k},$$

an MLF

$$k_\square \stackrel{\text{def}}{=} \mathbb{Q}_{p_k}(\zeta_{p_k}, a_\square^{1/p_k}),$$

and an algebraic closure

$$\bar{k}_\square$$

of k_\square . Here, let us recall that the MLF k_\square — that is one of the main *arithmetic holomorphic* objects [cf. [9], §2.7, (vii)] in this discussion — determines some *mono-analytic* objects [cf. [9], §2.7, (vii)]. For instance, we have

- the group of *MLF-type*

$$G_\square \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square)$$

obtained by forming the absolute Galois group of k_\square and

- the *MLF $^\diamond$ -pair* [cf. [3], Definition 5.3]

$$G_\square \curvearrowright \bar{k}_\square^\times$$

obtained by considering the natural action of G_\square on \bar{k}_\square^\times .

In the remainder of Remark 8.7.1, suppose that we are in a situation in which

we are interested in a certain “*characterization*” of the element $a_\square \in \mathbb{Q}_{p_k}^\times$ from the point of view of such *mono-analytic* objects associated to the *arithmetic holomorphic* object k_\square .

More specifically, suppose that we are in a situation in which

we are interested in a certain “*comparison*” between a_\circ and a_\bullet via a suitable “*link*” that relates such *mono-analytic* objects associated to k_\circ and k_\bullet .

a “link”, i.e., a suitable isomorphism
 mono-an. obj. of $k_\circ \xrightarrow{\sim}$ mono-an. obj. of k_\bullet
 $\xRightarrow{?}$ a certain “comparison” between a_\circ and a_\bullet .

(ii) Let us start by observing that, for each $\square \in \{\circ, \bullet\}$, the group G_\square of MLF-type does *not give* any “*characterization*” of the element $a_\square \in \mathbb{Q}_{p_k}^\times$. Indeed, one verifies easily that

$$k_\square^{(\text{ab})} = \mathbb{Q}_{p_k}(\zeta_{p_k}), \quad d_{k_\square} = p_k \cdot (p_k - 1), \quad d_{k_\square}^{(\text{ab})} = p_k - 1.$$

Thus, it follows immediately from Theorem 4.11 [cf. also Proposition 4.9, (i); [3], Proposition 3.6] that the isomorphism class of the group “ G_\square ” does *not depend* on the choice of “ a_\square ”. In particular, we have an *isomorphism* of groups

$$G_\circ \xrightarrow{\sim} G_\bullet.$$

As a consequence, one may conclude that

one *cannot obtain* “any information about a_\square ” if one considers only “ G_\square ”.

(iii) In order to obtain a certain “*comparison*” between a_\circ and a_\bullet , let us relate *mono-analytic* portions of the *arithmetic holomorphic* structures of k_\circ and k_\bullet as follows: In the remainder of Remark 8.7.1, let us fix an isomorphism of [abstract] groups [i.e., between *Frobenius-like portions* — cf. [3], Definition 5.4]

$$\alpha_{\text{Fr}}: \bar{k}_\circ^\times \xrightarrow{\sim} \bar{k}_\bullet^\times$$

such that the isomorphism $\text{Aut}(\bar{k}_\circ^\times) \xrightarrow{\sim} \text{Aut}(\bar{k}_\bullet^\times)$ obtained by conjugation by α_{Fr} *restricts* to an isomorphism of $G_\circ \subseteq \text{Aut}(\bar{k}_\circ^\times)$ with $G_\bullet \subseteq \text{Aut}(\bar{k}_\bullet^\times)$. Write

$$\alpha_{\text{ét}}: G_\circ \xrightarrow{\sim} G_\bullet$$

for the resulting isomorphism [i.e., between *étale-like portions* — cf. [3], Definition 5.4]. [So the pair $(\alpha_{\text{Fr}}, \alpha_{\text{ét}})$ determines an isomorphism

$$(\alpha_{\text{Fr}}, \alpha_{\text{ét}}): (G_\circ \curvearrowright \bar{k}_\circ^\times) \xrightarrow{\sim} (G_\bullet \curvearrowright \bar{k}_\bullet^\times)$$

of MLF^\diamond -pairs — cf. [3], Definition 5.1, (ii).] In particular, roughly speaking, we are in a situation in which the collection of data “ $G_\square \curvearrowright \bar{k}_\square^\times$ ” may be regarded as a *coric* object [cf. [9], §2.7, (iv)] of our “link” [i.e., the pair $(\alpha_{\text{Fr}}, \alpha_{\text{ét}})$].

Note that we have natural inclusions of groups

$$G_\square \subseteq G_\square^\Gamma \stackrel{\text{def}}{=} \text{Gal}(\bar{k}_\square/k_\square^{(\text{ab})}) \subseteq \text{Aut}(\bar{k}_\square) \subseteq \text{Aut}(\bar{k}_\square^\times).$$

Write

$$k_\square^{(d=1)} \stackrel{\text{def}}{=} (\bar{k}_\square)^{\text{Aut}(\bar{k}_\square)}.$$

[So $k_\square^{(d=1)} = \mathbb{Q}_{p_k}$ in \bar{k}_\square .] Thus, we have a natural identification

$$\text{Aut}(\bar{k}_\square) = \text{Gal}(\bar{k}_\square/k_\square^{(d=1)}).$$

In particular, the group $\text{Aut}(\bar{k}_\square)$ is a group of *MLF-type* such that $d(\text{Aut}(\bar{k}_\square)) = 1$ [cf. [3], Proposition 3.6].

(iv) Before proceeding, we pause to recall some of the discussions of §5 from the point of view of this situation. Let us recall that the *reconstruction algorithms* of Definition 5.2 and Definition 5.8, (i), assert that, in this situation,

(*) the $\text{MLF } k_\square^{(d=1)}$, as well as the collection of *strictly radical data* for the $\text{MLF } k_\square^{(d=1)}$ [cf. Definition 5.6, (i)] each member of which yields the *absolutely strictly radical* $\text{MLF } k_\square$ [e.g., the strictly radical data $(p_k; 1; p_k; a_\square)$ for $k_\square^{(d=1)}$], is “*intrinsic*” from the point of view of the collection of data

$$G_\square \hookrightarrow \text{Aut}(\bar{k}_\square) \curvearrowright \bar{k}_\square^\times$$

— i.e., the MLF^\diamond -pair $\text{Aut}(\bar{k}_\square) \curvearrowright \bar{k}_\square^\times$ equipped with the subgroup $G_\square \subseteq \text{Aut}(\bar{k}_\square)$ of the étale-like portion $\text{Aut}(\bar{k}_\square)$.

Here, suppose that the “link” of (iii) [i.e., the pair $(\alpha_{\text{Fr}}, \alpha_{\text{ét}})$] satisfies the condition that

(†₁): the isomorphism $\text{Aut}(\bar{k}_\circ^\times) \xrightarrow{\sim} \text{Aut}(\bar{k}_\bullet^\times)$ obtained by conjugation by α_{Fr} also *restricts* to an isomorphism of $\text{Aut}(\bar{k}_\circ) \subseteq \text{Aut}(\bar{k}_\circ^\times)$ with $\text{Aut}(\bar{k}_\bullet) \subseteq \text{Aut}(\bar{k}_\bullet^\times)$.

[Put another way, roughly speaking, we are in a situation in which the collection of data “ $\text{Aut}(\bar{k}_\square) \curvearrowright \bar{k}_\square^\times$ ” may be regarded as a *coric* object of our “link”.] Then we conclude immediately from the above (*) that

the $((k_{\circ}^{(d=1)})^{\times})^{p_k}$ -orbit of $a_{\circ} \in (k_{\circ}^{(d=1)})^{\times}$ *coincides* — relative to some *isomorphism* $k_{\circ} \xrightarrow{\sim} k_{\bullet}$ of fields [i.e., determined by either the restriction $\alpha_{\text{Fr}}|_{(k_{\circ}^{(d=1)})^{\times}}$ or the composite of $\alpha_{\text{Fr}}|_{(k_{\circ}^{(d=1)})^{\times}}$ and the automorphism of $(k_{\bullet}^{(d=1)})^{\times}$ given by “ $x \mapsto x^{-1}$ ” — cf. [3], Theorem 7.6, (ii)] — with the $((k_{\bullet}^{(d=1)})^{\times})^{p_k}$ -orbit of $a_{\bullet} \in (k_{\bullet}^{(d=1)})^{\times}$,

i.e., obtain a certain “*comparison*” between a_{\circ} and a_{\bullet} . As a result,

the field k_{\circ} is *isomorphic*, as an abstract field, to the field k_{\bullet} .

This is precisely what is achieved by the application of the “*tautological*” assertion in Remark 5.9.3, (i), to *absolutely strictly radical* MLF’s [cf. Remark 5.9.3, (ii)].

Put another way, roughly speaking,

in the situation of (iv), one may *characterize* the element a_{\square} up to the *indeterminacies* arise from the action of the group $((k_{\square}^{(d=1)})^{\times})^{p_k}$.

$$\begin{array}{ccc} \text{Aut}(\bar{k}_{\circ}) & \xrightarrow{\sim} & \text{Aut}(\bar{k}_{\bullet}) \\ \curvearrowright & & \curvearrowright \\ \bar{k}_{\circ}^{\times} & \xrightarrow{\sim} & \bar{k}_{\bullet}^{\times} \end{array} \implies a_{\circ} = a_{\bullet} \text{ up to } ((k_{\circ}^{(d=1)})^{\times})^{p_k} \xrightarrow{\sim} ((k_{\bullet}^{(d=1)})^{\times})^{p_k}$$

(v) Let us return to the situation of the present §8. Next, suppose that the “link” of (iii) [i.e., the pair $(\alpha_{\text{Fr}}, \alpha_{\text{ét}})$] satisfies the condition that

(†₂): the action of the subgroup $G_{\circ}^{\Gamma} \subseteq \text{Aut}(\bar{k}_{\circ}^{\times})$ on $\text{Aut}(\bar{k}_{\bullet}^{\times})$ by conjugation — relative to the isomorphism $\text{Aut}(\bar{k}_{\circ}^{\times}) \xrightarrow{\sim} \text{Aut}(\bar{k}_{\bullet}^{\times})$ by α_{Fr} — *preserves* the subgroup $\text{Aut}(\bar{k}_{\bullet}) \subseteq \text{Aut}(\bar{k}_{\bullet}^{\times})$.

[Note that one verifies easily that the condition (†₁) in (iv) implies this condition (†₂).]

In this situation, we are *not given* any isomorphism of $\text{Aut}(\bar{k}_{\circ})$ with $\text{Aut}(\bar{k}_{\bullet})$. [Put another way, roughly speaking, we cannot regard the collection of data “ $\text{Aut}(\bar{k}_{\square}) \curvearrowright \bar{k}_{\square}^{\times}$ ” as a *coric* object of our “link”.] In particular, we *cannot apply* the *reconstruction algorithm* of Definition 5.8, (i). Nevertheless, Theorem 8.5 allows one to conclude that

if the $((k_{\circ}^{(d=1)})^{\times})^{p_k}$ -orbit of $a_{\circ} \in (k_{\circ}^{(d=1)})^{\times}$ contains a *unit* of $\mathcal{O}_{k_{\circ}^{(d=1)}}$, then the $((k_{\bullet}^{(d=1)})^{\times})^{p_k}$ -orbit of $a_{\bullet} \in (k_{\bullet}^{(d=1)})^{\times}$ contains a *unit* of $\mathcal{O}_{k_{\bullet}^{(d=1)}}$,

which thus implies that

$$\text{if } \text{ord}_{k_{\circ}^{(d=1)}}(a_{\circ}) = 0, \text{ then } \text{ord}_{k_{\bullet}^{(d=1)}}(a_{\bullet}) \in p_k \mathbb{Z}.$$

In particular, we obtain a certain “*comparison*” between a_{\circ} and a_{\bullet} .

$$\begin{array}{ccc} G_{\circ}^{\Gamma} \supseteq G_{\circ} & \xrightarrow{\sim} & G_{\bullet} \subseteq \text{Aut}(\bar{k}_{\bullet}) \\ \curvearrowright & & \curvearrowright \\ \bar{k}_{\circ}^{\times} & \xrightarrow{\sim} & \bar{k}_{\bullet}^{\times}, \end{array} \quad G_{\circ}^{\Gamma} \curvearrowright \bar{k}_{\circ}^{\times} \text{ preserves } \text{Aut}(\bar{k}_{\bullet})$$

$$\text{Then: } \quad \text{ord}_{k_{\circ}^{(d=1)}}(a_{\circ}) = 0 \implies \text{ord}_{k_{\bullet}^{(d=1)}}(a_{\bullet}) \in p_k \mathbb{Z}$$

(vi) Finally, in order to obtain an application of the conclusion of the discussion of (v), let us take the “ (a_{\circ}, a_{\bullet}) ” to be $(p_k + 1, p_k)$. Then it follows from the discussion of (ii) that the group G_{\circ} is *isomorphic* to the group G_{\bullet} . Thus, by applying a technique of

mono-anabelian transport [cf. [3], Theorem 7.6, (i); also [3], Remark 7.6.1, (i)], we obtain a “link”

$$(\alpha_{\text{Fr}}, \alpha_{\text{ét}}): (G_{\circ} \curvearrowright \bar{k}_{\circ}^{\times}) \xrightarrow{\sim} (G_{\bullet} \curvearrowright \bar{k}_{\bullet}^{\times})$$

as in (iii). In particular, since $(a_{\circ}, a_{\bullet}) = (p_k + 1, p_k)$, it follows from the conclusion of the discussion of (v) that the “link” does *not satisfy* the condition (\dagger_2) in (v), i.e., that

the action of G_{\circ}^{Γ} on $\text{Aut}(\bar{k}_{\circ}^{\times})$ by conjugation — relative to the isomorphism $\text{Aut}(\bar{k}_{\circ}^{\times}) \xrightarrow{\sim} \text{Aut}(\bar{k}_{\bullet}^{\times})$ by α_{Fr} — does *not preserve* the subgroup $\text{Aut}(\bar{k}_{\bullet}) \subseteq \text{Aut}(\bar{k}_{\bullet}^{\times})$.

Therefore, we conclude that

the subgroup $\text{Aut}(\bar{k}_{\bullet}) \subseteq \text{Aut}(G_{\bullet})$ is *not normal*,

which thus implies — by considering the respective quotients by G_{\bullet} — that

the subgroup $\text{Aut}(k_{\bullet}) \subseteq \text{Out}(G_{\bullet})$ is *not normal*.

This situation is precisely the situation formulated in the “nonnormal portion” of Corollary 8.7.

REFERENCES

- [1] Y. Hoshi, A note on the geometricity of open homomorphisms between the absolute Galois groups of p -adic local fields, *Kodai Math. J.* **36** (2013), no. **2**, 284–298.
- [2] Y. Hoshi, *Mono-anabelian reconstruction of number fields*, RIMS Preprint **1819** (March 2015).
- [3] Y. Hoshi, *Introduction to mono-anabelian geometry*, RIMS Preprint **1868** (January 2017).
- [4] M. Jarden and J. Ritter, On the characterization of local fields by their absolute Galois groups, *J. Number Theory* **11** (1979), no. **1**, 1–13.
- [5] M. Jarden and J. Ritter, Normal automorphisms of absolute Galois groups of p -adic fields, *Duke Math. J.* **47** (1980), no. **1**, 47–56.
- [6] W. Jenkner, Les corps p -adiques dont les groupes de Galois absolus sont isomorphes, *Journées Arithmétiques*, 1991 (Geneva). *Astérisque* No. **209** (1992), **14**, 221–226.
- [7] S. Mochizuki, Topics in absolute anabelian geometry I: generalities, *J. Math. Sci. Univ. Tokyo* **19** (2012), no. **2**, 139–242.
- [8] S. Mochizuki, Topics in absolute anabelian geometry III: global reconstruction algorithms, *J. Math. Sci. Univ. Tokyo* **22** (2015), no. **4**, 939–1156.
- [9] S. Mochizuki, *The mathematics of mutually alien copies: from Gaussian integrals to inter-universal Teichmüller Theory*, RIMS Preprint **1854** (July 2016).
- [10] N. Nakagoshi, The structure of the multiplicative group of residue classes modulo \mathfrak{p}^{N+1} , *Nagoya Math. J.* **73** (1979), 41–60.
- [11] J. Neukirch, *Algebraic number theory*. Translated from the 1992 German original and with a note by Norbert Schappacher. With a foreword by G. Harder, *Grundlehren der Mathematischen Wissenschaften*, **322**. Springer-Verlag, Berlin, 1999.
- [12] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*. Second edition, *Grundlehren der Mathematischen Wissenschaften*, **323**. Springer-Verlag, Berlin, 2008.
- [13] J. Ritter, \mathfrak{P} -adic fields having the same type of algebraic extensions, *Math. Ann.* **238** (1978), no. **3**, 281–288.
- [14] J.-P. Serre, *Local fields*. Translated from the French by Marvin Jay Greenberg, *Graduate Texts in Mathematics*, **67**. Springer-Verlag, New York-Berlin, 1979.
- [15] J.-P. Serre, *Abelian l -adic representations and elliptic curves*. With the collaboration of Willem Kuyk and John Labute, Revised reprint of the 1968 original. *Research Notes in Mathematics*, **7**. A K Peters, Ltd., Wellesley, MA, 1998.
- [16] K. Uchida, Isomorphisms of Galois groups, *J. Math. Soc. Japan* **28** (1976), no. **4**, 617–620.

-
- [17] S. Yamagata, A counterexample for the local analogy of a theorem by Iwasawa and Uchida, *Proc. Japan Acad.* **52** (1976), no. **6**, 276–278.

(Yuichiro Hoshi) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

E-mail address: `yuichiro@kurims.kyoto-u.ac.jp`