

RIMS-1894

**Homotopy Sequences for Varieties over Curves**

By

Yuichiro Hoshi

November 2018



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

---

# HOMOTOPY SEQUENCES FOR VARIETIES OVER CURVES

YUICHIRO HOSHI

NOVEMBER 2018

---

ABSTRACT. — In the present paper, we discuss the homotopy sequences for varieties over curves. We prove that, for instance, for a morphism from a normal variety to a certain smooth curve over a field of characteristic zero, it holds that the induced outer homomorphism between the étale fundamental groups satisfies some group-theoretic conditions if and only if the homotopy sequence for the morphism of varieties is exact. Moreover, we also give a refinement of a known result concerning the Grothendieck conjecture for hyperbolic polycurves of dimension two by means of our study of homotopy sequences.

## CONTENTS

INTRODUCTION .....	1
§1. SOME GENERALITIES ON HOMOTOPY SEQUENCES .....	4
§2. HOMOTOPY SEQUENCES FOR VARIETIES OVER CURVES .....	8
§3. AN ANABELIAN RESULT ON HYPERBOLIC POLYCURVES OF DIMENSION TWO .....	15
REFERENCES .....	20

## INTRODUCTION

The exactness of the homotopy sequences for certain *proper* morphisms was established in [15], Exposé X, Corollaire 1.4. Moreover, *K. Mitsui* [cf. [10], Theorem 1.1; [10], Theorem 4.22] and the author of the present paper [cf. [7], Theorem 2; [8], Proposition 1.10, (i)] have established the exactness of the homotopy sequences for certain *not necessarily proper* morphisms. [Note that, in [7], Theorem 2, the morphism “ $f^{\log}$ ” of log schemes is assumed to be *proper*; on the other hand, by applying the *log purity theorem* of [7], Proposition B.7 — that gives a natural outer isomorphism between the log fundamental group of a suitable log scheme and the étale fundamental group of the interior of the log scheme — one may derive, from [7], Theorem 2, the exactness of the homotopy sequences for certain *not necessarily proper* morphisms of schemes — i.e., as opposed to log schemes — in suitable situations.] In the present paper, we discuss a “*group-theoretic*” approach to study of homotopy sequences. Here, let us observe that each of the above results concerning the exactness of homotopy sequences gives us a statement of the following form:

---

2010 MATHEMATICS SUBJECT CLASSIFICATION. — 14F35.

KEY WORDS AND PHRASES. — homotopy sequence.

If a morphism of schemes satisfies suitable *scheme-theoretic conditions*, then the homotopy sequence for the morphism of schemes is *exact*.

On the other hand, one main purpose of the study of the present paper is to give a statement of the following form:

For a morphism of schemes, if the induced outer homomorphism between the étale fundamental groups satisfies suitable *group-theoretic conditions*, then the homotopy sequence for the morphism of schemes is *exact*.

Note that a requirement for such a statement arises naturally in, for instance, an attempt to solve the *Grothendieck conjecture for hyperbolic polycurves* [cf., e.g., the various arguments of [8], [13]; also the arguments of §3 of the present paper].

Here, let us explain a property

$$\mathbb{P}_{\mathcal{F} \rightarrow \infty}$$

of a profinite group [cf. Definition 2.5, (i)] that plays an important role in the present paper. We shall say that a profinite group  $G$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$  if, for an arbitrary open subgroup  $H \subseteq G$  of  $G$ , there exists a prime number  $l_H$  such that  $H$  has no quotient which is free pro- $l_H$  and not topologically finitely generated. One verifies easily that if a profinite group  $G$  has one of the following three properties, then  $G$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$  [cf. Remark 2.5.1]:

- The profinite group  $G$  is *topologically finitely generated*.
- An open subgroup of the profinite group  $G$  is *abelian*.
- An open subgroup of the profinite group  $G$  is *pro- $\Sigma$*  for some set  $\Sigma$  of prime numbers *not equal* to the set of all prime numbers.

The first main result — in the case where the base field is *of characteristic zero* — is as follows [cf. Theorem 2.8, also Remark 2.8.1]. In the present paper, if  $X$  is a connected locally noetherian scheme, then we shall write

$$\Pi_X$$

for the étale fundamental group of  $X$  [for some choice of basepoint] [cf. Definition 1.1, (i)]; moreover, if  $f$  is a morphism between connected locally noetherian schemes, then we shall write

$$\Delta_f$$

for the kernel of the outer homomorphism between the étale fundamental groups induced by  $f$  [cf. Definition 1.1, (ii)].

**THEOREM A.** — *Let  $k$  be a field of characteristic zero,  $S$  a smooth curve over  $k$  [cf. Definition 2.2, (ii)] whose base-change to an algebraic closure of  $k$  is isomorphic neither to the projective line nor to the affine line,  $X$  a normal variety over  $k$  [cf. Definition 2.2, (i)],  $f: X \rightarrow S$  a morphism over  $k$ , and  $\bar{\eta} \rightarrow S$  a geometric generic point of  $S$ . Then the following four conditions are equivalent:*

(1) *The outer homomorphism  $\Pi_X \twoheadrightarrow \Pi_S$  induced by  $f$  is **surjective**. Moreover, the profinite group  $\Delta_f$  is **topologically finitely generated**.*

(2) *The outer homomorphism  $\Pi_X \twoheadrightarrow \Pi_S$  induced by  $f$  is **surjective**. Moreover, the profinite group  $\Delta_f$  **has the property**  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$  [cf. Definition 2.5, (i)].*

(3) *The morphism  $f$  is **surjective** and **generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes*

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

*induces an **exact** sequence of profinite groups*

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

(4) *The morphism  $f$  is **dominant** and **generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes*

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

*induces an **exact** sequence of profinite groups*

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

In §3 of the present paper, we apply [some considerations that lead us to] Theorem A to the study of the *Grothendieck conjecture for hyperbolic polycurves*. The second main result of the present paper is as follows [cf. Theorem 3.3].

**THEOREM B.** — *Let  $p$  be a prime number,  $k$  a **sub- $p$ -adic field** [cf., e.g., [8], Definition 3.1],  $\bar{k}$  an algebraic closure of  $k$ ,  $X$  a **hyperbolic polycurve of dimension 2** over  $k$  [cf. [8], Definition 2.1, (ii)], and  $Z$  a **normal variety** over  $k$  [cf. Definition 2.2, (i)]. Write  $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ . Let*

$$\alpha: \Pi_Z \longrightarrow \Pi_X$$

*be an **open** homomorphism of profinite groups over  $G_k$ . Suppose that the kernel of  $\alpha$  has the property  $\mathbb{P}_{\bar{\mathbb{Z}} \rightarrow \infty}$  [cf. Definition 2.5, (i)]. Then there exists a **uniquely determined dominant** morphism*

$$Z \longrightarrow X$$

*over  $k$  from which  $\alpha$  arises. In particular, the normal variety  $Z$  is of **dimension  $\geq 2$** .*

Note that Theorem B may be regarded as a *refinement* of [8], Theorem A, in the case where condition (2) is satisfied [cf. Remark 3.3.1]. Finally, in §3, we observe that if one drops one of

- the assumption that the hyperbolic polycurve  $X$  is of *dimension 2* and
- the assumption that the kernel of  $\alpha$  has the property  $\mathbb{P}_{\bar{\mathbb{Z}} \rightarrow \infty}$

in the statement of Theorem B, then the conclusion of Theorem B no longer holds in general [cf. Remark 3.3.2; Remark 3.3.4, (i)].

## ACKNOWLEDGMENTS

The author would like to thank *Ippeï Nagamachi* for some helpful discussions. This research was supported by JSPS KAKENHI Grant Number 18K03239 and by the Research Institute for Mathematical Sciences, a Joint Usage/Research Center located in Kyoto University.

## 1. SOME GENERALITIES ON HOMOTOPY SEQUENCES

In the present §1, we discuss some generalities on homotopy sequences [cf. Lemma 1.3 and Lemma 1.5 below].

**DEFINITION 1.1.**

- (i) Let  $X$  be a connected locally noetherian scheme. Then we shall write

$$\Pi_X$$

for the étale fundamental group of  $X$  [for some choice of basepoint].

- (ii) Let  $X, Y$  be connected locally noetherian schemes;  $f: X \rightarrow Y$  a morphism of schemes. Then we shall write

$$\Delta_f = \Delta_{X/Y} \subseteq \Pi_X$$

for the kernel of the outer homomorphism  $\Pi_X \rightarrow \Pi_Y$  induced by  $f$ .

**DEFINITION 1.2.** — We shall say that a scheme is of *CLN-type* if the scheme is connected, locally noetherian, and normal.

**LEMMA 1.3.** — *Let  $X, S$  be schemes of CLN-type;  $f: X \rightarrow S$  a morphism of schemes which is of finite type, separated, and dominant;  $\bar{\eta} \rightarrow S$  a geometric generic point of  $S$ . Suppose that the following two conditions are satisfied:*

- (a) *The outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is **surjective**.*
- (b) *The geometric generic fiber  $X \times_S \bar{\eta}$  of  $f$  is **reduced** [cf. Remark 1.3.1 below].*

*Consider the following four conditions:*

(1) *Let  $Y \rightarrow X$  be a connected finite étale covering of  $X$ . Then the normalization of  $S$  in  $Y$  is **étale** over  $S$ .*

(2) *Let  $Y \rightarrow X$  be a connected finite étale covering of  $X$ . If the base-change  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$  of  $Y \rightarrow X$  by  $\bar{\eta} \rightarrow S$  has a **splitting**, then the normalization of  $S$  in  $Y$  is **étale** over  $S$ .*

(3) *The morphism  $f$  is **generically geometrically irreducible**. Moreover, for a connected finite étale covering  $Y \rightarrow X$  of  $X$ , if the base-change  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$  of  $Y \rightarrow X$  by  $\bar{\eta} \rightarrow S$  has a **splitting**, then there exists a connected finite étale covering of  $S$  whose pull-back by  $X \rightarrow S$  is **isomorphic** to  $Y$  over  $X$ .*

(4) *The morphism  $f$  is **generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes*

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

*induces an **exact** sequence of profinite groups*

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

*Then the implications*

$$(1) \implies (2) \iff (3) \iff (4)$$

*hold.*

PROOF. — The implication

$$(1) \implies (2)$$

is immediate. Next, we verify the implication

$$(2) \implies (3).$$

To this end, suppose that condition (2) is satisfied, and take a connected finite étale covering  $Y \rightarrow X$  of  $X$  whose base-change  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$  by  $\bar{\eta} \rightarrow S$  has a *splitting*. Now let us write

$$T \longrightarrow S$$

for the normalization of  $S$  in  $Y$ . Let us first verify the following claim:

Claim 1.3.A: The normalization  $T \rightarrow S$  of  $S$  in  $Y$  is a *connected finite étale covering* of  $S$ .

To this end, let us observe that it is immediate that  $T$  is *connected*. Next, let us observe that it follows from [2], Proposition 4.6.1, together with the condition (b), that the function field of  $X$ , hence also of  $Y$ , is *separable* over the function field of  $S$ . Thus, it follows from [9], §33, Lemma 1, that the morphism  $T \rightarrow S$  is *finite*. In particular, since [it follows from condition (2) that]  $T \rightarrow S$  is *étale*, we conclude that Claim 1.3.A holds, as desired. This completes the proof of Claim 1.3.A.

Next, we verify the following claim:

Claim 1.3.B: The morphism  $f$  is *generically geometrically irreducible*.

To this end, let us observe that it follows from Claim 1.3.A [i.e., in the case where we take “ $Y$ ” of Claim 1.3.A to be  $X$ ] that the normalization of  $S$  in  $X$  is a *connected finite étale covering* of  $S$ . Thus, since the outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is *surjective* [cf. the condition (a)], we conclude that the normalization of  $S$  in  $X$  *coincides* with  $S$ , which thus implies that the function field of  $S$  is *algebraically closed* in the function field of  $X$ . In particular, it follows from [2], Proposition 4.5.9, and [3], Proposition 9.7.8, that  $f$  is *generically geometrically irreducible*, as desired. This completes the proof of Claim 1.3.B.

Next, we verify the following claim:

Claim 1.3.C: The fiber product  $X \times_S T$  is *of CLN-type*.

To this end, let us observe that one verifies easily from Claim 1.3.A that  $X \times_S T$  is *locally noetherian* and *normal*. Next, it follows from the *surjectivity* of the outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  [cf. the condition (a)], together with Claim 1.3.A, that  $X \times_S T$  is *connected*. This completes the proof of Claim 1.3.C.

Next, we verify the following claim:

Claim 1.3.D: The natural morphism  $Y \rightarrow X \times_S T$  is a *connected finite étale covering* [hence *closed* and *open*; thus, this morphism  $Y \rightarrow X \times_S T$  is *surjective* — cf. Claim 1.3.C].

To this end, let us first observe that since both  $Y$  and  $X \times_S T$  are *finite* over  $X$  [cf. Claim 1.3.A], it follows that the morphism  $Y \rightarrow X \times_S T$  is *finite*. In particular, since [one verifies easily that] both  $Y \rightarrow X$  and  $X \times_S T \rightarrow X$  are *surjective*, by considering the fibers of  $Y \rightarrow X \times_S T \rightarrow X$  at the generic point of  $X$  [cf. also Claim 1.3.C], we conclude that the morphism  $Y \rightarrow X \times_S T$  is *dominant*. On the other hand, since  $Y \rightarrow X$  is *unramified*, it follows from [4], Proposition 17.3.3, (v), that the morphism  $Y \rightarrow X \times_S T$  is *unramified*. Thus, since  $X \times_S T$  is *normal* [cf. Claim 1.3.C], it follows from [15], Exposé I, Corollaire 9.11, that the morphism  $Y \rightarrow X \times_S T$  is *étale*, as desired. This completes the proof of Claim 1.3.D.

Finally, we verify the following claim:

Claim 1.3.E: The natural morphism  $Y \rightarrow X \times_S T$  is an *isomorphism*.

To this end, write  $d$  for the degree of the connected finite covering  $T \rightarrow S$  [cf. Claim 1.3.A]. Then since  $X \times_S \bar{\eta}$  is *connected* [cf. Claim 1.3.B], one verifies easily that the number of the connected components of  $(X \times_S T) \times_S \bar{\eta}$  is *equal* to  $d$ . Moreover, since the geometric fiber of  $Y \rightarrow T$  at an arbitrary geometric generic point of  $T$  is *connected* [cf. [2], Proposition 4.5.9], the number of the connected components of  $Y \times_S \bar{\eta}$  is *equal* to  $d$ . Thus, since  $Y \rightarrow X \times_S T$  is *surjective* [cf. Claim 1.3.D], this morphism  $Y \rightarrow X \times_S T$  determines a *bijection* between the set of connected components of  $Y \times_S \bar{\eta}$  and the set of connected components of  $(X \times_S T) \times_S \bar{\eta}$ . On the other hand, let us recall that we have assumed that the base-change  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$  of  $Y \rightarrow X$  by  $\bar{\eta} \rightarrow S$  has a *splitting*. Thus, by considering the connected component of  $Y \times_S \bar{\eta}$  obtained by forming the image of a splitting of  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$ , we conclude that the connected finite étale covering  $Y \rightarrow X \times_S T$  [cf. Claim 1.3.D] is *of degree one*, i.e., an *isomorphism*, as desired. This completes the proof of Claim 1.3.E, hence also of the implication (2)  $\Rightarrow$  (3) [cf. also Claim 1.3.A].

Next, we verify the implication

$$(3) \implies (2).$$

To this end, suppose that condition (3) is satisfied, take a connected finite étale covering  $Y \rightarrow X$  of  $X$  whose base-change  $Y \times_S \bar{\eta} \rightarrow X \times_S \bar{\eta}$  by  $\bar{\eta} \rightarrow S$  has a *splitting*, and write  $T \rightarrow S$  for the normalization of  $S$  in  $Y$ . Then it follows from condition (3) that there exist a connected finite étale covering  $S_Y \rightarrow S$  and an isomorphism  $Y \xrightarrow{\sim} X \times_S S_Y$  over  $X$ . Now let us observe that it follows from [2], Proposition 4.6.1, together with the condition (b), that the function field of  $X$ , hence also of  $Y$ , is *separable* over the function field of  $S$ . Thus, since [one verifies easily that]  $S_Y$  is *normal*, and [we have assumed that] the

morphism  $X \rightarrow S$ , hence also  $Y \rightarrow S_Y$ , is *generically geometrically irreducible*, it follows from [2], Proposition 4.5.9, that the morphism  $S_Y \rightarrow S$  *factors* through a *birational* morphism  $S_Y \rightarrow T$  over  $S$ . In particular, since  $T$  is *normal*, and the morphism  $S_Y \rightarrow S$ , hence also  $S_Y \rightarrow T$ , is *finite*, it follows from [1], Corollaire 4.4.9, that the *birational finite* morphism  $S_Y \rightarrow T$  is an *isomorphism*, as desired. This completes the proof of the implication (3)  $\Rightarrow$  (2).

Finally, we verify the equivalence

$$(3) \iff (4).$$

To this end, suppose that the morphism  $f$  is *generically geometrically irreducible*. Now let us observe that since the composite  $X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$  *factors* through the geometric point  $\bar{\eta} \rightarrow S$ , the image of the composite  $\Pi_{X \times_S \bar{\eta}} \rightarrow \Pi_X \rightarrow \Pi_S$  is *trivial*. Moreover, we have assumed that the outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is *surjective* [cf. the condition (a)]. Thus, we conclude that condition (4) is *equivalent* to the inclusion

$$\Delta_f \subseteq \text{Im}(\Pi_{X \times_S \bar{\eta}} \rightarrow \Pi_X).$$

On the other hand, it is immediate that this inclusion is *equivalent* to condition (3). This completes the proof of the equivalence (3)  $\Leftrightarrow$  (4), hence also of Lemma 1.3.  $\square$

**REMARK 1.3.1.** — In the situation of Lemma 1.3, suppose that the function field of  $S$  is *of characteristic zero*. Then since [we have assumed that]  $X$  is *normal*, it follows from [2], Proposition 4.6.1, that the condition (b) is always satisfied.

**DEFINITION 1.4.** — Let  $\mathbb{P}$  be a property of a profinite group [e.g., “topologically finitely generated”, “abelian”, or “pro- $\Sigma$  for some set  $\Sigma$  of prime numbers”].

(i) We shall say that the property  $\mathbb{P}$  is *stable under taking open subquotients* if the following condition is satisfied: If a profinite group  $G$  has the property  $\mathbb{P}$ , then every quotient of an arbitrary open subgroup of  $G$  [i.e., by a normal closed subgroup of the open subgroup] has the property  $\mathbb{P}$ .

(ii) Let  $X$  be a scheme of CLN-type. Then we shall say that  $X$  *satisfies the condition*  $(*\mathbb{P})$  if the following condition is satisfied: For a scheme  $Y$  of CLN-type and a finite and generically étale surjective morphism  $f: Y \rightarrow X$ , if the profinite group  $\Delta_f$  has the property  $\mathbb{P}$ , then  $f$  is étale.

**LEMMA 1.5.** — *Let  $X, S$  be schemes of CLN-type;  $f: X \rightarrow S$  a morphism of schemes which is of finite type, separated, and dominant;  $\bar{\eta} \rightarrow S$  a geometric generic point of  $S$ . Suppose that the following two conditions are satisfied:*

- (a) *The outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is **surjective**.*
- (b) *The geometric generic fiber  $X \times_S \bar{\eta}$  of  $f$  is **reduced** [cf. Remark 1.3.1].*

*Suppose, moreover, that there exists a property  $\mathbb{P}$  of a profinite group such that*

- (1) *the property  $\mathbb{P}$  is **stable under taking open subquotients**,*



- (2) the kernel  $\Delta_f$  **has** the property  $\mathbb{P}$ , and, moreover,  
 (3) the scheme  $S$  of CLN-type **satisfies** the condition  $(*\mathbb{P})$ .

Then the following hold:

- (i) Let  $Y \rightarrow X$  be a connected finite étale covering of  $X$ . Then the normalization of  $S$  in  $Y$  is **étale** over  $S$ .  
 (ii) The morphism  $f$  is **generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

induces an **exact** sequence of profinite groups

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

PROOF. — First, we verify assertion (i). Write  $T \rightarrow S$  for the normalization of  $S$  in  $Y$ . First, let us verify the following claim:

Claim 1.5.A: The morphism  $T \rightarrow S$  is *finite, generically étale, and surjective*.

To this end, let us observe that it follows from [2], Proposition 4.6.1, together with the condition (b), that the function field of  $X$ , hence also of  $Y$ , is *separable* over the function field of  $S$ . Thus, the morphism  $T \rightarrow S$  is *finite* [cf. [9], §33, Lemma 1] and *generically étale*, which thus implies that  $f$  is *closed* and *dominant*. In particular, the morphism  $T \rightarrow S$  is *surjective*, as desired. This completes the proof of Claim 1.5.A.

Next, let us observe that we have a commutative diagram of profinite groups

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_{Y/S} & \longrightarrow & \Pi_Y & \longrightarrow & \Pi_S \\ & & \downarrow & & \downarrow & & \parallel \\ 1 & \longrightarrow & \Delta_{T/S} & \longrightarrow & \Pi_T & \longrightarrow & \Pi_S \end{array}$$

— where the horizontal sequences are *exact*. It follows immediately from [similar arguments to the arguments applied in the proofs of] [8], Lemma 1.6, and [8], Lemma 1.9, that the middle vertical arrow  $\Pi_Y \rightarrow \Pi_T$  of this diagram, hence also the left-hand vertical arrow  $\Delta_{Y/S} \rightarrow \Delta_{T/S}$  of this diagram, is *surjective*. In particular, since [it follows from conditions (1), (2) that] the profinite group  $\Delta_{Y/S}$  has the property  $\mathbb{P}$ , it follows from condition (1) that the profinite group  $\Delta_{T/S}$  has the property  $\mathbb{P}$ . Thus, it follows from Claim 1.5.A and condition (3) that the morphism  $T \rightarrow S$  is *étale*, as desired. This completes the proof of assertion (i).

Assertion (ii) follows from the implication (1)  $\Rightarrow$  (4) of Lemma 1.3, together with assertion (i). This completes the proof of Lemma 1.5.  $\square$

## 2. HOMOTOPY SEQUENCES FOR VARIETIES OVER CURVES

In the present §2, we obtain an application of the results of §1 concerning the homotopy sequences for normal varieties over smooth curves [cf. Theorem 2.8 below].

**LEMMA 2.1.** — *Let  $k$  be an algebraically closed field,  $X$  a hyperbolic curve over  $k$  [cf., e.g., [8], Definition 2.1, (i)], and  $l$  a prime number invertible in  $k$ . Then an arbitrary nontrivial normal closed subgroup of the maximal pro- $l$  quotient of  $\Pi_X$  of infinite index is free pro- $l$  and not topologically finitely generated.*

PROOF. — Let  $N$  be a nontrivial normal closed subgroup of the maximal pro- $l$  quotient of  $\Pi_X$  of infinite index. Let us first recall from [11], Proposition 2.3, (i), that  $N$  is not topologically finitely generated. Next, let us observe that since  $N$  is of infinite index, and the maximal pro- $l$  quotient of  $\Pi_X$  is topologically finitely generated [cf. [11], Proposition 2.2], there exists a sequence of open subgroups of the maximal pro- $l$  quotient of  $\Pi_X$

$$\cdots \subseteq H_{n+1} \subseteq H_n \subseteq \cdots \subseteq H_2 \subseteq H_1$$

such that  $H_i \neq H_{i+1}$  for an arbitrary positive integer  $i$ , and, moreover, the equality  $\bigcap_i H_i = N$  holds, which thus implies [cf. [12], Proposition 6.5.5] that the natural homomorphism

$$\varinjlim_i H^2(H_i, \mathbb{Z}/l\mathbb{Z}) \longrightarrow H^2(N, \mathbb{Z}/l\mathbb{Z})$$

— where the module  $\mathbb{Z}/l\mathbb{Z}$  is regarded as being equipped with the trivial actions of  $N$  and the  $H_i$ 's — is an isomorphism. In particular, to verify that  $N$  is free pro- $l$ , it follows from [12], Theorem 7.7.4, that it suffices to verify that the image of the natural homomorphism

$$H^2(H_n, \mathbb{Z}/l\mathbb{Z}) \longrightarrow H^2(H_{n+1}, \mathbb{Z}/l\mathbb{Z})$$

is zero for an arbitrary positive integer  $n$ . To this end, write  $X_{n+1} \rightarrow X_n \rightarrow X$  for the connected finite étale coverings of  $X$  that correspond to the open subgroups  $H_{n+1} \subseteq H_n$  of the maximal pro- $l$  quotient of  $\Pi_X$ , respectively. Now let us recall that the homomorphism under consideration fits into a commutative diagram

$$\begin{array}{ccc} H^2(H_n, \mathbb{Z}/l\mathbb{Z}) & \longrightarrow & H^2(H_{n+1}, \mathbb{Z}/l\mathbb{Z}) \\ \wr \downarrow & & \downarrow \wr \\ H_{\text{ét}}^2(X_n, \mathbb{Z}/l\mathbb{Z}) & \longrightarrow & H_{\text{ét}}^2(X_{n+1}, \mathbb{Z}/l\mathbb{Z}) \end{array}$$

— where the vertical arrows are isomorphisms [cf., e.g., [6], Lemma 4.2, (iii)], and the lower horizontal arrow is the homomorphism induced by the connected finite étale covering  $X_{n+1} \rightarrow X_n$ . Thus, since the module  $H^2(H_n, \mathbb{Z}/l\mathbb{Z})$  is annihilated by  $l$ , and [it is well-known that] the lower horizontal arrow of this diagram factors through the endomorphism of  $H_{\text{ét}}^2(X_n, \mathbb{Z}/l\mathbb{Z})$  given by multiplication by the degree of the covering  $X_{n+1} \rightarrow X_n$  — i.e., the index  $[H_n : H_{n+1}]$  ( $\neq 1$ ) — we conclude that the image of the upper horizontal arrow of the above diagram is zero, as desired. This completes the proof of Lemma 2.1.  $\square$

**DEFINITION 2.2.** — Let  $k$  be a field and  $X$  a scheme over  $k$ .

(i) We shall say that  $X$  is a normal variety over  $k$  if  $X$  is of finite type, separated, geometrically connected, and geometrically normal over  $k$ .

(ii) We shall say that  $X$  is a smooth curve over  $k$  if there exist a normal variety  $X^{\text{cpt}}$  over  $k$  which is proper over  $k$  and of dimension one, a [possible empty] closed subscheme  $D \subseteq X^{\text{cpt}}$  of  $X^{\text{cpt}}$  which is étale over  $k$ , and an isomorphism  $X \xrightarrow{\sim} X^{\text{cpt}} \setminus D$  over  $k$ .

**REMARK 2.2.1.** — One verifies easily that a *smooth curve* over a field is *smooth* and a *normal variety* over the field.

**LEMMA 2.3.** — *Let  $k$  be an algebraically closed field,  $X$  a smooth curve over  $k$ , and  $l$  a prime number invertible in  $k$ . Suppose that  $X$  is isomorphic neither to the projective line over  $k$  nor to the affine line over  $k$ . Then the maximal pro- $l$  quotient of  $\Pi_X$ , hence also  $\Pi_X$  itself, is infinite.*

PROOF. — This assertion follows from, for instance, the isomorphism (1-3) and the exact sequence (1-5) of [14], Remark 1.3.  $\square$

**LEMMA 2.4.** — *Let  $k$  be an algebraically closed field,  $X$  and  $Y$  smooth curves over  $k$ ,  $l$  a prime number invertible in  $k$ , and  $f: X \rightarrow Y$  a dominant and generically étale morphism over  $k$ . Suppose that the following two conditions are satisfied:*

(1) *The smooth curve  $Y$  is isomorphic neither to the projective line over  $k$  nor to the affine line over  $k$ .*

(2) *The outer homomorphism from the maximal pro- $l$  quotient of  $\Pi_X$  to the maximal pro- $l$  quotient of  $\Pi_Y$  induced by  $f$  is an outer isomorphism.*

*Then  $f$  is a finite étale covering.*

PROOF. — Let us write  $X^{\text{cpt}}$  (respectively,  $Y^{\text{cpt}}$ ) for the smooth compactification of  $X$  (respectively,  $Y$ ),  $g_X$  (respectively,  $g_Y$ ) for the genus of  $X^{\text{cpt}}$  (respectively,  $Y^{\text{cpt}}$ ), and  $r_X$  (respectively,  $r_Y$ ) for the cardinality of the complement in  $X^{\text{cpt}}$  (respectively,  $Y^{\text{cpt}}$ ) of  $X$  (respectively,  $Y$ ). Then since  $f$  is *dominant*, it follows from elementary theory of algebraic curves that

$$g_X \geq g_Y \text{ and } r_X \geq r_Y.$$

Thus, it follows immediately from conditions (1), (2), together with [14], Proposition 1.1 [cf. also the isomorphism (1-3) and the exact sequence (1-5) of [14], Remark 1.3], that

$$g_X = g_Y \text{ and } r_X = r_Y.$$

In particular, since  $f$  is *generically étale*, it follows immediately from condition (1), together with elementary theory of algebraic curves, that  $f$  is a *finite étale covering*, as desired. This completes the proof of Lemma 2.4.  $\square$

**DEFINITION 2.5.**

(i) Let  $\Sigma$  be a [possibly empty] set of prime numbers. Then we shall write

$$\mathbb{P}_{\mathcal{Z} \rightarrow \infty}^{\exists \notin \Sigma}$$

for the property of a profinite group defined as follows: A profinite group  $G$  has the property  $\mathbb{P}_{\mathcal{Z} \rightarrow \infty}^{\exists \notin \Sigma}$  if, for an arbitrary open subgroup  $H \subseteq G$  of  $G$ , there exists a prime

number  $l_H$  such that  $l_H \notin \Sigma$ , and, moreover, there is no quotient of  $H$  which is free pro- $l_H$  and not topologically finitely generated. Moreover, we shall write

$$\mathbb{P}_{\mathcal{F} \rightarrow \infty} \stackrel{\text{def}}{=} \mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \emptyset}.$$

(ii) Let  $\mathbb{P}$  be a property of a profinite group. Then we shall say that a profinite group  $G$  has the property *almost*  $\mathbb{P}$  if an open subgroup of  $G$  has the property  $\mathbb{P}$ .

**REMARK 2.5.1.** — One verifies easily that, in the situation of Definition 2.5, (i), if  $\Sigma$  is *not equal* to the set of all prime numbers, and a profinite group  $G$  has one of the following three properties, then  $G$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ :

- (1) The profinite group  $G$  is *topologically finitely generated*.
- (2) The profinite group  $G$  is *almost abelian*.
- (3) The profinite group  $G$  is *almost pro- $\Sigma'$*  for some set  $\Sigma'$  of prime numbers such that the union  $\Sigma \cup \Sigma'$  does *not coincide* with the set of all prime numbers.

**LEMMA 2.6.** — *Let  $\Sigma$  be a set of prime numbers and  $G$  a profinite group. Then the following hold:*

- (i) *Let  $H \subseteq G$  be an open subgroup of  $G$ . Then the profinite group  $G$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$  if and only if the profinite group  $H$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ .*
- (ii) *The property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$  is **stable under taking open subquotients**.*

PROOF. — First, we verify assertion (i). It is immediate that if  $G$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ , then  $H$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ . Suppose that  $H$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ . To complete the verification of assertion (i), assume that  $G$  does *not have* the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ , i.e., that there exists an open subgroup  $U \subseteq G$  of  $G$  such that, for an arbitrary prime number  $l \notin \Sigma$ , the profinite group  $U$  has a *free pro- $l$  and not topologically finitely generated* quotient  $U \twoheadrightarrow Q$ . Then since  $U \cap H$  is an *open* subgroup of  $U$ , the image of the composite  $U \cap H \hookrightarrow U \twoheadrightarrow Q$  is an *open* subgroup of  $Q$ . In particular, this image is *free pro- $l$*  [cf. [12], Corollary 7.7.5] and *not topologically finitely generated*. Thus, since  $U \cap H$  is an *open* subgroup of  $H$ , we obtain a *contradiction*. This completes the proof of assertion (i).

Next, we verify assertion (ii). It follows from assertion (i) that, to verify assertion (ii), it suffices to verify that if a profinite group has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ , then an arbitrary quotient of the profinite group has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}^{\exists \notin \Sigma}$ . On the other hand, this is immediate. This completes the proof of assertion (ii), hence also of Lemma 2.6.  $\square$

**LEMMA 2.7.** — *Let  $k$  be an algebraically closed field and  $X$  a **smooth curve** over  $k$ . Suppose that  $X$  is **isomorphic neither** to the projective line over  $k$  **nor** to the affine line over  $k$ . Write  $\Sigma$  for the set of prime numbers **not invertible** in  $k$ . [So  $\#\Sigma \leq 1$ .] Then the following hold:*

(i) Let  $Y$  be a **hyperbolic curve** over  $k$  and  $f: Y \rightarrow X$  a **dominant and generically étale** morphism over  $k$ . Suppose that there exists a prime number  $l \notin \Sigma$  such that there is **no** quotient of  $\Delta_f$  which is **free pro- $l$**  and **not topologically finitely generated**. Then  $f$  is a **finite étale covering**.

(ii) The smooth curve  $X$  **satisfies** the condition  $(*_\frac{\mathbb{P}^1 \setminus \Sigma}{l \rightarrow \infty})$  [cf. Definition 1.4, (ii)].

PROOF. — First, we verify assertion (i). Let us first observe that since [we have assumed that]  $X$  is *isomorphic neither* to the projective line over  $k$  nor to the affine line over  $k$ , it follows from elementary theory of algebraic curves that an arbitrary connected finite étale covering of  $X$  is *isomorphic neither* to the projective line over  $k$  nor to the affine line over  $k$ . Thus, we may assume without loss of generality, by replacing  $X$  by the connected finite étale covering of  $X$  that corresponds to the image of some homomorphism  $\Pi_Y \rightarrow \Pi_X$  induced by  $f$ , that

the outer homomorphism  $\Pi_Y \rightarrow \Pi_X$  is *surjective*.

Let  $l \notin \Sigma$  be as in the statement of assertion (i). Then we have an *exact* sequence of profinite groups

$$\Delta_f^{(l)} \longrightarrow \Pi_Y^{(l)} \longrightarrow \Pi_X^{(l)} \longrightarrow 1$$

— where we write “ $(-)^{(l)}$ ” for the maximal pro- $l$  quotient of “ $(-)$ ”. Now let us recall from Lemma 2.3 that the profinite group  $\Pi_X^{(l)}$  is *infinite*, which thus implies that the image of the first arrow of the above exact sequence is of *infinite index* in  $\Pi_Y^{(l)}$ . Thus, it follows immediately from Lemma 2.1, together with our assumption on the quotients of  $\Delta_f$ , that the image of the first arrow of the above exact sequence is *trivial*. In particular, the outer homomorphism  $\Pi_Y^{(l)} \rightarrow \Pi_X^{(l)}$  induced by  $f$  is an outer *isomorphism*. Thus, it follows from Lemma 2.4 that  $f$  is a *finite étale covering*, as desired. This completes the proof of assertion (i).

Next, we verify assertion (ii). Let us first observe that it follows immediately from assertion (i) that, to verify assertion (ii), it suffices to verify the following claim:

Claim 2.7.A: Let  $Y$  be a scheme of *CLN-type* and  $f: Y \rightarrow X$  a *finite and generically étale surjective* morphism. [So one verifies easily that  $Y$  has a natural structure of a smooth curve over  $k$  with respect to which  $f$  is a morphism over  $k$ .] If  $Y$ , hence also  $X$ , is *not a hyperbolic curve* over  $k$ , then  $f$  is *étale*.

To this end, let us first observe that since [we have assumed that]  $X$  is *isomorphic neither* to the projective line over  $k$  nor to the affine line over  $k$ , it follows immediately from elementary theory of algebraic curves that both  $Y$  and  $X$  are either

- *proper* over  $k$  and of *genus one*, or
- *isomorphic* to the complement in the projective line over  $k$  of *distinct two* closed points.

Thus, since [we have assumed that]  $f$  is *generically étale* and *finite*, again by elementary theory of algebraic curves, we conclude immediately that  $f$  is *étale*, as desired. This completes the proof of assertion (ii), hence also of Lemma 2.7. □

**THEOREM 2.8.** — *Let  $k$  be a field,  $S$  a smooth curve over  $k$  [cf. Definition 2.2, (ii)] whose base-change to an algebraic closure of  $k$  is **isomorphic neither** to the projective line **nor** to the affine line,  $X$  a **normal variety** over  $k$  [cf. Definition 2.2, (i)],  $f: X \rightarrow S$  a morphism over  $k$ , and  $\bar{\eta} \rightarrow S$  a geometric generic point of  $S$ . Suppose that the geometric generic fiber  $X \times_S \bar{\eta}$  of  $f$  is [either empty or] **reduced** [cf. Remark 2.8.1 below]. Consider the following four conditions:*

(1) *The outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is **surjective**. Moreover, the profinite group  $\Delta_f$  is **topologically finitely generated**.*

(2) *The outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is **surjective**. Moreover, if we write  $\Sigma$  for the set of prime numbers **not invertible** in  $k$  [so  $\#\Sigma \leq 1$ ], then the profinite group  $\Delta_f$  **has** the property  $\mathbb{P}_{\mathbb{F} \rightarrow \infty}^{\exists \notin \Sigma}$  defined in Definition 2.5, (i) [which is the case if, for instance, either*

- *the profinite group  $\Delta_f$  is **topologically finitely generated**,*

- *the profinite group  $\Delta_f$  is **almost abelian**, or*

- *the profinite group  $\Delta_f$  is **almost pro- $\Sigma'$**  for some [possibly empty] set  $\Sigma'$  of prime numbers such that there exists a prime number **not contained in  $\Sigma'$  and invertible in  $k$***

— *cf. Remark 2.5.1].*

(3) *The morphism  $f$  is **surjective and generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes*

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

*induces an **exact** sequence of profinite groups*

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

(4) *The morphism  $f$  is **dominant and generically geometrically irreducible**. Moreover, the sequence of irreducible [hence also connected] schemes*

$$X \times_S \bar{\eta} \xrightarrow{\text{pr}_1} X \xrightarrow{f} S$$

*induces an **exact** sequence of profinite groups*

$$\Pi_{X \times_S \bar{\eta}} \longrightarrow \Pi_X \longrightarrow \Pi_S \longrightarrow 1.$$

*Then the following hold:*

(i) *The implications*

$$(1) \implies (2) \implies (3) \implies (4)$$

*hold.*

(ii) *Suppose that either  $k$  is **of characteristic zero**, or the geometric generic fiber  $X \times_S \bar{\eta}$  of  $f$  is **proper** over  $\bar{\eta}$ . Then the equivalences*

$$(1) \iff (2) \iff (3) \iff (4)$$

*hold.*

PROOF. — Let us first observe that one verifies immediately from [15], Exposé IX, Théorème 6.1, that, to verify Theorem 2.8, we may assume without loss of generality, by replacing  $k$  by the algebraic closure of  $k$  determined by the composite  $\bar{\eta} \rightarrow S \rightarrow \text{Spec}(k)$ , that

$k$  is algebraically closed.

Now we verify assertion (i). The implications

$$(1) \implies (2), \quad (3) \implies (4)$$

are immediate. Next, to verify the implication

$$(2) \implies (4),$$

suppose that condition (2) is satisfied. Let us first recall from Lemma 2.3 that the profinite group  $\Pi_S$  is *nontrivial*. In particular, since [we have assumed that] the outer homomorphism  $\Pi_X \rightarrow \Pi_S$  is *surjective*, we conclude that the morphism  $f$  is *nonconstant*, i.e., *dominant*. Thus, the implication (2)  $\Rightarrow$  (4) follows from Lemma 1.5, (ii), and Lemma 2.7, (ii) [cf. also Lemma 2.6, (ii)]. This completes the proof of the implication (2)  $\Rightarrow$  (4).

Next, to verify the implication

$$(2) \implies (3),$$

suppose that condition (2), hence also condition (4) [cf. the preceding paragraph], is satisfied, and write  $U \subseteq S$  for the image of the morphism  $f: X \rightarrow S$ . [Thus,  $U$  is an *open* subscheme of  $S$  — cf. condition (4).] Then we have a commutative diagram of profinite groups

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_f & \longrightarrow & \Pi_X & \longrightarrow & \Pi_S \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \parallel \\ 1 & \longrightarrow & \Delta_{U/S} & \longrightarrow & \Pi_U & \longrightarrow & \Pi_S \longrightarrow 1 \end{array}$$

— where the horizontal sequences are *exact*. Next, let us observe that since the resulting [necessarily *surjective*] morphism  $X \rightarrow U$  is *generically geometrically irreducible* [cf. condition (4)], it follows from [a similar argument to the argument applied in the proof of] [8], Lemma 1.6, that the middle vertical arrow  $\Pi_X \rightarrow \Pi_U$  of this diagram, hence also the left-hand vertical arrow  $\Delta_f \rightarrow \Delta_{U/S}$  of this diagram, is *surjective*. In particular, since the profinite group  $\Delta_f$  has the property  $\mathbb{P}_{\bar{\mathcal{F}} \rightarrow \infty}^{\exists \notin \Sigma}$  [cf. condition (2)], it follows from Lemma 2.6, (ii), that the profinite group  $\Delta_{U/S}$  has the property  $\mathbb{P}_{\bar{\mathcal{F}} \rightarrow \infty}^{\exists \notin \Sigma}$ , which thus implies that there exists a prime number  $l \notin \Sigma$  such that there is *no* quotient of  $\Delta_{U/S}$  which is *free pro- $l$*  and *not topologically finitely generated*.

Next, let us observe that the lower horizontal sequence of the above diagram determines an *exact* sequence of profinite groups

$$\Delta_{U/S}^{(l)} \longrightarrow \Pi_U^{(l)} \longrightarrow \Pi_S^{(l)} \longrightarrow 1$$

— where we write “ $(-)^{(l)}$ ” for the maximal pro- $l$  quotient of “ $(-)$ ”. Now assume that  $U \neq S$ . Then since [we have assumed that]  $S$  is *isomorphic neither* to the projective line over  $k$  *nor* to the affine line over  $k$ , one verifies easily that  $U$  is a *hyperbolic curve*

over  $k$ . Moreover, it follows from Lemma 2.3 that the profinite group  $\Pi_S^{(l)}$  is *infinite*, which thus implies that the image of the first arrow of the above exact sequence is of *infinite index* in  $\Pi_U^{(l)}$ . Thus, it follows immediately from Lemma 2.1, together with our assumption on the quotients of  $\Delta_{U/S}$ , that the image of the first arrow of the above exact sequence is *trivial*. In particular, the outer homomorphism  $\Pi_U^{(l)} \rightarrow \Pi_S^{(l)}$  induced by the open immersion  $U \hookrightarrow S$  is an outer *isomorphism*. Thus, it follows from Lemma 2.4 that  $U = S$ , in *contradiction* to our assumption that  $U \neq S$ . This completes the proof of the implication (2)  $\Rightarrow$  (3), hence also of assertion (i).

Next, we verify assertion (ii). Suppose that condition (4) is satisfied. Thus, the outer homomorphism  $\Pi_X \rightarrow \Pi_S$  induced by  $f$  is *surjective*. Next, let us observe that it follows from condition (4) that the outer homomorphism  $\Pi_{X \times_S \bar{\eta}} \rightarrow \Pi_X$  induced by  $X \times_S \bar{\eta} \rightarrow X$  determines a  $\Pi_X$ -conjugacy class of *surjective* homomorphisms  $\Pi_{X \times_S \bar{\eta}} \rightarrow \Delta_f$ . Thus, since  $\Pi_{X \times_S \bar{\eta}}$  is *topologically finitely generated* [cf. [8], Lemma 1.7, and [15], Exposé X, Théorème 2.9], we conclude that condition (1) is satisfied, as desired. This completes the proof of assertion (ii), hence also of Theorem 2.8.  $\square$

**REMARK 2.8.1.** — In the situation of Theorem 2.8, suppose that  $k$  is of *characteristic zero*. Then it follows from Remark 1.3.1 that the assumption that the geometric generic fiber  $X \times_S \bar{\eta}$  of  $f$  is [either empty or] *reduced* in the statement of Theorem 2.8 is always satisfied.

**REMARK 2.8.2.** — Theorem 2.8 may be regarded as a *refinement* of the implication (2)  $\Rightarrow$  (3) of [8], Lemma 2.11.

**REMARK 2.8.3.** — Let  $k$  be an algebraically closed field of characteristic zero. Write  $S$  for *either the projective line over  $k$  or the affine line over  $k$* . Let  $f: X \stackrel{\text{def}}{=} \text{Spec}(k) \rightarrow S$  be a  $k$ -rational point of  $S$  [i.e., a splitting of the structure morphism of  $S$ ]. Then one verifies easily that the morphism  $f$  *satisfies* condition (1) of Theorem 2.8 but does *not satisfy* condition (4) of Theorem 2.8.

### 3. AN ANABELIAN RESULT ON HYPERBOLIC POLYCURVES OF DIMENSION TWO

In the present §3, we obtain an application of the results of §1 concerning the Grothendieck conjecture for hyperbolic polycurves of dimension two [cf. Theorem 3.3 below].

**DEFINITION 3.1.** — Let  $\mathbb{P}$  be a property of a profinite group. Then we shall say that the property  $\mathbb{P}$  is *stable under taking extensions of topologically finitely generated profinite groups* if the following condition is satisfied: If a profinite group  $G$  has the property  $\mathbb{P}$ , then an arbitrary extension of a topologically finitely generated profinite group by  $G$  [i.e, a profinite group  $E$  that fits into an exact sequence of profinite groups

$$1 \longrightarrow G \longrightarrow E \longrightarrow H \longrightarrow 1$$

— where  $H$  is topologically finitely generated] has the property  $\mathbb{P}$ .



**REMARK 3.1.1.** — Let  $\mathbb{P}$  be a property of a profinite group which is *stable under taking open subquotients* and *stable under taking extensions of topologically finitely generated profinite groups*. Suppose that there exists a profinite group that has the property  $\mathbb{P}$ . Then one verifies easily that an *arbitrary topologically finitely generated* profinite group has the property  $\mathbb{P}$ .

**LEMMA 3.2.** — *Let  $\Sigma$  be a set of prime numbers. Then the property  $\mathbb{P}_{\neq\infty}^{\exists\neq\Sigma}$  on a profinite group defined in Definition 2.5, (i), is **stable under taking extensions of topologically finitely generated profinite groups**.*

PROOF. — Let  $G$  be a profinite group that has the property  $\mathbb{P}_{\neq\infty}^{\exists\neq\Sigma}$  and

$$1 \longrightarrow G \longrightarrow E \longrightarrow H \longrightarrow 1$$

an exact sequence of profinite groups such that  $H$  is *topologically finitely generated*. Now, to verify Lemma 3.2, assume that the profinite group  $E$  does *not* have the property  $\mathbb{P}_{\neq\infty}^{\exists\neq\Sigma}$ , i.e., that there exists an open subgroup  $U \subseteq E$  of  $E$  such that, for an arbitrary prime number  $l \notin \Sigma$ , the profinite group  $U$  has a *free pro- $l$*  and *not topologically finitely generated* quotient  $U \twoheadrightarrow Q_l$ .

Now let us observe that since  $G$  is *normal* in  $E$ , the image of the composite  $U \cap G \hookrightarrow U \twoheadrightarrow Q_l$  is *normal* in  $Q_l$ . In particular, if this image of  $U \cap G$  in  $Q_l$  is *nontrivial*, then this image in  $Q_l$  is *free pro- $l$*  [cf. [12], Corollary 7.7.5] and *not topologically finitely generated* [cf. [12], Theorem 8.6.5]. Thus, since  $U \cap G$  is an *open* subgroup of  $G$ , and [we have assumed that]  $G$  has the property  $\mathbb{P}_{\neq\infty}^{\exists\neq\Sigma}$ , we conclude that there exists a prime number  $l \notin \Sigma$  such that the image of the composite  $U \cap G \hookrightarrow U \twoheadrightarrow Q_l$  is *trivial*, which thus implies that we have a *surjective* homomorphism  $U/(U \cap G) \twoheadrightarrow Q_l$ .

On the other hand, since  $U/(U \cap G)$  may be regarded as an *open* subgroup of the *topologically finitely generated* profinite group  $H$ , the quotient  $U/(U \cap G)$  is *topologically finitely generated*. Thus, since  $Q_l$  is *not topologically finitely generated*, we obtain a *contradiction* by the above *surjective* homomorphism  $U/(U \cap G) \twoheadrightarrow Q_l$ . This completes the proof of Lemma 3.2.  $\square$

**THEOREM 3.3.** — *Let  $p$  be a prime number,  $k$  a sub- $p$ -adic field [cf., e.g., [8], Definition 3.1],  $\bar{k}$  an algebraic closure of  $k$ ,  $X$  a **hyperbolic polycurve of dimension 2** over  $k$  [cf. [8], Definition 2.1, (ii)], and  $Z$  a **normal variety** over  $k$  [cf. Definition 2.2, (i)]. Write  $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ . Let*

$$\alpha: \Pi_Z \longrightarrow \Pi_X$$

*be an **open** homomorphism of profinite groups over  $G_k$ . Suppose that the kernel  $\text{Ker}(\alpha)$  of  $\alpha$  has the property  $\mathbb{P}_{\neq\infty}$  defined in Definition 2.5, (i) [which is the case if, for instance, either*

- *the profinite group  $\text{Ker}(\alpha)$  is **topologically finitely generated**,*
- *the profinite group  $\text{Ker}(\alpha)$  is **almost abelian**, or*
- *the profinite group  $\text{Ker}(\alpha)$  is **almost pro- $\Sigma$**  for some [possibly empty] set  $\Sigma$  of prime numbers **not equal** to the set of all prime numbers*

— cf. Remark 2.5.1]. Then there exists a **uniquely determined dominant morphism**

$$Z \longrightarrow X$$

over  $k$  from which  $\alpha$  arises. In particular, the normal variety  $Z$  is of dimension  $\geq 2$ .

PROOF. — Let us first observe that one verifies easily [cf. also [8], Proposition 2.3] that, to verify Theorem 3.3, we may assume without loss of generality, by replacing  $X$  by the connected finite étale covering of  $X$  that corresponds to the image of  $\alpha$ , that

the homomorphism  $\alpha$  is *surjective*.

Fix a parametrizing morphism  $X \rightarrow S$  for  $X$  [cf. [8], Definition 2.1, (ii)] and a [necessarily *surjective* — cf. [8], Proposition 2.4, (i)] homomorphism  $\Pi_X \rightarrow \Pi_S$  over  $G_k$  induced by this parametrizing morphism  $X \rightarrow S$ . Write  $\beta: \Pi_Z \xrightarrow{\alpha} \Pi_X \rightarrow \Pi_S$  for the composite of  $\alpha$  and the fixed surjective homomorphism  $\Pi_X \rightarrow \Pi_S$ . Then it follows from [8], Theorem A, in the case where condition (1) is satisfied that there exists a *uniquely determined dominant morphism*

$$f: Z \longrightarrow S$$

over  $k$  from which  $\beta$  arises.

Now we verify the following claim:

Claim 3.3.A: The kernel  $\Delta_f$  [i.e., the kernel of  $\beta$ ] has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ .

To this end, let us first observe that we have a commutative diagram of profinite groups

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_f & \longrightarrow & \Pi_Z & \xrightarrow{\beta} & \Pi_S \longrightarrow 1 \\ & & \downarrow & & \downarrow \alpha & & \parallel \\ 1 & \longrightarrow & \Delta_{X/S} & \longrightarrow & \Pi_X & \longrightarrow & \Pi_S \longrightarrow 1 \end{array}$$

— where the horizontal sequences are *exact*. Thus, the kernel  $\Delta_f$  is *isomorphic* to an extension of  $\Delta_{X/S}$  by the kernel of  $\alpha$ . Now let us recall from [8], Proposition 2.4, (iii), that the profinite group  $\Delta_{X/S}$  is *topologically finitely generated*. Thus, since the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$  is *stable under taking extensions of topologically finitely generated profinite groups* [cf. Lemma 3.2], and [we have assumed that] the kernel of  $\alpha$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ , the profinite group  $\Delta_f$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ , as desired. This completes the proof of Claim 3.3.A.

Next, we verify the following claim:

Claim 3.3.B: Both the morphism  $X \rightarrow S$  and  $f: Z \rightarrow S$  are *generically geometrically irreducible*.

To this end, let us recall that it follows from the definition of a hyperbolic curve that the morphism  $X \rightarrow S$  is *generically geometrically irreducible*. On the other hand, the assertion that the morphism  $f: Z \rightarrow S$  is *generically geometrically irreducible* follows from Lemma 1.5, (ii) [cf. also Lemma 2.6, (ii), and Lemma 2.7, (ii)], together with Claim 3.3.A. This completes the proof of Claim 3.3.B.

Write  $\eta \rightarrow S$  for the [morphism of schemes determined by the] generic point of  $S$ . Let  $\bar{\eta} \rightarrow (\eta \rightarrow S)$  be a geometric generic point of  $S$ . Next, we verify the following claim:

Claim 3.3.C: There exists a *surjective* homomorphism

$$\alpha_\eta: \Pi_{Z \times_S \eta} \longrightarrow \Pi_{X \times_S \eta}$$

[cf. Claim 3.3.B] over  $\Pi_\eta$  such that the diagram of profinite groups

$$\begin{array}{ccc} \Pi_{Z \times_S \eta} & \xrightarrow{\alpha_\eta} & \Pi_{X \times_S \eta} \\ \downarrow & & \downarrow \\ \Pi_Z & \xrightarrow{\alpha} & \Pi_X \end{array}$$

— where the vertical arrows are some [necessarily *surjective* — cf. [8], Lemma 1.2] homomorphisms over  $\Pi_S$  induced by  $Z \times_S \eta \xrightarrow{\text{pr}_1} Z$ ,  $X \times_S \eta \xrightarrow{\text{pr}_1} X$ , respectively — *commutes*.

To this end, let us first observe that it follows immediately from the various definitions involved that we have a sequence of profinite groups over  $\Pi_\eta$

$$\Pi_{Z \times_S \eta} \longrightarrow \Pi_Z \times_{\Pi_S} \Pi_\eta \xrightarrow{\alpha \times_{\text{id}_{\Pi_S}} \text{id}_{\Pi_\eta}} \Pi_X \times_{\Pi_S} \Pi_\eta \longleftarrow \Pi_{X \times_S \eta}.$$

On the other hand, it follows from [8], Proposition 2.4, (ii), that the third arrow of this sequence is an *isomorphism*; moreover, since  $\alpha$  is *surjective*, one verifies easily that the second arrow of this sequence is *surjective*. Thus, to complete the verification of Claim 3.3.C, it suffices to verify that the first arrow of this sequence is *surjective*, or, alternatively [cf. [8], Lemma 1.5], the  $\Pi_Z$ -conjugacy class of homomorphisms  $\Pi_{Z \times_S \eta} \rightarrow \Delta_{Z/S}$  induced by  $Z \times_S \eta \xrightarrow{\text{pr}_1} Z$  is *surjective*. On the other hand, this follows from Lemma 1.5, (ii) [cf. also Lemma 2.6, (ii), and Lemma 2.7, (ii)], together with Claim 3.3.A. This completes the proof of Claim 3.3.C.

Thus, since [one verifies easily that]  $Z \times_S \eta$  is a *normal variety* over  $\eta$  [i.e., over the function field of  $S$ ], and the function field of  $S$  is *sub- $p$ -adic*, it follows immediately from [8], Theorem A, in the case where condition (1) is satisfied and [8], Lemma 2.10, that there exists a *uniquely determined dominant* morphism

$$Z \longrightarrow X$$

over  $S$  from which  $\alpha$  arises, as desired. This completes the proof of Theorem 3.3.  $\square$

**REMARK 3.3.1.** — Let us recall that [8], Theorem A, in the case where condition (2) is satisfied is nothing but Theorem 3.3 under the assumption that the kernel of  $\alpha$  is *topologically finitely generated*. Thus, Theorem 3.3 may be regarded as a *refinement* of [8], Theorem A, in the case where condition (2) is satisfied.

**REMARK 3.3.2.** — Suppose that we are in the situation of Theorem 3.3. Then one may conclude that

if one drops the assumption that the hyperbolic polycurve  $X$  is of *dimension 2*, then the conclusion of Theorem 3.3 no longer holds in general

as follows: Let  $C_1, C_2, C_3$  be proper hyperbolic curves over  $k$ . Then  $Y \stackrel{\text{def}}{=} C_1 \times_k C_2 \times_k C_3$  is a *hyperbolic polycurve of dimension 3* over  $k$ . Now it follows from [5], Exposé XII, Corollaire 3.5, that there exist a *normal variety  $H$  of dimension 2* over  $k$  and a closed immersion  $i: H \hookrightarrow Y$  over  $k$  such that  $i$  induces an outer *isomorphism*

$$\Pi_H \xrightarrow{\sim} \Pi_Y.$$

Now let us observe that

(a) the triple consisting of the hyperbolic polycurve  $Y$  over  $k$ , the normal variety  $H$  over  $k$ , and some fixed isomorphism  $\Pi_H \xrightarrow{\sim} \Pi_Y$  over  $G_k$  contained in the outer isomorphism induced by  $i$  satisfies the condition imposed on the triple “ $(X, Z, \alpha)$ ” in the statement of Theorem 3.3 *except for the condition that “ $X$ ” is of dimension 2*. On the other hand, there is *no dominant* morphism from  $H$  to  $X$  [cf.  $\dim(H) = 2 < 3 = \dim(X)$ ], which thus implies that the fixed isomorphism  $\Pi_H \xrightarrow{\sim} \Pi_Y$  over  $G_k$  *never arises* from a *dominant* morphism over  $k$ .

Moreover, let  $C_4$  be a hyperbolic curve over  $k$ . Then one verifies easily [cf. [8], Proposition 2.4, (i)] that the composite  $H \times_k C_4 \rightarrow Y$  of the first projection  $H \times_k C_4 \rightarrow H$  and the closed immersion  $i: H \hookrightarrow Y$  induces an outer *surjective* homomorphism

$$\Pi_{H \times_k C_4} \twoheadrightarrow \Pi_Y$$

whose kernel is *isomorphic* to  $\Delta_{C_4/k}$ , hence also *topologically finitely generated* [cf. [8], Proposition 2.4, (iii)]. Now let us also observe that

(b) the triple consisting of the hyperbolic polycurve  $Y$  over  $k$ , the normal variety  $H \times_k C_4$  over  $k$ , and some fixed surjective homomorphism  $\Pi_{H \times_k C_4} \twoheadrightarrow \Pi_Y$  over  $G_k$  contained in the outer surjective homomorphism induced by the above composite satisfies the condition imposed on the triple “ $(X, Z, \alpha)$ ” in the statement of Theorem 3.3 *except for the condition that “ $X$ ” is of dimension 2*. On the other hand, it follows immediately from [8], Proposition 3.2, (ii), that the fixed surjective homomorphism  $\Pi_{H \times_k C_4} \twoheadrightarrow \Pi_Y$  over  $G_k$  *never arises* from a *dominant* morphism over  $k$ .

**REMARK 3.3.3.** — Let us observe that it follows from [8], Lemma 2.6, (i), and [8], Proposition 2.7, that the normal variety “ $H$ ” over  $k$  of Remark 3.3.2 is *of LFG-type* [cf. [8], Definition 2.5]. In particular, it follows from a similar argument to the argument applied in the proof of [8], Proposition 2.7, that the normal variety “ $H \times_k C_4$ ” over  $k$  of Remark 3.3.2 is *of LFG-type*. Thus, the observation (a) of Remark 3.3.2 implies that

if one drops assumption (3-iv) in the statement of [8], Theorem A, then the conclusion of [8], Theorem A [i.e., in the case where condition (3) is satisfied], no longer holds in general.

Moreover, the observation (b) of Remark 3.3.2 implies that

if one drops assumption (3-ii) in the statement of [8], Theorem A, then the conclusion of [8], Theorem A [i.e., in the case where condition (3) is satisfied], no longer holds in general.

Finally, one may also conclude from a similar argument to the argument applied in the consideration related to (b) of Remark 3.3.2 that

if one drops assumption (4-ii) in the statement of [8], Theorem A, then the conclusion of [8], Theorem A [i.e., in the case where condition (4) is satisfied], no longer holds in general.

**REMARK 3.3.4.**

- (i) Suppose that we are in the situation of Theorem 3.3. Then one may conclude that if one drops the assumption that the kernel of  $\alpha$  has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ , then the conclusion of Theorem 3.3 no longer holds in general

as follows: Let  $C_1$  and  $C_2$  be proper hyperbolic curves over  $k$ . Then  $W \stackrel{\text{def}}{=} C_1 \times_k C_2$  is a hyperbolic polycurve of dimension 2 over  $k$ . Now it follows from [5], Exposé XII, Corollaire 3.5, that there exist a normal variety  $J$  over  $k$  and a closed immersion  $j: J \hookrightarrow W$  over  $k$  such that  $j$  is not an isomorphism [or, alternatively, not dominant] — which thus implies that  $J$  is of dimension  $\leq 1$  — but induces an outer surjective homomorphism

$$\Pi_J \twoheadrightarrow \Pi_W.$$

Now let us observe that the triple consisting of the hyperbolic polycurve  $W$  over  $k$ , the normal variety  $J$  over  $k$ , and some fixed surjective homomorphism  $\Pi_J \rightarrow \Pi_W$  over  $G_k$  contained in the outer surjective homomorphism induced by  $j$  satisfies the condition imposed on the triple “ $(X, Z, \alpha)$ ” in the statement of Theorem 3.3 *except for the condition that the kernel of “ $\alpha$ ” has the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$* . On the other hand, there is no dominant morphism from  $J$  to  $W$  [cf.  $\dim(J) \leq 1 < 2 = \dim(W)$ ], which thus implies that the fixed surjective homomorphism  $\Pi_J \twoheadrightarrow \Pi_W$  over  $G_k$  never arises from a dominant morphism over  $k$ .

- (ii) One may find

an example of a triple consisting of a hyperbolic polycurve  $X$  of dimension 2 over  $k$ , a normal variety  $Z$  over  $k$ , and a dominant morphism  $f: Z \rightarrow X$  over  $k$  such that the kernel  $\Delta_f$  does not have the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ .

as follows: Let  $C_1, C_2, C_3$  be hyperbolic curves over  $k$ ;  $f_C: C_2 \rightarrow C_3$  a dominant morphism over  $k$  that is not a finite étale covering. Then both  $W_{1,2} \stackrel{\text{def}}{=} C_1 \times_k C_2$  and  $W_{1,3} \stackrel{\text{def}}{=} C_1 \times_k C_3$  are hyperbolic polycurves of dimension 2 over  $k$ , and the morphism  $f_W: W_{1,2} \rightarrow W_{1,3}$  over  $C_1$ , hence also over  $k$ , induced by  $f_C$  is dominant. Moreover, one verifies easily [cf. [8], Proposition 2.4, (i)] that  $\Delta_{f_W} \cong \Delta_{f_C}$ . On the other hand, it follows immediately from Lemma 2.7, (i), that the profinite group  $\Delta_{f_C}$ , hence also  $\Delta_{f_W}$ , does not have the property  $\mathbb{P}_{\mathcal{F} \rightarrow \infty}$ .

REFERENCES

- [1] A. Grothendieck, *Eléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I*, *Inst. Hautes Études Sci. Publ. Math.* No. **11** (1961).
- [2] A. Grothendieck, *Eléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II*, *Inst. Hautes Études Sci. Publ. Math.* No. **24** (1965).
- [3] A. Grothendieck, *Eléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III*, *Inst. Hautes Études Sci. Publ. Math.* No. **28** (1966).

- [4] A. Grothendieck, *Eléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. IV*, *Inst. Hautes Études Sci. Publ. Math.* No. **32** (1967).
- [5] A. Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Séminaire de Géométrie Algébrique du Bois Marie, 1962. Augmenté d'un exposé de Michèle Raynaud. With a preface and edited by Yves Laszlo. Revised reprint of the 1968 French original. Documents Mathématiques (Paris), **4**. Société Mathématique de France, Paris, 2005.
- [6] Y. Hoshi, On the fundamental groups of log configuration schemes, *Math. J. Okayama Univ.* **51** (2009), 1–26.
- [7] Y. Hoshi, The exactness of the log homotopy sequence, *Hiroshima Math. J.* **39** (2009), no. **1**, 61–121.
- [8] Y. Hoshi, The Grothendieck conjecture for hyperbolic polycurves of lower dimension, *J. Math. Sci. Univ. Tokyo* **21** (2014), no. **2**, 153–219.
- [9] H. Matsumura, *Commutative ring theory*, Translated from the Japanese by M. Reid. Cambridge Studies in Advanced Mathematics, **8**. Cambridge University Press, Cambridge, 1986.
- [10] K. Mitsui, Homotopy exact sequences and orbifolds, *Algebra Number Theory* **9** (2015), no. **5**, 1089–1136.
- [11] S. Mochizuki, Topics in absolute anabelian geometry I: generalities, *J. Math. Sci. Univ. Tokyo* **19** (2012), no. **2**, 139–242.
- [12] L. Ribes and P. Zalesskii, *Profinite groups*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, **40**. Springer-Verlag, Berlin, 2000.
- [13] K. Sawada, Pro- $p$  Grothendieck conjecture for hyperbolic polycurves, *Publ. Res. Inst. Math. Sci.* **54** (2018), no. **4**, 729–780.
- [14] A. Tamagawa, The Grothendieck conjecture for affine curves, *Compositio Math.* **109** (1997), no. **2**, 135–194.
- [15] *Revêtements étales et groupe fondamental (SGA 1)*, Séminaire de géométrie algébrique du Bois Marie 1960–61. Directed by A. Grothendieck. With two papers by M. Raynaud. Updated and annotated reprint of the 1971 original. Documents Mathématiques (Paris), **3**. Société Mathématique de France, Paris, 2003.

(Yuichiro Hoshi) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

*E-mail address:* `yuichiro@kurims.kyoto-u.ac.jp`