

# Free boundary problems for a viscous incompressible fluid

Yoshihiro SHIBATA\*

Department of Mathematical Sciences, School of Science and Engineering,  
Waseda University, Ohkubo 3-4-1, Shinjuku-ku, Tokyo 169-8555, Japan.  
e-mail address: yshibata@waseda.jp

Senjo SHIMIZU†

Faculty of Engineering, Shizuoka University  
Hamamatsu, Shizuoka 432-8561, Japan.  
e-mail address: tssshim@ipc.shizuoka.ac.jp

## 1 A review of free boundary problems

In this section we review free boundary problems for a viscous incompressible fluid. In writing the review we are indebted to the works due to Zadrzyńska [54], Solonnikov [36] and Nishida [21] that we consulted.

The domain  $\Omega_t \subset \mathbb{R}^d$  occupied by the fluid is given only on the initial time  $t = 0$ , while for  $t > 0$  it is to be determined. The velocity vector field  $v(x, t) = (v_1, \dots, v_d)$  and the pressure  $p(x, t)$  satisfy the Navier-Stokes equations

$$v_t + (v \cdot \nabla)v - \nu \Delta v + \nabla p = f, \quad \nabla \cdot v = 0 \quad x \in \Omega_t, \quad t > 0 \quad (1.1)$$

and suitable initial and boundary conditions, where  $\nu > 0$  is a constant coefficient of viscosity and  $f$  is a vector field of external forces. We classify three kinds of free boundary problems with respect to the geometry of the domain  $\Omega_t$ .

### 1.1 The motion of an isolated mass of a viscous fluid

This is the problem of describing the motion of an isolated mass of viscous fluid bounded by a free boundary. In the problem  $\Omega_t$  is a bounded domain, a free surface  $\partial\Omega_t = \Gamma_t$  is a compact, and initial and boundary conditions are given by

$$\begin{aligned} v|_{t=0} &= v_0(x) & x \in \Omega_0 \\ S n - \sigma H n|_{x \in \Gamma_t} &= 0. \end{aligned} \quad (1.2)$$

Here  $\sigma \geq 0$  is a constant coefficient of the surface tension.  $n$  is the unit outward normal vector to  $\Gamma_t$ .  $S = \nu D(v) - pI$  is the stress tensor,  $D(v)$  is the deformation tensor with the elements

---

\*Partly supported by Grants-in-Aid for Scientific Research (B) - 15340204, Japan Society for the Promotion of Science.

†Partly supported by Grants-in-Aid for Scientific Research (C) - 17540156, Japan Society for the Promotion of Science.

*2000 Mathematics Subject Classification.* 35Q30, 76D07.

*Key Words and Phrases.* Free boundary problem, Navier-Stokes equations, global in time unique existence, local in time unique existence, maximal regularity.

$\{D(v)\}_{ij} = \partial v_i / \partial x_j + \partial v_j / \partial x_i$ ,  $I$  is the identity matrix,  $H/(d-1)$  is the mean curvature of  $\Gamma_t$ . The sign of  $H$  is chosen in such a way that  $Hn = \Delta(t)x$ , where  $\Delta(t)$  is the Laplace-Beltrami operator on  $\Gamma_t$ .

The most of the existence results are obtained after transforming the free boundary problem written by Eulerian coordinates  $x$  to a problem in a fixed domain written by Lagrangean coordinates  $\xi$ . Let  $x(\xi, t)$  be a solution of the Cauchy problem

$$\frac{dx}{dt} = v(x, t), \quad x(0) = \xi. \quad (1.3)$$

Integrating (1.3), we connect Eulerian coordinates  $x$  with Lagrangean coordinates  $\xi$  by the formula

$$x = \xi + \int_0^t u(\xi, \tau) d\tau = X_u(\xi, t), \quad (1.4)$$

where  $u(\xi, t) = v(X_u(\xi, t), t)$ . A kinematic boundary condition at the free surface  $\Gamma_t = \{x = x(\xi, t) \mid \xi \in \Gamma\}$  is assumed. This expresses the fact that the free surface  $\Gamma_t$  consists for all  $t > 0$  of the same fluid particles, which do not leave it and are not incident on it from inside  $\Omega_t$ . Passing to Lagrangean coordinates  $x \in \Omega$  in (1.1) and (1.2), and setting  $p(X_u(\xi, t), t) = q(\xi, t)$ , we obtain

$$\begin{aligned} u_t - \nu \nabla_u^2 u + \nabla_u q &= f(X_u, t), \quad \nabla_u \cdot u = 0 \quad \xi \in \Omega, \quad t > 0 \\ u|_{t=0} &= v_0(\xi), \quad S_u n - \sigma \Delta(t) X_u|_{\xi \in \Gamma} = 0. \end{aligned} \quad (1.5)$$

Here

$$\begin{aligned} \nabla_u &= \mathcal{A} \nabla = \left\{ \sum_m A_{im} \frac{\partial}{\partial \xi_m} \right\}_{i=1, \dots, d} \\ S_u &= \nu D_u(u) - qI, \quad \{D_u(u)\}_{ij} = \sum_m \left( A_{im} \frac{\partial u_j}{\partial \xi_m} + A_{jm} \frac{\partial u_i}{\partial \xi_m} \right). \end{aligned}$$

$\mathcal{A}$  is the matrix with elements  $A_{im} = \partial \xi_m / \partial x_i|_{x=X_u(\xi, t)}$ ,  $n = n(X_u) = \mathcal{A}n_o / |\mathcal{A}n_o|$ , where  $n_o(\xi)$  is the unit outward normal to  $\partial\Omega = \Gamma$ .

### 1.1.1 The case of $\sigma = 0$

In this case the effect of surface tension on  $\Gamma_t$  is excluded. The pioneer work was done by Solonnikov [33] in 1977, he proved the local in time unique solvability of (1.5) in the framework of Hölder spaces  $C^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  by using the results of the linearized problem [32]. Later on, Solonnikov proved in [37] the local in time unique solvability of (1.5) for arbitrary initial data and the global in time unique solvability of (1.5) for  $f = 0$  and sufficiently small initial data in the class of isotropic Sobolev spaces  $W_p^{2,1}$  with  $d < p < \infty$  when  $d = 2, 3$ . Solonnikov used the technique of the hydrodynamical potentials with the estimates of the kernels of the corresponding singular integrals. Recently, we extended the Solonnikov results in [37] in the class of anisotropic Sobolev spaces  $W_{q,p}^{2,1}$  with  $d < q < \infty$  and  $2 < p < \infty$  of functions whose  $L_q$  norm with respect to the space variable together with the corresponding norms of the first and second spatial derivatives and the first time derivative are integrable with respect to time with the exponent  $p$ <sup>1</sup>.

The novelty of our results consists of two moments. First, we extend the results of Solonnikov by unbinding of the exponents of integrability with respect to the space and time variables.

<sup>1</sup>The results were announced in [29].

Second, we allow the exponent of integrability with respect to time  $p$  go down up to 2 providing the weaker setting than one that was allowed by Solonnikov whose case was  $p = q > d$ . We develop the semigroup approach which consists on the following. The linear operator obtained under the linearization of the free boundary problem generates the analytic semigroup on the functional space of  $q$ -integrable divergence free functions which was known due to Grubb and Solonnikov [16] (cf. Theorem 2.1 in § 2). This should allow us to apply some facts of the general analytic semigroup theory to obtain two coercive estimates (Theorems 2.3 and 2.6 in § 2) which are crucial for the study of the nonlinear problem. The core of our approach is to show the  $L_p$ - $L_q$  maximal regularity of the linearized problem global in time with exponential decay, which is stated in Theorem 2.3. One of our main issues to prove Theorem 2.3 is to use  $\mathcal{R}$ -boundedness and operator valued Fourier multiplier theorem which are recently developed by Weis [53], Denk, Hieber and Prüss [14] and Amann [4]. Methodologically, our approach seems to be simpler and more demonstrative than the estimates of kernel of singular integrals in the anisotropic Sobolev spaces. Thanks to the global in time  $L_p$ - $L_q$  maximal regularity of the linearized problem on the whole time interval  $(0, \infty)$  with exponential decay (Theorem 2.3), our proof of the global in time existence theorem (Theorem 2.5 in § 2) for the nonlinear problem (1.5) is much simpler than Solonnikov's proof [37]. In fact, we can show Theorem 2.5 simply by contraction mapping principle (cf. [31]). In the next section we state our results precisely.

Mucha and Zajączkowski considered the case where the self-gravitational force exists, namely  $f = \kappa \nabla U$ , where  $\kappa$  is the gravitational constant and  $U$  is the Newtonian potential. They proved in [20] the local in time unique solvability of (1.5) in  $W_p^{2,1}$  with  $d = 3$  and  $3 < p < \infty$  for arbitrary initial data. They used the local in time unique solvability result of the linearized problem of (1.5) which was proved in [19].

### 1.1.2 The case of $\sigma > 0$

In this case the effect of surface tension on  $\Gamma_t$  is included. Solonnikov formulated the local in time solvability of (1.5) in the Sobolev-Slobodetskii space  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $f = 0$  and arbitrary initial data in [34]. In [35], Solonnikov proved the global in time solvability of (1.5) in  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $f = 0$  provided that initial data are sufficiently small and the initial domain  $\Omega_0$  is sufficiently close to a ball. In [38] and [41], Solonnikov considered the case where the self-gravitational force exists. He proved in [41] the local in time unique solvability of (1.5) in  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  for arbitrary initial data, and in [38] the global in time unique solvability of (1.5) in  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $f = \kappa \nabla U$ , where  $\kappa$  is the gravitational constant and  $U$  is the Newtonian potential, provided that initial data are sufficiently small and the initial domain is sufficiently close to a ball. In [34, 35, 41, 38], he used the local in time unique solvability result of the linearized problem of (1.5) which was proved in [39].

Mogilievskiĭ and Solonnikov [18] proved the local in time solvability of (1.5) in Hölder spaces. Schweizer [27] proved the local in time unique existence of (1.5) for small initial data by using the semigroup approach. Padula and Solonnikov [26] proved the global in time unique solvability of (1.1) and (1.2) in Hölder spaces by using the mapping of  $\Omega_t$  on a ball instead of Lagrangean coordinates.

### 1.1.3 The case of $\sigma$ dependent on the temperature

This is an evolution problem of thermocapillary convection. Besides  $v$  and  $p$  satisfy (1.1), the temperature of the fluid  $\theta(x, t)$  satisfies

$$\theta_t + (v \cdot \nabla)\theta - \kappa\Delta\theta = \lambda|D(v)|^2 \quad x \in \Omega_t, \quad t > 0$$

and they satisfy the initial and boundary conditions, where  $\kappa$  and  $\lambda$  are positive constants or small positive functions of  $\theta$ .

Solonnikov proved in [40] the local in time unique solvability in  $C^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$ , and proved in [42] the global in time unique solvability in  $C^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  provided that the initial velocity and the initial temperature are sufficiently small and the domain is close to a ball. In the case  $\lambda = 0$ , Lagnova and Solonnikov [17] obtained the local in time unique solvability in Hölder spaces, and Wagner [52] obtained the local in time unique solvability in Sobolev spaces.

## 1.2 Two phase problems

This problem describes the motion of two liquids separated by free interface. Let  $\Omega$  be a bounded domain in  $\mathbb{R}^d$  ( $d \geq 2$ ) or the whole space. Let  $\Omega_t^+ \subset \Omega$  be occupied by the fluid of viscosity  $\nu^+ > 0$  and  $\Gamma_t$  be a boundary of  $\Omega_t^+$  which is strictly contained in  $\Omega$ . Put  $\Omega_t^- = \Omega \setminus (\Omega_t^+ \cup \Gamma_t)$ .  $\Omega_t^-$  is occupied by the fluid of viscosity  $\nu^- > 0$ . Given functions  $w^\pm$  defined on  $\Omega_t^\pm$ , we put

$$w = \begin{cases} w^+ & x \in \Omega_t^+, \quad t > 0 \\ w^- & x \in \Omega_t^-, \quad t > 0. \end{cases}$$

Moreover given function  $w$  defined on  $\Omega$ ,  $w^\pm$  denote the restriction of  $w$  to  $\Omega_t^\pm$ . The velocity vector field  $v^\pm$  and the pressure  $p^\pm$  satisfy the Navier-Stokes equations

$$\partial_t v + (v \cdot \nabla)v - \text{Div } S^\pm(v, p) = f, \quad \nabla \cdot v = 0 \quad \text{in } \Omega_t^\pm, \quad t > 0 \quad (1.6)$$

and the boundary and the initial conditions

$$\begin{aligned} & \left[ \lim_{x \rightarrow x_0 \in \Gamma_t, x \in \Omega_t^+} S^+(v, p) - \lim_{x \rightarrow x_0 \in \Gamma_t, x \in \Omega_t^-} S^-(v, p) \right] n|_{\Gamma_t} = \sigma H n|_{\Gamma_t} \\ & \lim_{x \rightarrow x_0 \in \Gamma_t, x \in \Omega_t^+} v|_{\Gamma_t} = \lim_{x \rightarrow x_0 \in \Gamma_t, x \in \Omega_t^-} v|_{\Gamma_t} \\ & v|_{\partial\Omega} = 0, \quad v|_{t=0} = v_0, \end{aligned} \quad (1.7)$$

where  $\sigma \geq 0$  is coefficient of the surface tension,  $n$  is the unit outward normal to  $\Gamma_t$  of  $\Omega_t^+$ , and  $S^\pm(v, p)$  are stress tensors defined by  $S^\pm(v, p) = \nu^\pm D(v) - pI$ . A kinematic boundary condition at the free interface  $\Gamma_t$  is assumed. When  $\Omega$  is the whole space, additional conditions are necessary.

In the case of a bounded  $\Omega$ , Tanaka [46] proved the global in time solvability of (1.6) and (1.7) in  $W_2^{2+\alpha}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $\sigma > 0$ ,  $d = 3$  and sufficiently small data with discontinuity of densities. Giga and Takahashi [15] and Takahashi [45] proved the global in time existence of weak solutions of (1.6) and (1.7) in the spaces such that the first derivative of the velocity in  $L_p$  with  $p > 2(d+1)$  with respect to time and space for  $\sigma = 0$  provided that  $\nu^+$  is close to  $\nu^-$ . Nouri and Poupanand [25] proved the local in time existence of a weak solution of the Navier-Stokes equation describing a multi fluid flow for arbitrary initial data for  $\sigma = 0$ . Recently, one

of the authors Shimizu obtained the global in time unique solvability of (1.6) and (1.7) for  $f = 0$  and sufficiently small initial data, and the local in time unique solvability of (1.6) and (1.7) for arbitrary initial data in  $W_{q,p}^{2,1}$  with  $2 < p < \infty$  and  $d < q < \infty$  when  $\sigma = 0$ . The proofs of the result will be given in a forthcoming paper.

In the case of the whole space  $\Omega$ , Denisova [10] proved the local in time unique solvability for arbitrary initial data in  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $\sigma \geq 0$  and  $d = 3$  with discontinuity of densities by using the local in time unique solvability result of the linearized problem in [8] and [11]. Denisova and Solonnikov [13] proved the local in time unique solvability for arbitrary initial data in the Hölder spaces with a power-like weight for  $\sigma > 0$  and  $d = 3$  with discontinuity of densities by using the local in time unique solvability result of the linearized problem in [9] and [12]. Recently, Abels [2] proved the existence of varifold and measure-valued varifold solutions for singular free interfaces for  $\sigma \geq 0$ .

### 1.3 Surface wave problems

This problem describes the motion of a fluid which occupies a semi-infinite domain in  $\mathbb{R}^d$  ( $d = 2, 3$ ) between the moving upper surface and a fixed bottom. Let

$$\Omega_t = \{x = (x', x_d) \in \mathbb{R}^d \mid x' \in \mathbb{R}^{d-1}, -b(x') < x_d < \eta(x', t)\}.$$

The velocity vector  $v$  and the pressure  $p$  satisfy the Navier-Stokes equation (1.1). The upper free surface  $S_F : x_d = \eta(x', t)$  satisfies the kinematic boundary condition

$$\eta_t = u_d - \sum_{k=1}^{d-1} (\partial_k \eta) u_k \quad \text{on } S_F. \quad (1.8)$$

The boundary condition on  $S_F$  is given by

$$pn_i - \nu D_{ij}(v)n_j = [g\eta - \sigma \nabla \cdot \{(1 + |\nabla \eta|^2)^{-\frac{1}{2}} \nabla \eta\}] n_i \quad \text{on } S_F, \quad (1.9)$$

where  $n = (n_1, \dots, n_d)$  is the outward normal to  $S_F$ ,  $g$  is the gravitation constant, and  $\sigma$  is the coefficient of surface tension. On the bottom surface  $S_B : x_d = -b(x')$  the boundary is impenetrable

$$v = 0 \quad \text{on } S_B. \quad (1.10)$$

The initial condition is the following

$$\eta = \eta_0(x') \quad x \in \mathbb{R}^{d-1}, \quad u = u_0(x) \quad x \in \Omega_0 \quad \text{at } t = 0. \quad (1.11)$$

The pioneer work of this problem was done by Beale [5] in 1980. Beale proved the local in time unique solvability for  $\sigma = 0$  and  $d = 3$  in the Bessel potential spaces  $H_2^{\ell, \frac{\ell}{2}}$  with  $3 < \ell < \frac{7}{2}$ . In [6], Beale proved the global in time unique solvability in  $H_2^{\ell, \frac{\ell}{2}}$  with  $3 < \ell < \frac{7}{2}$  for  $\sigma > 0$ ,  $d = 3$  and  $f = 0$  provided that the initial data  $\eta_0$  and  $u_0$  are sufficiently small. Beale and Nishida [7] obtained the asymptotic power-like in time decay of the global solutions. The local existence theorem for  $\sigma > 0$  and  $d = 2$  was established by Allain [3]. Tani [47] proved the local in time unique solvability for  $\sigma > 0$  and  $d = 3$  in  $W_2^{2+\alpha, 1+\frac{\alpha}{2}}$  with  $\alpha \in (\frac{1}{2}, 1)$ . Sylvester [44] showed the global in time solvability in  $H_2^{\ell, \frac{\ell}{2}}$  with  $\frac{9}{2} < \ell < 5$  for  $\sigma = 0$  and  $d = 3$  provided that initial data are sufficiently small by using Beale's method. Tani and Tanaka [48] proved the global in time solvability in  $W_2^{2+\alpha}$  with  $\alpha \in (\frac{1}{2}, 1)$  for  $\sigma \geq 0$  and  $d = 3$  provided that initial data are

sufficiently small by using Solonnikov's method. Recently, Abels [1] proved the local in time unique solvability in the isotropic Sobolev spaces  $W_p^{2,1}$  with  $d < p < \infty$ .

Nishida, Teramoto and Yoshihara [22] considered this problem under the assumption that the motion of fluid is horizontally periodic and that spatial mean of the motion of unknown free surface over the space period is equal to zero. They proved the global in time unique solvability and exponential stability in  $H_2^{\ell, \frac{\ell}{2}}$  with  $3 < \ell < \frac{7}{2}$  for sufficiently small initial data.

Teramoto has studied the motion of a viscous incompressible fluid which flows down an inclined plane under the effect of gravity. The fluid is bounded from below by a fixed plane which is inclined at an angle  $0 < \phi < \pi/2$  to the horizontal plane. He proved the local in time unique solvability for  $\sigma = 0$  and  $d = 3$  in [49], for  $\sigma > 0$  and  $d = 3$  in [50] by using Beale's idea. Nishida, Teramoto and Win [24] proved the global in time unique existence and stability for  $d = 2$  and sufficiently small initial data. Nishida, Teramoto and Yoshihara provided the Hopf bifurcation theorem in [23].

## 2 Results

As mentioned in the previous section, in this section we state our results precisely. We consider a time dependent problem with free surface for the Navier-Stokes equations which describes the motion of an isolated finite volume of viscous incompressible fluid without taking surface tension into account. The region  $\Omega_t \subset \mathbb{R}^d$ ,  $d \geq 2$ , occupied by the fluid is given only on the initial time  $t = 0$ , while for  $t > 0$  it is to be determined. The velocity vector field  $v(x, t) = (v_1, \dots, v_d)^*$ <sup>2</sup> and the pressure  $\theta(x, t)$  for  $x \in \Omega_t$  satisfy the Navier-Stokes equations:

$$\begin{aligned} v_t + (v \cdot \nabla)v - \operatorname{Div} S(v, \theta) &= f(x, t) && \text{in } \Omega_t, t > 0 \\ \operatorname{div} v &= 0 && \text{in } \Omega_t, t > 0 \\ S(v, \theta)n_t + \theta_0(x, t)n_t &= 0 && \text{in } \Gamma_t, t > 0 \\ v|_{t=0} &= v_0 && \text{on } \Omega. \end{aligned} \tag{2.1}$$

Here,  $\Gamma_t$  denotes the boundary of  $\Omega_t$ ,  $n_t(x)$  is the unit outward normal to  $\Gamma_t$  at the point  $x \in \Gamma_t$ ,  $\nabla = (\partial_1, \dots, \partial_d)$  with  $\partial_i = \partial/\partial x_i$ , and  $S(v, \theta)$  is the stress tensor defined by  $S(v, \theta) = D(v) - \theta I$  where  $D(v)$  is the deformation tensor of the velocities with elements  $D_{ij}(v) = \partial_i v_j + \partial_j v_i$  and  $I$  is the  $d \times d$  identity matrix. The external force  $f(x, t)$  and the pressure  $\theta_0(x, t)$  are functions defined on the whole space. In what follows, we may always assume that  $\theta_0(x, t) = 0$ , since we can arrive at this case by replacing  $\theta(x, t)$  by  $\theta + \theta_0$ .

Aside from the dynamical boundary condition, a further kinematic condition for  $\Gamma_t$  is satisfied. We write  $\Omega = \Omega_0$  and  $\Gamma = \Gamma_0$  and we assume that  $\Gamma$  is a  $C^{2,1}$  compact hypersurface.

Passing to Lagrangean coordinates in (2.1) and setting  $\theta(X_u(\xi, t), t) = \pi(\xi, t)$ , we obtain

$$\begin{aligned} u_t - \operatorname{Div} [S(u, \pi) + U(u, \pi)] &= f(X_u(\xi, t), t) && \text{in } \Omega \times (0, T) \\ \operatorname{div} u + E(u) = \operatorname{div} [u + \tilde{E}(u)] &= 0 && \text{in } \Omega \times (0, T) \\ [S(u, \pi) + U(u, \pi)]n &= 0 && \text{on } \Gamma \times (0, T) \\ u|_{t=0} &= u_0 && \text{in } \Omega, \end{aligned} \tag{2.2}$$

where  $u_0(\xi) = v_0(x)$ . Here and hereafter,  $n$  denotes the unit outward normal to  $\Gamma$ , and  $U(u, \pi)$ ,

---

<sup>2</sup> $M^*$  denotes the transpose of  $M$

$E(u)$  and  $\tilde{E}(u)$  are nonlinear terms of the following forms:

$$\begin{aligned} U(u, \pi) &= V_1\left(\int_0^t \nabla u \, d\tau\right) \nabla u + V_2\left(\int_0^t \nabla u \, d\tau\right) \pi \\ E(u) &= V_3\left(\int_0^t \nabla u \, d\tau\right) \nabla u, \quad \tilde{E}(u) = V_4\left(\int_0^t \nabla u \, d\tau\right) u \end{aligned}$$

with some polynomials  $V_j(\cdot)$  of  $\int_0^t \nabla u \, d\tau$ ,  $j = 1, 2, 3, 4$ , such as  $V_j(0) = 0$ . As a linearized problem of (2.2), we obtain the following Stokes equation with Neumann boundary condition:

$$\begin{aligned} u_t - \operatorname{Div} S(u, \pi) &= f && \text{in } \Omega \times (0, T) \\ \operatorname{div} u &= g = \operatorname{div} \tilde{g} && \text{in } \Omega \times (0, T) \\ S(u, \pi)n|_{\Gamma} &= h, \quad u|_{t=0} = u_0. \end{aligned} \tag{2.3}$$

In order to state our main results precisely, we introduce the function spaces and some symbols which will be used throughout the paper. For any domain  $D$  in  $\mathbb{R}^d$ , integer  $m$  and  $1 \leq q \leq \infty$ ,  $L_q(D)$  and  $W_q^m(D)$  denote the usual Lebesgue space and Sobolev space of functions defined on  $D$  with norms:  $\|\cdot\|_{L_q(D)}$  and  $\|\cdot\|_{W_q^m(D)}$ , respectively. And also, for any Banach space  $X$ , interval  $I$ , integer  $\ell$  and  $1 \leq p \leq \infty$ ,  $L_p(I, X)$  and  $W_p^\ell(I, X)$  denote the usual Lebesgue space and Sobolev space of the  $X$ -valued functions defined on  $I$  with norms:  $\|\cdot\|_{L_p(I, X)}$  and  $\|\cdot\|_{W_p^\ell(I, X)}$ , respectively. Set

$$\begin{aligned} W_{q,p}^{m,\ell}(D \times I) &= L_p(I, W_q^m(D)) \cap W_p^\ell(I, L_q(D)) \\ \|u\|_{W_{q,p}^{m,\ell}(D \times I)} &= \|u\|_{L_p(I, W_q^m(D))} + \|u\|_{W_p^\ell(I, L_q(D))} \\ W_{p,0}^\ell((0, T), X) &= \{u \in W_p^\ell((-\infty, T), X) \mid u = 0 \text{ for } t < 0\} \\ W_q^0(D) &= L_q(D), \quad W_p^0(I, X) = L_p(I, X), \quad W_{p,0}^0((0, T), X) = L_{p,0}((0, T), X). \end{aligned}$$

Given  $\alpha \geq 0$ , we set

$$\begin{aligned} \langle D_t \rangle^\alpha u(t) &= \mathcal{F}^{-1}[(1 + s^2)^{\alpha/2} \mathcal{F}u(s)](t) \\ H_p^\alpha(\mathbb{R}, X) &= \{u \in L_p(\mathbb{R}, X) \mid \langle D_t \rangle^\alpha u \in L_p(\mathbb{R}, X)\} \\ \|u\|_{H_p^\alpha(\mathbb{R}, X)} &= \|\langle D_t \rangle^\alpha u\|_{L_p(\mathbb{R}, X)} + \|u\|_{L_p(\mathbb{R}, X)}. \end{aligned}$$

Here and hereafter,  $\mathcal{F}$  and  $\mathcal{F}^{-1}$  denote the Fourier transform and its inverse formula, respectively. Set

$$\begin{aligned} H_{q,p}^{1,1/2}(D \times \mathbb{R}) &= H_p^{1/2}(\mathbb{R}, L_q(D)) \cap L_p(\mathbb{R}, W_q^1(D)) \\ H_{q,p,0}^{1,1/2}(D \times \mathbb{R}_+) &= \{u \in H_{q,p}^{1,1/2}(D \times \mathbb{R}) \mid u = 0 \text{ for } t < 0\} \\ \|u\|_{H_{q,p}^{1,1/2}(D \times \mathbb{R})} &= \|u\|_{H_p^{1/2}(\mathbb{R}, L_q(D))} + \|u\|_{L_p(\mathbb{R}, W_q^1(D))}. \end{aligned}$$

Finally, given  $0 < T < \infty$  we set

$$\begin{aligned} H_{q,p}^{1,1/2}(D \times (0, T)) &= \{u \mid \exists v \in H_{q,p}^{1,1/2}(D \times \mathbb{R}), \quad u = v \text{ on } D \times (0, T)\} \\ \|u\|_{H_{q,p}^{1,1/2}(D \times (0, T))} &= \inf \{\|v\|_{H_{q,p}^{1,1/2}(D \times \mathbb{R})} \mid \forall v \in H_{q,p}^{1,1/2}(D \times \mathbb{R}), \quad v = u \text{ on } D \times (0, T)\} \\ H_{q,p,0}^{1,1/2}(D \times (0, T)) &= \{u \mid \exists v \in H_{q,p,0}^{1,1/2}(D \times \mathbb{R}_+), \quad u = v \text{ on } D \times (0, T)\} \end{aligned}$$

$$\|u\|_{H_{q,p,0}^{1,1/2}(D \times (0,T))} = \inf \{ \|v\|_{H_{q,p}^{1,1/2}(D \times \mathbb{R})} \mid \forall v \in H_{q,p,0}^{1,1/2}(D \times \mathbb{R}_+), v = u \text{ on } D \times (0,T) \}.$$

Given Banach space  $X$  with norm  $\|\cdot\|_X$ , we set  $X^d = \{v = (v_1, \dots, v_d)^* \mid v_j \in X\}$ ,  $\|v\|_X = \sum_{j=1}^d \|v_j\|_X$ . The dot  $\cdot$  denotes the inner-product of  $\mathbb{R}^d$ .  $F = (F_{ij})$  means an  $d \times d$  matrix whose  $i$ -th column and  $j$ -th row component is  $F_{ij}$ . For the differentiation of an  $d \times d$  matrix of functions  $F = (F_{ij})$ , an  $d$ -vector of functions  $u = (u_1, \dots, u_d)^*$  and a scalar function  $\theta$ , we use the following symbols:  $\theta_t = \partial_t \theta = \partial \theta / \partial t$ ,  $\partial_j \theta = \partial \theta / \partial x_j$ ,

$$\begin{aligned} \nabla \theta &= (\partial_1 \theta, \dots, \partial_d \theta)^*, \quad \nabla^k \theta = (\partial_x^\alpha \theta \mid |\alpha| = k), \quad u_t = \partial_t u = (\partial_t u_1, \dots, \partial_t u_d), \quad \nabla u = (\partial_i u_j), \\ \nabla^k u &= (\partial_x^\alpha u_i, |\alpha| = k, i = 1, \dots, d), \quad \operatorname{div} u = \sum_{j=1}^d \partial_j u_j, \quad \operatorname{Div} F = \left( \sum_{j=1}^d \partial_j F_{1j}, \dots, \sum_{j=1}^d \partial_j F_{dj} \right)^*. \end{aligned}$$

The inner products  $(\cdot, \cdot)_\Omega$  and  $(\cdot, \cdot)_\Gamma$  are defined by  $(u, v)_\Omega = \int_\Omega u(x) \cdot v(x) dx$  and  $(u, v)_\Gamma = \int_\Gamma u(x) \cdot v(x) d\sigma$  where  $d\sigma$  denotes the surface element of  $\Gamma$ . By  $C$  we denote a generic constant and  $C_{a,b,\dots}$  denotes the constant depending on the quantities  $a, b, \dots$ . The constants  $C$  and  $C_{a,b,\dots}$  may change from line to line.

In order to introduce our class of initial data for (2.2), we discuss an analytic semigroup approach to the initial boundary value problem:

$$\begin{aligned} u_t - \operatorname{Div} S(u, \pi) &= 0, \quad \operatorname{div} u = 0 \quad \text{in } \Omega \times (0, \infty) \\ S(u, \pi)n|_\Gamma &= 0, \quad u|_{t=0} = u_0. \end{aligned} \quad (2.4)$$

Since the time derivative of  $\pi$  is missing in (2.4), to obtain the evolution equation for  $u$  we have to eliminate  $\pi$  from (2.4). To do this, for a while instead of (2.4) we shall consider the resolvent problem:

$$\lambda u - \operatorname{Div} S(u, \pi) = f, \quad \operatorname{div} u = 0 \quad \text{in } \Omega \times (0, \infty), \quad S(u, \pi)n|_\Gamma = 0 \quad (2.5)$$

and we shall discuss how to eliminate  $\pi$  from (2.5). We introduce the second Helmholtz decomposition corresponding to (2.4). Set

$$\begin{aligned} J_q(\Omega) &= \{u = (u_1, \dots, u_d)^* \in L_q(\Omega)^d \mid \operatorname{div} u = 0 \text{ in } \Omega\} \\ G_q(\Omega) &= \{\nabla \pi \mid \pi \in W_q^1(\Omega), \pi|_\Gamma = 0\}. \end{aligned}$$

Then, by Grubb and Solonnikov [16] (cf. also Shibata and Shimizu [28]) we know that

$$L_q(\Omega)^d = J_q(\Omega) \oplus G_q(\Omega)$$

for  $1 < q < \infty$ , where  $\oplus$  denotes the direct sum. Let  $P_q$  be the solenoidal projection:  $L_q(\Omega)^d \rightarrow J_q(\Omega)$  along  $G_q(\Omega)$ . Then, substituting the 2nd Helmholtz decomposition of  $f$ :  $f = P_q f + \nabla \theta$ , where  $\theta \in W_q^1(\Omega)$  and  $\theta|_\Gamma = 0$ , into (2.5), we have

$$\lambda v - \operatorname{Div} S(v, \pi - \theta) = P_q f, \quad \operatorname{div} v = 0 \quad \text{in } \Omega, \quad S(v, \pi - \theta)n|_\Gamma = 0. \quad (2.6)$$

Denoting  $\pi - \theta$  by  $\pi$  in (2.6) again, from now on we consider (2.5) under the condition that  $\operatorname{div} f = 0$ . Then, applying the divergence to (2.5) and multiplying the boundary condition by  $n$ , we have

$$\Delta \theta = 0 \quad \text{in } \Omega, \quad \theta|_\Gamma = [D(v)n] \cdot n - \operatorname{div} v|_\Gamma, \quad (2.7)$$

where we have used the facts that  $\operatorname{div} v = 0$  in  $\Omega$  and  $n \cdot n = 1$  on  $\Gamma$ . We know that given  $v \in W_q^2(\Omega)^d$  there exists a unique  $\theta \in W_q^1(\Omega)$  which solves (2.7) and enjoys the estimate:

$$\|\theta\|_{W_q^1(\Omega)} \leq C \|v\|_{W_q^2(\Omega)}.$$

From this point of view, let us define the map  $K : W_q^2(\Omega)^d \rightarrow W_q^1(\Omega)$  by  $\theta = K(v)$  for  $v \in W_q^2(\Omega)^d$ . By using this symbol, the equation (2.5) is rewritten in the form:

$$\lambda v - \operatorname{Div} S(v, K(v)) = f \quad \text{in } \Omega, \quad S(v, K(v))n|_\Gamma = 0 \quad (2.8)$$

for  $f \in J_q(\Omega)$ . We set

$$\begin{aligned} A_q u &= -\operatorname{Div} S(u, K(u)) \quad \text{for } u \in \mathcal{D}(A_q) \\ \mathcal{D}(A_q) &= \{u \in J_q(\Omega) \cap W_q^2(\Omega)^d \mid S(u, K(u))n|_\Gamma = 0\}. \end{aligned}$$

From Grubb and Solonnikov [16] and Shibata and Shimizu [28], we know the following theorem.

**Theorem 2.1.** *Let  $1 < q < \infty$ . Then,  $A_q$  generates an analytic semigroup  $\{e^{-A_q t}\}_{t \geq 0}$  on  $J_q(\Omega)$ .*

**Remark 2.2.** The function  $e^{-A_q t} u_0$  is an initial flow for (2.2).

Now, we shall state our results. For the initial data, we introduce the following space:

$$\mathcal{D}_{q,p}(\Omega) = [J_q(\Omega), \mathcal{D}(A_q)]_{1-1/p,p}.$$

Here and hereafter,  $[\cdot, \cdot]_{\theta,p}$  denotes the real interpolation functor. In order to state a global in time existence theorem for (2.2), we introduce the rigid space  $\mathcal{R}_{gd}$  which is defined by the relation:

$$\mathcal{R}_{gd} = \{Ax + b \mid A : d \times d \text{ anti-symmetric matrix, } b \in \mathbb{R}^d\}.$$

In what follows, we denote the basis of  $\mathcal{R}_{gd}$  by  $\{p_\ell\}_{\ell=1}^M$ , which are normalized such as  $(p_\ell, p_m)_\Omega = \delta_{\ell m}$  ( $\ell, m = 1, \dots, M$ ), where  $\delta_{\ell m}$  are Kronecker's delta symbols.

The first theorem is the main result which shows the global in time  $L_p$ - $L_q$  maximal regularity of (2.3) with exponential stability.

**Theorem 2.3.** *Let  $1 < p, q < \infty$ . Then, there exists a  $\gamma_0 > 0$  such that if  $u_0, f, g, \tilde{g}$  and  $h$  satisfy the following conditions:*

$$\begin{aligned} u_0 &\in \mathcal{D}_{q,p}(\Omega), \quad e^{\gamma t} f \in L_p((0, \infty), L_q(\Omega))^d, \quad e^{\gamma t} g \in L_{p,0}((0, \infty), W_q^1(\Omega)) \\ e^{\gamma t} \tilde{g} &\in W_{p,0}^1((0, \infty), L_q(\Omega))^d, \quad e^{\gamma t} h \in H_{q,p,0}^{1,1/2}(\Omega \times (0, \infty))^d \end{aligned}$$

for some  $\gamma \in [0, \gamma_0]$ , and

$$(u_0, p_\ell)_\Omega = 0, \quad (f(\cdot, t), p_\ell)_\Omega + (h(\cdot, t), g_\ell)_\Gamma = 0$$

for  $t \geq 0$  and  $\ell = 1, \dots, M$ , then the equation (2.3) with  $T = \infty$  admits a unique solution

$$(u, \pi) \in W_{q,p}^{2,1}(\Omega \times (0, \infty))^d \times L_p((0, \infty), W_q^1(\Omega)).$$

Moreover there exists  $\tilde{\pi} \in H_{q,p}^{1,1/2}(\Omega \times (0, \infty))$  such that  $\tilde{\pi} = \pi$  on  $\Gamma \times (0, \infty)$ . The solution satisfies the estimates:

$$\begin{aligned} &\|e^{\gamma t} u\|_{W_{q,p}^{2,1}(\Omega \times (0, \infty))} + \|e^{\gamma t} \pi\|_{L_p((0, \infty), W_q^1(\Omega))} + \|e^{\gamma t} \tilde{\pi}\|_{H_{q,p}^{1,1/2}(\Omega \times (0, \infty))} \leq C \{ \|u_0\|_{\mathcal{D}_{q,p}(\Omega)} \\ &\quad + \|e^{\gamma t} f\|_{L_p((0, \infty), L_q(\Omega))} + \|e^{\gamma t} g\|_{L_p(\mathbb{R}, W_q^1(\Omega))} + \|e^{\gamma t} \tilde{g}\|_{W_p^1(\mathbb{R}, L_q(\Omega))} + \|e^{\gamma t} h\|_{H_{q,p,0}^{1,1/2}(\Omega \times (0, \infty))} \} \end{aligned}$$

and the condition:  $(u(\cdot, t), p_\ell)_\Omega = 0$  for  $t \geq 0$  and  $\ell = 1, \dots, M$ .

**Remark 2.4.** Let us define the Besov space  $B_{q,p}^{2(1-1/p)}(\Omega)$  by the real interpolation:

$$B_{q,p}^{2(1-1/p)}(\Omega) = [L_q(\Omega), W_q^2(\Omega)]_{1-1/p,p}$$

and set

$$JB_{q,p}^{2(1-1/p)}(\Omega) = \{u \in B_{q,p}^{2(1-1/p)}(\Omega)^d \mid \operatorname{div} u = 0 \text{ in } \Omega\}.$$

Then, we see that  $\mathcal{D}_{q,p}(\Omega) \subset JB_{q,p}^{2(1-1/p)}(\Omega)$ . Moreover, from Proposition 2.13 combined with Remarks 2.7 (c) in Steiger [43] (cf. also Triebel [51]) it follows that

$$\mathcal{D}_{q,p}(\Omega) = \begin{cases} \{v \in JB_{q,p}^{2(1-1/p)}(\Omega) \mid S(v, K(v))n|_{\Gamma} = 0\} & 2(1-1/p) > 1+1/q \\ JB_{q,p}^{2(1-1/p)}(\Omega) & 2(1-1/p) < 1+1/q. \end{cases}$$

The next theorem shows the global in time unique solvability of (2.2) for  $f = 0$  and sufficiently small initial data which are orthogonal to the rigid space, which is proved by using the contraction mapping principle based on Theorem 2.3.

**Theorem 2.5.** *Let  $2 < p < \infty$  and  $d < q < \infty$ . We consider the case where  $T = \infty$  and  $f = 0$  in (2.2). Then, there exist positive numbers  $\epsilon$  and  $\gamma$  such that if  $u_0 \in \mathcal{D}_{q,p}(\Omega)$ ,  $\|u_0\|_{\mathcal{D}_{q,p}(\Omega)} \leq \epsilon$  and  $(u_0, p_\ell)_\Omega = 0$  for  $\ell = 1, \dots, M$ , then the equation (2.2) with  $T = \infty$  and  $f = 0$  admits a unique solution*

$$(u, \pi) \in W_{q,p}^{2,1}(\Omega \times (0, \infty))^d \times L_p((0, \infty), W_q^1(\Omega)).$$

Moreover there exists  $\tilde{\pi} \in H_{q,p}^{1,1/2}(\Omega \times (0, \infty))$  such that  $\tilde{\pi} = \pi$  on  $\Gamma \times (0, \infty)$ . The solution satisfies the estimate:

$$\|e^{\gamma t} u\|_{W_{q,p}^{2,1}(\Omega \times (0, \infty))} + \|e^{\gamma t} \pi\|_{L_p((0, \infty), W_q^1(\Omega))} + \|e^{\gamma t} \tilde{\pi}\|_{H_{q,p}^{1,1/2}(\Omega \times (0, \infty))} \leq C\epsilon$$

for some  $\gamma > 0$  and the condition:

$$(u(\cdot, t), p_\ell)_\Omega = 0 \quad \text{for } \ell = 1, \dots, M \text{ and } t \geq 0.$$

The next theorem shows the  $L_p$ - $L_q$  maximal regularity of (2.3) local in time.

**Theorem 2.6.** *Let  $1 < p, q < \infty$  and  $0 < T < \infty$ . If  $u_0, f, g, \tilde{g}$  and  $h$  satisfy the condition:*

$$\begin{aligned} u_0 \in \mathcal{D}_{q,p}(\Omega), \quad f \in L_p((0, T), L_q(\Omega))^d, \quad g \in L_{p,0}((0, T), W_q^1(\Omega)) \\ \tilde{g} \in W_{p,0}^1((0, T), L_q(\Omega))^d, \quad h \in H_{q,p,0}^{1,1/2}(\Omega \times (0, T))^d \end{aligned}$$

then the equation (2.3) admits a unique solution

$$(u, \pi) \in W_{q,p}^{2,1}(\Omega \times (0, T))^d \times L_p((0, T), W_q^1(\Omega)).$$

Moreover there exists  $\tilde{\pi} \in H_{q,p}^{1,1/2}(\Omega \times (0, T))$  such that  $\tilde{\pi} = \pi$  on  $\Gamma \times (0, T)$ . The solution satisfies the estimate:

$$\begin{aligned} \|u\|_{W_{q,p}^{2,1}(\Omega \times (0, T))} + \|\pi\|_{L_p((0, T), W_q^1(\Omega))} + \|\tilde{\pi}\|_{H_{q,p}^{1,1/2}(\Omega \times (0, T))} \leq C(1+T) \{ \|u_0\|_{\mathcal{D}_{q,p}(\Omega)} \\ + \|f\|_{L_p((0, T), L_q(\Omega))} + \|g\|_{L_p((0, T), W_q^1(\Omega))} + \|\tilde{g}\|_{W_{p,0}^1((0, T), L_q(\Omega))} + \|h\|_{H_{q,p,0}^{1,1/2}(\Omega \times [0, T])} \} \end{aligned} \quad (2.9)$$

where the constant  $C$  is independent of  $T, u, \pi, u_0, f, g, \tilde{g}$  and  $h$ .

The next theorem shows the local in time unique solvability of (2.2) for any initial data and right member of  $f$ , which is proved by using the contraction mapping principle based on Theorem 2.6.

**Theorem 2.7.** *Let  $2 < p < \infty$  and  $d < q < \infty$ . Then, for any  $R > 0$  and  $R' > 0$  there exists a time  $T > 0$  depending on  $R$  and  $R'$  such that the equation (2.2) admits a unique solution*

$$(u, \pi) \in W_{q,p}^{2,1}(\Omega \times (0, T))^d \times L_p((0, T), W_q^1(\Omega)).$$

Moreover there exists  $\tilde{\pi} \in H_{q,p}^{1,1/2}(\Omega \times (0, T))$  such that  $\tilde{\pi} = \pi$  on  $\Gamma \times (0, T)$ . The solution satisfies the estimate:

$$\|u\|_{W_{q,p}^{2,1}(\Omega \times (0, T))} + \|\pi\|_{L_p((0, T), W_q^1(\Omega))} + \|\tilde{\pi}\|_{H_{q,p}^{1,1/2}(\Omega \times (0, T))} \leq CR$$

for some constant  $C$  depending essentially only on  $p$  and  $q$  provided that  $u_0 \in \mathcal{D}_{q,p}(\Omega)$ ,  $f \in L_p(\mathbb{R}_+, L_q(\mathbb{R}^d))$ ,  $\nabla f \in L_\infty(\mathbb{R}^d \times \mathbb{R}_+)$  and

$$\|u_0\|_{\mathcal{D}_{q,p}(\Omega)} + \|f\|_{L_p(\mathbb{R}_+, L_q(\mathbb{R}^d))} \leq R, \quad \|\nabla f\|_{L_\infty(\mathbb{R}^d \times \mathbb{R}_+)} \leq R'.$$

The proofs of Theorems 2.3 and 2.6 are given in [30] and those of Theorems 2.5 and 2.7 are given in [31].

## References

- [1] H. Abels, *The initial-Value problem for the Navier-Stokes equations with a free surface in  $L_q$  Sobolev spaces*, Adv. Differential Equations, **10** (2005) 45–64.
- [2] H. Abels, *On generalized solutions of two-phase flows for viscous incompressible fluids*, Preprint (2005).
- [3] G. Allain, *Small-time existence for the Navier-Stokes equations with a free surface*, Appl. Math. Optim., **16** (1987) 37–50.
- [4] H. Amann, *Linear and Quasilinear Parabolic Problems Vol. I Abstract Linear Theory*, Monographs in Math., Vol 89, 1995, Birkhäuser Verlag, Basel·Boston·Berlin
- [5] J. T. Beale, *The initial value problem for the Navier-Stokes equations with a free boundary*, Comm. Pure Appl. Math., **31** (1980) 359–392.
- [6] J. T. Beale, *Large time regularity of viscous surface waves*, Arch. Rat. Mech. Anal., **84** (1984) 307–352.
- [7] J. T. Beale and T. Nishida, *Large time behavior of viscous surface waves*, Lecture Notes in Numer. Appl. Anal., **8** (1985) 1–14.
- [8] I. V. Denisova, *A priori estimates for the solution of a linear time-dependent problem connected with the motion of a drop in a fluid medium*, Trudy Mat. Inst. Steklov., **188** (1990) 3–21 (in Russian); English transl.: Proc. Steklov Inst. Math., **188** (1991) 1–24.
- [9] I. V. Denisova, *Solvability in Hölder spaces of a linear problem concerning the motion of two fluids separated by a closed surface*, Algebra i Analiz, **5** (1993) 122–148 (in Russian); English transl.: St.Petersburg Math. J., **5** (1994) 765–787.

- [10] I. V. Denisova, *Problem of the motion of two viscous incompressible fluids separated by a closed free interface*, Acta Appl. Math., **37** (1994) 31–40.
- [11] I. V. Denisova and V. A. Solonnikov, *Solvability of a linearized problem on the motion of a drop in a fluid flow*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), **171** (1989) 53–65 (in Russian); English transl.: J. Soviet Math., **56** (1991) 2309–2316.
- [12] I. V. Denisova and V. A. Solonnikov, *Solvability in Hölder spaces of a model initial-boundary value problem generated by a problem on the motion of two fluids*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), **181** (1991) 5–44 (in Russian); English transl.: J. Math. Sci., **70** (1994) 1717–1746.
- [13] I. V. Denisova and V. A. Solonnikov, *Classical solvability of the problem on the the motion of two viscous incompressible fluids*, Algebra i Analiz **7** (1995) 101–142 (in Russian); English transl.: St.Petersburg Math. J., **7** (1996) 755–786.
- [14] R. Denk, M. Hieber and J. Prüss,  *$\mathcal{R}$ -boundedness, Fourier multipliers and problems of elliptic and parabolic type*, Memoirs of AMS, Vol 166, No. 788, 2003.
- [15] Y. Giga and Sh. Takahashi, *On global weak solutions of the nonstationary two-phase Stokes flow*, SIAM J. Math. Anal., **25** (1994) 876–893.
- [16] G. Grubb and V. A. Solonnikov, *Boundary value problems for the nonstationary Navier-Stokes equations treated by pseudo-differential method*, Math. Scand., **69** (1991) 217–290.
- [17] M. V. Lagunova and V. A. Solonnikov, *Nonstationary problem of thermocapillary convection*, Stability Appl. Anal. Contin. Media, **1** (1991) 47–72.
- [18] I. Sh. Mogilevskii and V. A. Solonnikov, *On the solvability of a free boundary problem for the Navier-Stokes equations in the Hölder spaces of functions*, Nonlinear Analysis. A Tribute in Honour of Giovanni Prodi, Quaderni, Pisa (1991) 257–272.
- [19] P. B. Mucha and W. Zajączkowski, *On the existence for the Cauchy-Neumann problem for the Stokes system in the  $L_p$ -framework*, Studia Mathematica, **143** (2000) 75–101.
- [20] P. B. Mucha and W. Zajączkowski, *On local existence of solutions of the free boundary problem for an incompressible viscous self-gravitating fluid motion*, Applicationes Mathematicae, **27** (2000) 319–333.
- [21] T. Nishida, *Equations of fluid dynamics – free surface problems*, Comm. Pure Appl. Math., **39** (1986) 221–238.
- [22] T. Nishida, Y. Teramoto and H. Yoshihara *Global in time behavior of viscous surface waves: horizontally periodic motion*, J. Math. Kyoto Univ., **44** (2004) 271–323.
- [23] T. Nishida, Y. Teramoto and H. Yoshihara *Hopf bifurcation in viscous incompressible flow down an inclined plane*, J. math. fluid mech., **7** (2005) 29–71.
- [24] T. Nishida, Y. Teramoto and H. A. Win *Navier-Stokes flow down an inclined plane: downward periodic motion*, J. Math. Kyoto Univ., **33** (1993) 787–801.
- [25] A. Nouri and F. Poupaud *An existence theorem for the multifluid Navier-Stokes problem*, J. Differential Equations, **123** (1995) 71–88.

- [26] M. Padula, V. A. Solonnikov *On the global existence of nonsteady motions of a fluid drop and their exponential decay to a uniform rigid rotation*, Quad. Mat., **10** (2002) 185–218.
- [27] B. Schweizer, *Free boundary fluid systems in a semigroup approach and oscillatory behavior*, SIAM J. Math. Anal., **28** (1997) 1135–1157.
- [28] Y. Shibata and S. Shimizu, *On a resolvent estimate for the Stokes system with Neumann boundary condition*, Differential Integral Equations, **16** (2003) 385–426.
- [29] Y. Shibata and S. Shimizu,  *$L_p$ - $L_q$  maximal regularity and viscous incompressible flows with free surface*, Proc. Japan Acad. Ser. A Math. Sci., **81** (2005) 151–155.
- [30] Y. Shibata and S. Shimizu, *A On the  $L_p$ - $L_q$  maximal regularity of the Neumann problem for the Stokes equations in a bounded domain*, Preprint (2005).
- [31] Y. Shibata and S. Shimizu, *On some free boundary problem for the Navier-Stokes equations*, Preprint (2005).
- [32] V. A. Solonnikov, *On the solvability of the second initial-boundary value problem for the linear nonstationary Navier-Stokes system*, Zap. Nauchn. Sem. (LOMI), **69** (1977), 1388–1424 (in Russian); English transl.: J. Soviet Math., **10** (1978) 141–193.
- [33] V. A. Solonnikov, *Solvability of a problem on the motion of a viscous incompressible fluid bounded by a free surface*, Izv. Akad. Nauk SSSR Ser. Mat., **41** (1977), 1388–1424 (in Russian); English transl.: Math. USSR Izv., **11** (1977) 1323–1358.
- [34] V. A. Solonnikov, *Solvability of the evolution problem for an isolated mass of a viscous incompressible capillary liquid*, Zap. Nauchn. Sem. (LOMI), **140** (1984) 179–186 (in Russian); English transl.: J. Soviet Math., **32** (1986) 223–238.
- [35] V. A. Solonnikov, *Unsteady motion of a finite mass of fluid, bounded by a free surface*, Zap. Nauchn. Sem. (LOMI), **152** (1986) 137–157 (in Russian); English transl.: J. Soviet Math., **40** (1988) 672–686.
- [36] V. A. Solonnikov, *Free boundary problems and problems in non compact domains for the Navier-Stokes equations*, Proc. Inter. Congress of Math., (Berkeley, 1986) 1113–1122.
- [37] V. A. Solonnikov, *On the transient motion of an isolated volume of viscous incompressible fluid*, Izv. Akad. Nauk SSSR Ser. Mat., **51** (1987) 1065–1087 (in Russian); English transl.: Math. USSR Izv., **31** (1988) 381–405.
- [38] V. A. Solonnikov, *On nonstationary motion of a finite isolated mass of self-gravitating fluid*, Algebra i Analiz, **1** (1989) 207–249 (in Russian); English transl.: Leningrad Math. J., **1** (1990) 227–276.
- [39] V. A. Solonnikov, *On an initial-boundary value problem for the Stokes system arising in the study of a problem with a free boundary*, Trudy Mat. Inst. Steklov., **188** (1990) 150–188 (in Russian); English transl.: Proc. Steklov Inst. Math., **3** (1991) 191–239.
- [40] V. A. Solonnikov, *On an evolution problem of thermocapillary convection*, Le Matematiche, **66** (1991) 449–460.

- [41] V. A. Solonnikov, *Solvability of the problem of evolution of a viscous incompressible fluid bounded by a free surface on a finite time interval*, Algebra i Analiz, **3** (1991) 222–257 (in Russian); English transl.: St. Petersburg Math. J., **3** (1992) 189–220.
- [42] V. A. Solonnikov, *On an evolution problem of thermocapillary convection*, Internat. Ser. Numer. Math., **106**, Birkhäuser (1992) 301–317.
- [43] O. Steiger, *On Navier-Stokes equations with first order boundary conditions*, Ph.D. thesis, Universität Zürich, 2004.
- [44] D. Sylvester, *Large time existence of small viscous surface waves without surface tension*, Commun. Partial Differential Equations, **15** (1990) 823–903.
- [45] Sh. Takahashi, *On global weak solutions of the nonstationary two-phase Navier-Stokes flow*, Adv. Math. Sci. Appl., **5** (1995) 321–342.
- [46] N. Tanaka, *Global existence of two phase non-homogeneous viscous incompressible weak fluid flow*, Commun. Partial Differential Equations, **18** (1993) 41–81.
- [47] A. Tani, *Small-time existence for the three-dimensional incompressible Navier-Stokes equations with a free surface*, Arch. Rat. Mech. Anal., **133** (1996) 299–331.
- [48] A. Tani and N. Tanaka, *Large time existence of surface waves in incompressible viscous fluids with or without surface tension*, Arch. Rat. Mech. Anal., **130** (1995) 303–314.
- [49] Y. Teramoto, *The initial value problem for a viscous incompressible flow down an inclined plane*, Hiroshima Math. J., **15** (1985) 619–643.
- [50] Y. Teramoto, *On the Navier-Stokes flow down an inclined plane*, J. Math. Kyoto Univ., **32** (1992) 593–619.
- [51] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, 2nd edition, Johann Ambrosius Barth, Heidelberg, 1995.
- [52] A. Wagner, *Nonstationary Marangoni convection*, Appl. Math., **26** (1999) 195–220.
- [53] L. Weis, *Operator-valued Fourier multiplier theorems and maximal  $L_p$ -regularity*, Math. Ann., **319** (2001) 735–758.
- [54] E. Zadrzyńska, *Free boundary problems for nonstationary Navier-Stokes equations*, Dissertationes Mathematicae Vol 424, Polish Academy of Sciences (2004).