# On the $K(\pi, 1)$ -property for rings of integers in the mixed case

By

Alexander SCHMIDT<sup>\*</sup>

#### Abstract

We investigate the Galois group  $G_S(p)$  of the maximal *p*-extension unramified outside a finite set *S* of primes of a number field in the (mixed) case, when there are primes dividing *p* inside and outside *S*. We show that the cohomology of  $G_S(p)$  is 'often' isomorphic to the étale cohomology of the scheme  $Spec(\mathcal{O}_k \setminus S)$ , in particular,  $G_S(p)$  is of cohomological dimension 2 then. We deduce this from the results in our previous paper [Sch2], which mainly dealt with the tame case.

# §1. Introduction

Let Y be a connected locally noetherian scheme and let p be a prime number. We denote the étale fundamental group of Y by  $\pi_1(Y)$  and its maximal pro-p factor group by  $\pi_1(Y)(p)$ . The Hochschild-Serre spectral sequence induces natural homomorphisms

$$\phi_i: H^i(\pi_1^{et}(Y)(p), \mathbb{Z}/p\mathbb{Z}) \longrightarrow H^i_{et}(Y, \mathbb{Z}/p\mathbb{Z}), \ i \ge 0,$$

and we call Y a ' $K(\pi, 1)$  for p' if all  $\phi_i$  are isomorphisms; see [Sch2] Proposition 2.1 for equivalent conditions. See [Wi2] for a purely Galois cohomological approach to the  $K(\pi, 1)$ -property. Our main result is the following

**Theorem 1.1.** Let k be a number field and let p be a prime number. Assume that k does not contain a primitive p-th root of unity and that the class number of k is prime to p. Then the following holds:

Let S be a finite set of primes of k and let T be a set of primes of k of Dirichlet density  $\delta(T) = 1$ . Then there exists a finite subset  $T_1 \subset T$  such that  $Spec(\mathcal{O}_k) \setminus (S \cup T_1)$ is a  $K(\pi, 1)$  for p.

Received January 15, 2008. Revised June 2, 2008.

<sup>2000</sup> Mathematics Subject Classification(s): 11R34, 12G10

<sup>\*</sup>NWF I-Mathematik, Universität Regensburg, 93040 Regensburg, Deutschland.

e-mail: alexander.schmidt@mathematik.uni-regensburg.de

<sup>© 2009</sup> Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

**Remarks.** 1. If S contains the set  $S_p$  of primes dividing p, then Theorem 1.1 holds with  $T_1 = \emptyset$  and even without the condition  $\zeta_p \notin k$  and Cl(k)(p) = 0, see [Sch2], Proposition 2.3. In the tame case  $S \cap S_p = \emptyset$ , the statement of Theorem 1.1 is the main result of [Sch2]. Here we provide the extension to the 'mixed' case  $\emptyset \subsetneq S \cap S_p \subsetneq S_p$ .

2. For a given number field k, all but finitely many prime numbers p satisfy the condition of Theorem 1.1. We conjecture that Theorem 1.1 holds without the restricting assumption on p.

Let S be a finite set of places of a number field k. Let  $k_S(p)$  be the maximal p-extension of k unramified outside S and put  $G_S(p) = Gal(k_S(p)|k)$ . If  $S_{\mathbb{R}}$  denotes the set of real places of k, then  $G_{S \cup S_{\mathbb{R}}}(p) \cong \pi_1(Spec(\mathcal{O}_k) \setminus S)(p)$  (we have  $G_S(p) = G_{S \cup S_{\mathbb{R}}}(p)$ if p is odd or k is totally imaginary). The following Theorem 1.2 sharpens Theorem 1.1.

**Theorem 1.2.** The set  $T_1 \subset T$  in Theorem 1.1 may be chosen such that

- (i)  $T_1$  consists of primes  $\mathfrak{p}$  of degree 1 with  $N(\mathfrak{p}) \equiv 1 \mod p$ ,
- (ii)  $(k_{S\cup T_1}(p))_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$  for all primes  $\mathfrak{p} \in S \cup T_1$ .

Note that Theorem 1.2 provides nontrivial information even in the case  $S \supset S_p$ , where assertion (ii) was only known when k contains a primitive p-th root of unity (Kuz'min's theorem, see [Kuz] or [NSW], 10.6.4 or [NSW<sup>2</sup>], 10.8.4, respectively) and for certain CM fields (by a result of Mukhamedov, see [Muk] or [NSW], X §6 exercise or [NSW<sup>2</sup>], X §8 exercise, respectively).

By Theorem 3.3 below, Theorem 1.2 provides many examples of  $G_S(p)$  being a duality group. If  $\zeta_p \notin k$ , this is interesting even in the case that  $S \supset S_p$ , where examples of  $G_S(p)$  being a duality group were previously known only for real abelian fields and for certain CM-fields (see [NSW], 10.7.15 and [NSW<sup>2</sup>], 10.9.15, respectively, and the remark following there).

Previous results in the mixed case had been achieved by K. Wingberg [Wi1], Ch. Maire [Mai] and D. Vogel [Vog]. Though not explicitly visible in this paper, the present progress in the subject was only possible due to the results on mild pro-p groups obtained by J. Labute in [Lab].

I would like to thank K. Wingberg for pointing out that the proof of Proposition 8.1 in my paper [Sch2] did not use the assumption that the sets S and S' are disjoint from  $S_p$ . This was the key observation for the present paper. The main part of this text was written while I was a guest at the Department of Mathematical Sciences of Tokyo University and of the Research Institute for Mathematical Sciences in Kyoto. I want to thank these institutions for their kind hospitality.

### $\S 2$ . Proof of Theorems 1.1 and 1.2

We start with the observation that the proofs of Proposition 8.1 and Corollary 8.2 in [Sch2] did not use the assumption that the sets S and S' are disjoint from  $S_p$ . Therefore, with the same proof (which we repeat for the convenience of the reader) as in loc. cit., we obtain

**Proposition 2.1.** Let k be a number field and let p be a prime number. Assume k to be totally imaginary if p = 2. Put  $X = Spec(\mathcal{O}_k)$  and let  $S \subset S'$  be finite sets of primes of k. Assume that  $X \setminus S$  is a  $K(\pi, 1)$  for p and that  $G_S(p) \neq 1$ . Further assume that each  $\mathfrak{p} \in S' \setminus S$  does not split completely in  $k_S(p)$ . Then the following hold.

- (i)  $X \smallsetminus S'$  is a  $K(\pi, 1)$  for p.
- (ii)  $k_{S'}(p)_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$  for all  $\mathfrak{p} \in S' \smallsetminus S$ .

Furthermore, the arithmetic form of Riemann's existence theorem holds, i.e., setting  $K = k_S(p)$ , the natural homomorphism

$$*_{\mathfrak{p}\in S'\setminus S(K)} T(K_{\mathfrak{p}}(p)|K_{\mathfrak{p}}) \longrightarrow Gal(k_{S'}(p)|K)$$

is an isomorphism. Here  $T(K_{\mathfrak{p}}(p)|K_{\mathfrak{p}})$  is the inertia group and \* denotes the free prop-product of a bundle of pro-p-groups, cf. [NSW], Ch. IV, §3. In particular, the group  $Gal(k_{S'}(p)|k_S(p))$  is a free pro-p-group.

*Proof.* The  $K(\pi, 1)$ -property implies

$$H^{i}(G_{S}(p), \mathbb{Z}/p\mathbb{Z}) \cong H^{i}_{et}(X \setminus S, \mathbb{Z}/p\mathbb{Z}) = 0 \text{ for } i \geq 4,$$

hence  $cd \ G_S(p) \leq 3$ . Let  $\mathfrak{p} \in S' \setminus S$ . Since  $\mathfrak{p}$  does not split completely in  $k_S(p)$ and since  $cd \ G_S(p) < \infty$ , the decomposition group of  $\mathfrak{p}$  in  $k_S(p)|k$  is a non-trivial and torsion-free quotient of  $\mathbb{Z}_p \cong Gal(k_{\mathfrak{p}}^{nr}(p)|k_{\mathfrak{p}})$ . Therefore  $k_S(p)_{\mathfrak{p}}$  is the maximal unramified *p*-extension of  $k_{\mathfrak{p}}$ . We denote the normalization of an integral normal scheme Y in an algebraic extension L of its function field by  $Y_L$ . Then  $(X \setminus S)_{k_S(p)}$ is the universal pro-*p* covering of  $X \setminus S$ . We consider the étale excision sequence for the pair  $((X \setminus S)_{k_S(p)}, (X \setminus S')_{k_S(p)})$ . By assumption,  $X \setminus S$  is a  $K(\pi, 1)$  for *p*, hence  $H^i_{et}((X \setminus S)_{k_S(p)}, \mathbb{Z}/p\mathbb{Z}) = 0$  for  $i \geq 1$  by [Sch2], Proposition 2.1. Omitting the coefficients  $\mathbb{Z}/p\mathbb{Z}$  from the notation, this implies isomorphisms

$$H^{i}_{et}((X \smallsetminus S')_{k_{S}(p)}) \xrightarrow{\sim} \bigoplus_{\mathfrak{p} \in S' \smallsetminus S(k_{S}(p))}' H^{i+1}_{\mathfrak{p}}(((X \smallsetminus S)_{k_{S}(p)})_{\mathfrak{p}})$$

for  $i \geq 1$ . Here (and in variants also below) we use the notational convention

$$\bigoplus_{\mathfrak{p}\in S'\smallsetminus S(k_S(p))}' H^{i+1}_{\mathfrak{p}}\big(((X\smallsetminus S)_{k_S(p)})_{\mathfrak{p}}\big) := \varinjlim_{K\subset k_S(p)} \bigoplus_{\mathfrak{p}\in S'\smallsetminus S(K)} H^{i+1}_{\mathfrak{p}}\big(((X\smallsetminus S)_K)_{\mathfrak{p}}\big),$$

where K runs through the finite extensions of k inside  $k_S(p)$ . As  $k_S(p)$  realizes the maximal unramified p-extension of  $k_p$  for all  $\mathfrak{p} \in S' \setminus S$ , the schemes  $((X \setminus S)_{k_S(p)})_{\mathfrak{p}}, \mathfrak{p} \in S' \setminus S(k_S(p))$ , have trivial cohomology with values in  $\mathbb{Z}/p\mathbb{Z}$  and we obtain isomorphisms

$$H^{i}((k_{S}(p))_{\mathfrak{p}}) \xrightarrow{\sim} H^{i+1}_{\mathfrak{p}}(((X \smallsetminus S)_{k_{S}(p)})_{\mathfrak{p}})$$

for  $i \ge 1$ . These groups vanish for  $i \ge 2$ . This implies

$$H^i_{et}((X \smallsetminus S')_{k_S(p)}) = 0$$

for  $i \ge 2$ . Since the scheme  $(X \smallsetminus S')_{k_{S'}(p)}$  is the universal pro-*p* covering of  $(X \smallsetminus S')_{k_S(p)}$ , the Hochschild-Serre spectral sequence

$$E_2^{ij} = H^i\big(\operatorname{Gal}(k_{S'}(p)|k_S(p)), H^j_{et}((X \smallsetminus S')_{k_{S'}(p)})\big) \Rightarrow H^{i+j}_{et}((X \smallsetminus S')_{k_S(p)})$$

yields an inclusion

$$H^2(Gal(k_{S'}(p)|k_S(p))) \hookrightarrow H^2_{et}((X \setminus S')_{k_S(p)}) = 0.$$

Hence  $Gal(k_{S'}(p)|k_S(p))$  is a free pro-*p*-group and

$$H^{1}(\operatorname{Gal}(k_{S'}(p)|k_{S}(p))) \xrightarrow{\sim} H^{1}_{et}((X \smallsetminus S')_{k_{S}(p)}) \cong \bigoplus_{\mathfrak{p} \in S' \smallsetminus S(k_{S}(p))}' H^{1}(k_{S}(p)_{\mathfrak{p}}).$$

We set  $K = k_S(p)$  and consider the natural homomorphism

$$\phi: \underset{\mathfrak{p}\in S'\setminus S(K)}{\ast} T(K_{\mathfrak{p}}(p)|K_{\mathfrak{p}}) \longrightarrow Gal(k_{S'}(p)|K).$$

By the calculation of the cohomology of a free product ([NSW], 4.3.10 and 4.1.4),  $\phi$  is a homomorphism between free pro-*p*-groups which induces an isomorphism on mod *p* cohomology. Therefore  $\phi$  is an isomorphism. In particular,  $k_{S'}(p)_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$  for all  $\mathfrak{p} \in S' \setminus S$ . Using that  $Gal(k_{S'}(p)|k_S(p))$  is free, the Hochschild-Serre spectral sequence induces an isomorphism

$$0 = H^2_{et}((X \smallsetminus S')_{k_S(p)}) \xrightarrow{\sim} H^2_{et}((X \smallsetminus S')_{k_{S'}(p)})^{\operatorname{Gal}(k_{S'}(p)|k_S(p))}$$

Hence  $H^2_{et}((X \smallsetminus S')_{k_{S'}(p)}) = 0$ , since  $Gal(k_{S'}(p)|k_S(p))$  is a pro-*p*-group. Now [Sch2], Proposition 2.1 implies that  $X \smallsetminus S'$  is a  $K(\pi, 1)$  for p.

In order to prove Theorem 1.1, we first provide the following lemma. For an extension field K|k and a set of primes T of k, we write T(K) for the set of prolongations of primes in T to K and  $\delta_K(T)$  for the Dirichlet density of the set of primes T(K) of K.

**Lemma 2.2.** Let k be a number field, p a prime number and S a finite set of nonarchimedean primes of k. Let T be a set of primes of k with  $\delta_{k(\mu_p)}(T) = 1$ . Then there exists a finite subset  $T_0 \subset T$  such that all primes  $\mathfrak{p} \in S$  do not split completely in the extension  $k_{T_0}(p)|k$ .

*Proof.* By [NSW], 9.2.2 (ii) or [NSW<sup>2</sup>], 9.2.3 (ii), respectively, the restriction map

$$H^1(G_{T\cup S\cup S_p\cup S_{\mathbb{R}}}(p), \mathbb{Z}/p\mathbb{Z}) \longrightarrow \prod_{\mathfrak{p}\in S\cup S_p\cup S_{\mathbb{R}}} H^1(k_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z})$$

is surjective. A class in  $\alpha \in H^1(G_{T \cup S \cup S_p \cup S_{\mathbb{R}}}(p), \mathbb{Z}/p\mathbb{Z})$  which restricts to an unramified class  $\alpha_{\mathfrak{p}} \in H^1_{nr}(k_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z})$  for all  $\mathfrak{p} \in S \cup S_p \cup S_{\mathbb{R}}$  is contained in  $H^1(G_T(p), \mathbb{Z}/p\mathbb{Z})$ . Therefore the image of the composite map

$$H^1(G_T(p), \mathbb{Z}/p\mathbb{Z}) \hookrightarrow H^1(G_{T \cup S \cup S_p \cup S_{\mathbb{R}}}(p), \mathbb{Z}/p\mathbb{Z}) \to \prod_{\mathfrak{p} \in S} H^1(k_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z})$$

contains the subgroup  $\prod_{\mathfrak{p}\in S} H^1_{nr}(k_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z})$ . As this group is finite, it is already contained in the image of  $H^1(G_{T_0}(p), \mathbb{Z}/p\mathbb{Z})$  for some finite subset  $T_0 \subset T$ . We conclude that no prime in S splits completely in the maximal elementary abelian p-extension of k unramified outside  $T_0$ .

Proof of Theorems 1.1 and 1.2. As  $p \neq 2$ , we may ignore archimedean primes. Furthermore, we may remove the primes in  $S \cup S_p$  and all primes of degree greater than 1 from T. In addition, we remove all primes  $\mathfrak{p}$  with  $N(\mathfrak{p}) \not\equiv 1 \mod p$  from T. After these changes, we still have  $\delta_{k(\mu_p)}(T) = 1$ .

By Lemma 2.2, we find a finite subset  $T_0 \subset T$  such that no prime in S splits completely in  $k_{T_0}(p)|k$ . Put  $X = Spec(\mathcal{O}_k)$ . By [Sch2], Theorem 6.2, applied to  $T_0$  and  $T \setminus T_0$ , we find a finite subset  $T_2 \subset T \setminus T_0$  such that  $X \setminus (T_0 \cup T_2)$  is a  $K(\pi, 1)$  for p. Then Proposition 2.1 applied to  $T_0 \cup T_2 \subset S \cup T_0 \cup T_2$ , shows that also  $X \setminus (S \cup T_0 \cup T_2)$ is a  $K(\pi, 1)$  for p. Now put  $T_1 = T_0 \cup T_2 \subset T$ .

It remains to show Theorem 1.2. Assertion (i) holds by construction of  $T_1$ . Again by construction,  $X \setminus T_1$  is a  $K(\pi, 1)$  for p. By [Sch2], Theorem 3, the field  $k_{T_1}(p)$  realizes  $k_{\mathfrak{p}}(p)$  for  $\mathfrak{p} \in T_1$ , showing (ii) for these primes. Finally, assertion (ii) for  $\mathfrak{p} \in S$  follows from Proposition 2.1.

# §3. Duality

We start by investigating the relation between the  $K(\pi, 1)$ -property and the universal norms of global units.

Let us first remove redundant primes from S: If  $\mathfrak{p} \nmid p$  is a prime with  $\zeta_p \notin k_{\mathfrak{p}}$ , then every *p*-extension of the local field  $k_{\mathfrak{p}}$  is unramified (see [NSW], 7.5.1 or [NSW<sup>2</sup>], 7.5.9, respectively). Therefore primes  $\mathfrak{p} \notin S_p$  with  $N(\mathfrak{p}) \not\equiv 1 \mod p$  cannot ramify in a *p*-extension. Removing all these redundant primes from S, we obtain a subset  $S_{\min} \subset S$ , which has the property that  $G_S(p) = G_{S_{\min}}(p)$ . Furthermore, by [Sch2], Lemma 4.1,  $X \setminus S$  is a  $K(\pi, 1)$  for p if and only if  $X \setminus S_{\min}$  is a  $K(\pi, 1)$  for p.

**Theorem 3.1.** Let k be a number field and let p be a prime number. Assume that k is totally imaginary if p = 2. Let S be a finite set of nonarchimedean primes of k. Then any two of the following conditions (a) – (c) imply the third.

- (a)  $Spec(\mathcal{O}_k) \smallsetminus S$  is a  $K(\pi, 1)$  for p.
- (b)  $\varprojlim_{K \subset k_S(p)} \mathcal{O}_K^{\times} \otimes \mathbb{Z}_p = 0.$
- (c)  $(k_S(p))_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$  for all primes  $\mathfrak{p} \in S_{\min}$ .

The limit in (b) runs through all finite extensions K of k inside  $k_S(p)$ . If (a)–(c) hold, then also

$$\lim_{K \subset k_S(p)} \mathcal{O}_{K,S_{\min}}^{\times} \otimes \mathbb{Z}_p = 0.$$

**Remarks:** 1. Assume that  $\zeta_p \in k$  and  $S \supset S_p$ . Then (a) holds and condition (c) holds for p > 2 if  $\#S > r_2 + 2$  (see [NSW<sup>2</sup>], Remark 2 after 10.9.3). In the case  $k = \mathbb{Q}(\zeta_p)$ ,  $S = S_p$ , condition (c) holds if and only if p is an irregular prime number.

2. Assume that  $S \cap S_p = \emptyset$  and  $S_{\min} \neq \emptyset$ . If condition (a) holds, then either  $G_S(p) = 1$  (which only happens in very special situations, see [Sch2], Proposition 7.4) or (c) holds by [Sch2], Theorem 3 (or by Proposition 3.2 below).

Proof of Theorem 3.1. We may assume  $S = S_{\min}$  in the proof. Let K run through the finite extensions of k in  $k_S(p)$  and put  $X_K = Spec(\mathcal{O}_K)$ . Applying the topological Nakayama-Lemma ([NSW], 5.2.18) to the compact  $\mathbb{Z}_p$ -module  $\varprojlim \mathcal{O}_K^{\times} \otimes \mathbb{Z}_p$ , we see that condition (b) is equivalent to

(b)' 
$$\varprojlim_{K \subset k_S(p)} \mathcal{O}_K^{\times}/p = 0.$$

Furthermore, by [Sch2], Proposition 2.1, condition (a) is equivalent to

(a)' 
$$\varinjlim_{K \subset k_S(p)} H^i_{et}((X \setminus S)_K, \mathbb{Z}/p\mathbb{Z}) = 0 \text{ for } i \ge 1.$$

Condition (a)' always holds for  $i = 1, i \ge 4$ , and it holds for i = 3 provided that  $G_S(p)$  is infinite or S is nonempty or  $\zeta_p \notin k$  (see [Sch2], Lemma 3.7). The flat Kummer sequence  $0 \to \mu_p \to \mathbb{G}_m \xrightarrow{\cdot p} \mathbb{G}_m \to 0$  induces exact sequences

$$0 \longrightarrow \mathcal{O}_K^{\times}/p \longrightarrow H^1_{fl}(X_K, \mu_p) \longrightarrow {}_p Pic(X_K) \to 0$$

for all K. As the field  $k_S(p)$  does not have nontrivial unramified p-extensions, class field theory implies

$$\lim_{K \subset k_S(p)} {}_p Pic(X_K) \subset \lim_{K \subset k_S(p)} Pic(X_K) \otimes \mathbb{Z}_p = 0.$$

As we assumed k to be totally imaginary if p = 2, the flat duality theorem of Artin-Mazur ([Mil], III Corollary 3.2) induces natural isomorphisms

$$H^2_{et}(X_K, \mathbb{Z}/p\mathbb{Z}) = H^2_{ft}(X_K, \mathbb{Z}/p\mathbb{Z}) \cong H^1_{ft}(X_K, \mu_p)^{\vee}.$$

We conclude that

(\*) 
$$\lim_{K \subset k_S(p)} H^2_{et}(X_K, \mathbb{Z}/p\mathbb{Z}) \cong \Big(\lim_{K \subset k_S(p)} \mathcal{O}_K^{\times}/p\Big)^{\vee}.$$

We first show the equivalence of (a) and (b) in the case  $S = \emptyset$ . If (a)' holds, then (\*) shows (b)'. If (b) holds, then  $\zeta_{\mathfrak{p}} \notin k$  or  $G_S(p)$  is infinite. Hence we obtain (a)' for i = 3. Furthemore, (b)' implies (a)' for i = 2 by (\*). This finishes the proof of the case  $S = \emptyset$ .

Now we assume that  $S \neq \emptyset$ . For  $\mathfrak{p} \in S(K)$ , a standard calculation of local cohomology shows that

$$H^{i}_{\mathfrak{p}}(X_{K}, \mathbb{Z}/p\mathbb{Z}) \cong \begin{cases} 0 & \text{for } i \leq 1, \\ H^{1}(K_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z})/H^{1}_{nr}(K_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z}) & \text{for } i = 2, \\ H^{2}(K_{\mathfrak{p}}, \mathbb{Z}/p\mathbb{Z}) & \text{for } i = 3. \\ 0 & \text{for } i \geq 4. \end{cases}$$

For  $\mathfrak{p} \in S = S_{\min}$ , every proper Galois subextension of  $k_{\mathfrak{p}}(p)|k_{\mathfrak{p}}$  admits ramified *p*-extensions. Hence condition (c) is equivalent to

(c)' 
$$\varinjlim_{K \subset k_S(p)} \bigoplus_{\mathfrak{p} \in S(K)} H^i_{\mathfrak{p}}(X_K, \mathbb{Z}/p\mathbb{Z}) = 0 \text{ for all } i,$$

and to

(c)" 
$$\varinjlim_{K \subset k_S(p)} \bigoplus_{\mathfrak{p} \in S(K)} H^2_{\mathfrak{p}}(X_K, \mathbb{Z}/p\mathbb{Z}) = 0$$

Consider the direct limit over all K of the excision sequences

$$\cdots \to \bigoplus_{\mathfrak{p} \in S(K)} H^i_{\mathfrak{p}}(X_K, \mathbb{Z}/p\mathbb{Z}) \to H^i_{et}(X_K, \mathbb{Z}/p\mathbb{Z}) \to H^i_{et}((X \setminus S)_K, \mathbb{Z}/p\mathbb{Z}) \to \cdots$$

Assume that (a)' holds, i.e. the right hand terms vanish in the limit for  $i \ge 1$ . Then (\*) shows that (b)' is equivalent to (c)".

Now assume that (b) and (c) hold. As above, (b) implies the vanishing of the middle term for i = 2 in the limit. Condition (c)' then shows (a)'.

We have proven that any two of the conditions (a)-(c) imply the third.

#### Alexander Schmidt

Finally, assume that (a)–(c) hold. Tensoring the exact sequences (cf. [NSW], 10.3.11 or  $[NSW^2]$ , 10.3.12, respectively)

$$0 \to \mathcal{O}_K^{\times} \to \mathcal{O}_{K,S}^{\times} \to \bigoplus_{\mathfrak{p} \in S(K)} (K_\mathfrak{p}^{\times}/U_\mathfrak{p}) \to Pic(X_K) \to Pic((X \smallsetminus S)_K) \to 0$$

by (the flat Z-algebra)  $\mathbb{Z}_p$ , we obtain exact sequences of finitely generated, hence compact,  $\mathbb{Z}_p$ -modules. Passing to the projective limit over the finite extensions K of k inside  $k_S(p)$  and using  $\varprojlim Pic(X_K) \otimes \mathbb{Z}_p = 0$ , we obtain the exact sequence

$$0 \to \varprojlim_{K \subset k_S(p)} \mathcal{O}_K^{\times} \otimes \mathbb{Z}_p \to \varprojlim_{K \subset k_S(p)} \mathcal{O}_{K,S}^{\times} \otimes \mathbb{Z}_p \to \varprojlim_{K \subset k_S(p)} \bigoplus_{\mathfrak{p} \in S(K)} (K_\mathfrak{p}^{\times}/U_\mathfrak{p}) \otimes \mathbb{Z}_p \to 0.$$

Condition (c) and local class field theory imply the vanishing of the right hand limit. Therefore (b) implies the vanishing of the projective limit in the middle.  $\Box$ 

If  $G_S(p) \neq 1$  and condition (a) of Theorem 1.1 holds, then the failure in condition (c) can only come from primes dividing p. This follows from the next

**Proposition 3.2.** Let k be a number field and let p be a prime number. Assume that k is totally imaginary if p = 2. Let S be a finite set of nonarchimedean primes of k. If  $Spec(\mathcal{O}_k) \setminus S$  is a  $K(\pi, 1)$  for p and  $G_S(p) \neq 1$ , then every prime  $\mathfrak{p} \in S$  with  $\zeta_p \in k_{\mathfrak{p}}$  has an infinite inertia group in  $G_S(p)$ . Moreover, we have

$$k_S(p)_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$$

for all  $\mathfrak{p} \in S_{\min} \setminus S_p$ .

*Proof.* We may assume  $S = S_{\min}$ . Suppose  $\mathfrak{p} \in S$  with  $\zeta_p \in k_{\mathfrak{p}}$  does not ramify in  $k_S(p)|k$ . Setting  $S' = S \setminus \{\mathfrak{p}\}$ , we have  $k_{S'}(p) = k_S(p)$ , in particular,

$$H^1_{et}(X \smallsetminus S', \mathbb{Z}/p\mathbb{Z}) \xrightarrow{\sim} H^1_{et}(X \smallsetminus S, \mathbb{Z}/p\mathbb{Z}).$$

In the following, we omit the coefficients  $\mathbb{Z}/p\mathbb{Z}$  from the notation. Using the vanishing of  $H^3_{et}(X \setminus S)$ , the étale excision sequence yields a commutative exact diagram

$$H^{2}(G_{S'}(p)) \xrightarrow{\sim} H^{2}(G_{S}(p))$$

$$\downarrow^{\wr}$$

$$H^{2}_{\mathfrak{p}}(X) \xrightarrow{\sim} H^{2}_{et}(X \smallsetminus S') \xrightarrow{\alpha} H^{2}_{et}(X \smallsetminus S) \longrightarrow H^{3}_{\mathfrak{p}}(X) \longrightarrow H^{3}_{et}(X \smallsetminus S')$$

Hence  $\alpha$  is split-surjective and  $\mathbb{Z}/p\mathbb{Z} \cong H^3_{\mathfrak{p}}(X) \xrightarrow{\sim} H^3_{et}(X \setminus S')$ . This implies  $S' = \emptyset$ , hence  $S = \{\mathfrak{p}\}$ , and  $\zeta_p \in k$ . The same applies to every finite extension of k in  $k_S(p)$ , hence  $\mathfrak{p}$  is inert in  $k_S(p) = k_{\emptyset}(p)$ . This implies that the natural homomorphism

$$Gal(k_{\mathfrak{p}}^{nr}(p)|k_{\mathfrak{p}}) \longrightarrow G_{\varnothing}(k)(p)$$

is surjective. Therefore  $G_S(p) = G_{\emptyset}(p)$  is abelian, hence finite by class field theory. Since this group has finite cohomological dimension by the  $K(\pi, 1)$ -property, it is trivial, in contradiction to our assumptions.

This shows that all  $\mathfrak{p} \in S$  with  $\zeta_p \in k_\mathfrak{p}$  ramify in  $k_S(p)$ . As this applies to every finite extension of k inside  $k_S(p)$ , the inertia groups must be infinite. For  $\mathfrak{p} \in S_{\min} \setminus S_p$  this implies  $k_S(p)_\mathfrak{p} = k_\mathfrak{p}(p)$ .

**Theorem 3.3.** Let k be a number field and let p be a prime number. Assume that k is totally imaginary if p = 2. Let S be a finite nonempty set of nonarchimedean primes of k. Assume that conditions (a)–(c) of Theorem 3.1 hold and that  $\zeta_p \in k_p$  for all  $p \in S$ . Then  $G_S(p)$  is a pro-p duality group of dimension 2.

*Proof.* Condition (a) implies  $H^3(G_S(p), \mathbb{Z}/p\mathbb{Z}) \xrightarrow{\sim} H^3_{et}(X \setminus S, \mathbb{Z}/p\mathbb{Z}) = 0$ . Hence *cd*  $G_S(p) \leq 2$ . On the other hand, by (c), the group  $G_S(p)$  contains  $Gal(k_{\mathfrak{p}}(p)|k_{\mathfrak{p}})$  as a subgroup for all  $\mathfrak{p} \in S$ . As  $\zeta_p \in k_{\mathfrak{p}}$  for  $\mathfrak{p} \in S$ , these local groups have cohomological dimension 2, hence so does  $G_S(p)$ .

In order to show that  $G_S(p)$  is a duality group, we have to show that

$$D_i(G_S(p), \mathbb{Z}/p\mathbb{Z}) := \varinjlim_{\substack{U \subset G_S(p)\\cor^{\vee}}} H^i(U, \mathbb{Z}/p\mathbb{Z})^{\vee}$$

vanish for i = 0, 1, where U runs through the open subgroups of  $G_S(p)$  and the transition maps are the duals of the corestriction homomorphisms; see [NSW], 3.4.6. The vanishing of  $D_0$  is obvious, as  $G_S(p)$  is infinite. We therefore have to show that

$$\lim_{K \subset k_S(p)} H^1((X \setminus S)_K, \mathbb{Z}/p\mathbb{Z})^{\vee} = 0.$$

We put  $X = Spec(\mathcal{O}_k)$  and denote the embedding by  $j : (X \setminus S)_K \to X_K$ . By the flat duality theorem of Artin-Mazur, we have natural isomorphisms

$$H^1((X \setminus S)_K, \mathbb{Z}/p\mathbb{Z})^{\vee} \cong H^2_{fl,c}((X \setminus S)_K, \mu_p) = H^2_{fl}(X_K, j_!\mu_p).$$

The excision sequence together with a straightforward calculation of local cohomology groups shows an exact sequence

(\*) 
$$\bigoplus_{\mathfrak{p}\in S(K)} K_{\mathfrak{p}}^{\times}/K_{\mathfrak{p}}^{\times p} \to H_{fl}^2(X_K, j_!\mu_p) \to H_{fl}^2((X \smallsetminus S)_K, \mu_p).$$

As  $\zeta_p \in k_{\mathfrak{p}}$  and  $k_S(p)_{\mathfrak{p}} = k_{\mathfrak{p}}(p)$  for  $\mathfrak{p} \in S$  by assumption, the left hand term of (\*) vanishes when passing to the limit over all K. We use the Kummer sequence to obtain an exact sequence

$$(**) \qquad Pic((X \smallsetminus S)_K)/p \longrightarrow H^2_{fl}((X \smallsetminus S)_K, \mu_p) \longrightarrow {}_pBr((X \smallsetminus S)_K).$$

The left hand term of (\*\*) vanishes in the limit by the principal ideal theorem. The Hasse principle for the Brauer group induces an injection

$$_{p}Br((X \smallsetminus S)_{K}) \hookrightarrow \bigoplus_{\mathfrak{p} \in S(K)} {_{p}Br(K_{\mathfrak{p}})}.$$

As  $k_S(p)$  realizes the maximal unramified *p*-extension of  $k_p$  for  $p \in S$ , the limit of the middle term in (\*\*), and hence also the limit of the middle term in (\*) vanishes. This shows that  $G_S(p)$  is a duality group of dimension 2.

**Remark:** The dualizing module can be calculated to

$$D \cong \operatorname{tor}_p(C_S(k_S(p))),$$

i.e. D is isomorphic to the p-torsion subgroup in the S-idèle class group of  $k_S(p)$ . The proof is the same as in ([Sch1], Proof of Thm. 5.2), where we dealt with the tame case.

#### References

- [Kuz] L. V. Kuz'min Local extensions associated with l-extensions with given ramification (in Russian). Izv. Akad. Nauk SSSR 39 (1975) no. 4, 739–772. English transl. in Math. USSR Izv. 9 (1975), no. 4, 693–726.
- [Lab] J. P. Labute Mild pro-p-groups and Galois groups of p-extensions of Q. J. Reine und angew. Math. 596 (2006), 155–182.
- [NSW] J. Neukirch, A. Schmidt, K. Wingberg, Cohomology of Number Fields, Grundlehren der math. Wiss. Bd. 323, Springer-Verlag 2000.
- [NSW<sup>2</sup>] J. Neukirch, A. Schmidt, K. Wingberg, *Cohomology of Number Fields, 2nd ed.*, Grundlehren der math. Wiss. Bd. 323, Springer-Verlag 2008.
  - [Maz] B. Mazur Notes on étale cohomology of number fields. Ann. Sci. École Norm. Sup. (4)
     6 (1973), 521–552.
  - [Mai] Ch. Maire Sur la dimension cohomologique des pro-p-extensions des corps de nombres.
     J. Théor. Nombres Bordeaux 17 (2005), no. 2, 575–606.
  - [Mil] J.S. Milne Arithmetic duality theorems. Academic Press 1986.
  - [Muk] V. G. Mukhamedov, Local extensions associated with the l-extensions of number fields with restricted ramification (in Russian). Mat. Zametki 35 (1984), no. 4, 481–490, English transl. in Math. Notes 35, no. 3–4, 253–258.
  - [Sch1] A. Schmidt Circular sets of prime numbers and p-extension of the rationals. J. Reine und angew. Math. 596 (2006), 115–130.
  - [Sch2] A. Schmidt Rings of integers of type  $K(\pi, 1)$ . Doc. Math. 12 (2007), 441–471.
  - [Vog] D. Vogel p-extensions with restricted ramification the mixed case. Preprints der Forschergruppe Algebraische Zykel und L-Funktionen Regensburg/Freiburg Nr. 11, 2007: http://www.mathematik.uni-regensburg.de/FGAlgZyk/.
  - [Wi1] K. Wingberg Galois groups of number fields generated by torsion points of elliptic curves. Nagoya Math. J. 104 (1986), 43–53.
  - [Wi2] K. Wingberg Riemann's existence theorem and the  $K(\pi, 1)$ -property of rings of integers. Preprint 2007.