

Invariants of Unitary Reflection Groups

寺尾 宏明
(国際基督教大学(ICU))


有限ユニタリ鏡映群は, Shephard-Todd [3] によ, 2
分類され, 1~37 までの名前が与えられている。本
稿では, それらの不変量の表を作り, そこから得られる
observation を紹介したい。

§ 1. Notations and Table

1. The names of finite unitary reflection groups are due to Shephard-Todd [3],
2. m_1, \dots, m_ℓ are the exponents of group (i.e., the degrees of basis of invariant polynomials -1); ($m_1 \leq m_2 \leq \dots \leq m_\ell$)
 n_1, \dots, n_ℓ are the exponents of arrangement (i.e., the degrees of basis of logarithmic derivations $+1$). (See [4]) ($n_1 \leq n_2 \leq \dots \leq n_\ell$)
3. $m = \sum_{i=1}^{\ell} m_i$, $n = \sum_{i=1}^{\ell} n_i$.

4. $r_i = \#\{\text{reflections of order } i\}$,
 $s_i = \#\{\text{reflecting hyperplane } H \text{ such that}$
 there exist exactly i
 fixing H pointwise}\}
5. (d_1, \dots, d_ℓ) is self-dual $\stackrel{\text{def}}{\iff} d_i + d_{\ell+1-i} = \text{const.}$
 $(1 \leq i \leq \ell)$
6. (m_1, \dots, m_ℓ) and (n_1, \dots, n_ℓ) are codual
 $\stackrel{\text{def}}{\iff} m_i + n_{\ell+1-i} = \text{const.}$ $(1 \leq i \leq \ell)$
7. $t := \frac{2n}{\ell}$, $H := \frac{2(m+n)}{2\ell + (m-n)}$

8. type:



9. Condition (A): Are (m_i) & (n_i) both self-dual?
 Condition (B): If yes, $m_1 + m_\ell = n_1 + n_\ell$?
 Condition (C): Are (m_i) & (n_i) codual?
 Condition (D): Is G generated by ℓ reflections?
 Condition (E): Is G generated by reflections of order 2?

Group	m_1, \dots, m_r	m	r_2, r_3, \dots	t	m_{r+1}	(A)	(B)	(C)	(D)	(E)	type
Γ (A ₀)	m_1, \dots, m_r	n	S_2, S_3, \dots	H	m_{r+1}	(A)	(B)	(C)	(D)	(E)	Coxeter
Γ	$1, 2, \dots, r$	$\frac{r}{2} (r+1)$	$\frac{r}{2} (r+1), 0, 0, \dots$	$r+1$	$r+1$	Y	Y	Y	Y	Y	Coxeter
Γ	$1, 2, \dots, r$	$\frac{r}{2} (r+1)$	$\frac{r}{2} (r+1), 0, 0, \dots$	$r+1$	$r+1$	Y	Y	Y	Y	Y	Coxeter
$G(r, p, l)$											
$r = p^2$											
$r = q = 2$ (B ₂)	$1, 3, 5, \dots, 2l-1$	l^2	$l^2, 0, 0, \dots$	$2l$	$2l$	Y	Y	Y	Y	Y	Coxeter
$r = q = 2$	$1, 3, 5, \dots, 2l-1$	l^2	$l^2, 0, 0, \dots$	$2l$	$2l$	Y	Y	Y	Y	Y	Coxeter
$r = 4$	$3, 7, \dots, 4l-5 (=m_2), 2l-1$	$l(2l-1)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-4$	Y	Y	n	n	Y	II
$r = q = 2$	$1, 5, \dots, 4l-3$	$l(2l-1)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	Y	n	n	Y	II
$r = q > 2$	$r-1, 2r-1, \dots, lr-1$	$l \left(\frac{(l+1)r}{2} - 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-4$	Y	Y	n	n	Y	II
$r = q > 2$	$1, r+1, \dots, (l-1)r+1$	$l \left(\frac{(l-1)r}{2} + 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	Y	Y	n	III
$r = 2q > 4$	$r-1, 2r-1, \dots, (l-1)r-1, \frac{lr}{2}-1$	$l \left(\frac{(l-1)r}{2} - 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	n	n	n	IV
$r = 2q > 4$	$1, r+1, \dots, (l-1)r+1$	$l \left(\frac{(l-1)r}{2} + 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	n	n	n	IV
$q \neq r \neq 2q$	$r-1, 2r-1, \dots, (l-1)r-1, lq-1$	$l \left(\frac{(l-1)r}{2} + q - 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$q > 2$	$1, r+1, \dots, (l-1)r+1$	$l \left(\frac{(l-1)r}{2} + 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$q \neq r \neq 2q$	$r-1, 2r-1, \dots, (l-1)r-1, 2l-1$	$l \left(\frac{(l-1)r}{2} + 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$q = 2$	$1, r+1, \dots, (l-1)r+1$	$l \left(\frac{(l-1)r}{2} + 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$q \neq r \neq 2q$	$r-1, 2r-1, \dots, (l-1)r-1, l-1$	$l \left(\frac{(l-1)r}{2} - 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$q > 2$	$1, r+1, \dots, (l-2)r+1, (l-1)r-1$	$l \left(\frac{(l-1)r}{2} - 1 \right)$	$l(2l-1), 0, 0, \dots$	$4l-2$	$4l-2$	Y	n	-	n	n	IV
$p = r = 2$ (D ₂)	$1, 3, 5, \dots, 2l-3, l-1$	$l(l-1)$	$l(l-1), 0, 0, \dots$	$2l$	$2l$	Y	Y	Y	Y	Y	Coxeter
$p = r = 2$ (D ₂)	$1, 3, 5, \dots, 2l-3, l-1$	$l(l-1)$	$l(l-1), 0, 0, \dots$	$2l$	$2l$	Y	Y	Y	Y	Y	Coxeter

Group	m_1, \dots, m_r	m	k_2, k_3, \dots s_2, s_3, \dots	t	m_{r+1}	(A)	(B)	(C)	(D)	(E)	type
3	$r-1$	$r-1$	$0, 0, 0, 1, 0, \dots$	2	r	y	n	y	y	n	III
$r > 2$	$r-1$	1		2	r	y	n	y	y	n	III
4	$3, 5$ $1, 3$	8	$0, 8, 0, \dots$ $0, 4, 0, \dots$	4	6	y	n	y	y	n	III
5	$5, 11$ $1, 7$	16	$0, 16, 0, \dots$ $0, 8, 0, \dots$	8	12	y	n	y	y	n	III
6	$3, 11$ $1, 9$	14	$6, 8, 0, \dots$ $6, 4, 0, \dots$	10	12	y	n	y	y	n	III
7	$11, 11$ $1, 13$	22	$6, 16, 0, \dots$ $6, 8, 0, \dots$	14	12	y	n	n	n	n	IV
8	$7, 11$ $1, 5$	18	$6, 0, 12, \dots$ $0, 0, 6, \dots$	6	12	y	n	y	y	n	III
9	$7, 23$ $1, 17$	30	$18, 0, 12, \dots$ $12, 0, 6, \dots$	18	24	y	n	y	y	n	III
10	$11, 23$ $1, 13$	34	$6, 16, 12, \dots$ $0, 8, 6, \dots$	14	24	y	n	y	y	n	III

1
4

Group	m_1, \dots, m_r	m	K_2, K_3, \dots S_2, S_3, \dots	\pm	m_{e+1} m_{e+1}	(A) (B) (C) (D) (E) type
11	23, 23 1, 25	46 26	18, 16, 12, ... 12, 8, 6, ...	26 6	24 26	Y N N N N IV
12	5, 7 1, 11	12	12, 0, ... 12, 0, ...	12	8 12	Y Y N N Y II
13	7, 11 1, 17	18	18, 0, ... 18, 0, ...	18	12 18	Y Y N N Y II
14	5, 23 1, 19	28 20	12, 16, ... 12, 8, ...	20	24 20	Y N Y Y N III
15	11, 23 1, 25	34 26	18, 16, ... 18, 8, ...	26	24 26	Y N N N N IV
16	19, 29 1, 11	48 12	0, 0, 0, 48 0, 0, 0, 12	12	30 12	Y N Y Y N III
17	19, 59 1, 41	78 42	30, 0, 0, 48 30, 0, 0, 12	42	60 42	Y N Y Y N III
18	29, 59 1, 31	88 32	0, 40, 0, 48 0, 20, 0, 12	32	60 32	Y N Y Y N III
19	59, 59 1, 61	118 62	30, 40, 0, 48 30, 20, 0, 12	62	60 62	Y N N N N IV
20	11, 29 1, 19	40 20	0, 40, ... 0, 20, ...	20	30 20	Y N Y Y N III

15

Group	m_1, \dots, m_r	m	k_2, k_3, \dots	T	m_{r+1}	(A)	(B)	(C)	(D)	(E)	type
21	1, 5, 9 1, 4, 9	70 50	30, 40, ... 30, 20, ...	50 10	60 50	Y	m	Y	Y	m	III
22	1, 1, 1, 9 1, 2, 9	30 30	30, 0, ... 30, 0, ...	30 30	20 30	Y	Y	m	m	Y	II
23 (CH ₃)	1, 5, 9 1, 5, 9	15 15	15, 0, ... 15, 0, ...	10 10	10 10	Y	Y	Y	Y	Y	I
24	3, 5, 13 1, 9, 11	21 21	21, 0, ... 21, 0, ...	14 14	14 12	m	-	Y	Y	Y	I
25	5, 8, 11 1, 4, 7	24 12	0, 24, ... 0, 12, ...	8 4	12 8	Y	m	Y	Y	m	III
26	5, 11, 17 1, 7, 13	33 21	9, 24, ... 9, 12, ...	14 6	18 14	Y	m	Y	Y	m	III
27	5, 11, 29 1, 19, 25	45 45	45, 0, ... 45, 0, ...	30 30	30 26	m	-	Y	Y	Y	I
28 (F ₄)	1, 5, 7, 11 1, 5, 7, 11	24 24	24, 0, ... 24, 0, ...	12 12	12 12	Y	Y	Y	Y	Y	Complex
29	3, 7, 11, 19 1, 9, 13, 17	40 40	40, 0, ... 40, 0, ...	20 20	20 18	m	-	Y	Y	Y	I
30 (CH ₄)	1, 1, 1, 19, 29 1, 1, 1, 19, 29	60 60	60, 0, ... 60, 0, ...	30 30	30 30	Y	Y	Y	Y	Y	Complex

Groups	m_1, \dots, m_r	m	V_2, V_3, \dots	t	m_{r+1}	(A)	(B)	(C)	(D)	(E)	type
31	7, 11, 19, 23 1, 13, 17, 29	60	60, 0, ... 60, 0, ...	30	34	Y	Y	m	m	Y	II
32	11, 17, 23, 29 1, 7, 13, 19	80	0, 80, ... 0, 40, ...	20	30	Y	m	Y	Y	m	III
33	3, 5, 9, 11, 17 1, 7, 9, 13, 15	45	45, 0, ... 45, 0, ...	18	18	m	-	Y	Y	Y	I
34	5, 11, 17, 23, 29, 41 1, 13, 19, 25, 31, 37	126	126, 0, ... 126, 0, ...	42	38	m	-	Y	Y	Y	I
35 (E ₆)	1, 4, 5, 7, 8, 11 1, 4, 5, 7, 8, 11	36	36, 0, ... 36, 0, ...	12	12	Y	Y	Y	Y	Y	Correct
36 (E ₇)	1, 5, 7, 9, 11, 13, 17 1, 5, 7, 9, 11, 13, 17	63	63, 0, ... 63, 0, ...	18	18	Y	Y	Y	Y	Y	Correct
37 (E ₈)	1, 7, 11, 13, 17, 19, 23, 29 1, 7, 11, 13, 17, 19, 23, 29	120	120, 0, ... 120, 0, ...	30	30	Y	Y	Y	Y	Y	Correct

§ 2. Observations

1. (Orlik-Solomon [1])

(m) & (n) are codual (condition (C))

$$\begin{array}{c} \Downarrow \\ G \text{ is generated by } l \text{ reflections (condition (D))} \\ \Downarrow \end{array}$$

$$m_e \geq n_e$$

2. H is always a natural number. (This invariant was introduced by T. Yano.)
 t has been proved to be a natural number [5].

3. There are five cases:

$$t=H=m_e+1=n_e+1 \Leftrightarrow \begin{array}{ccccc} & (A) & (B) & (C) & (D) & (E) \\ \gamma & \gamma & \gamma & \gamma & \gamma & \end{array} \Leftrightarrow \text{Coxeter}$$

$$t=H=m_e+1 > n_e+1 \Leftrightarrow n \quad - \quad \gamma \quad \gamma \quad \gamma \Leftrightarrow \text{I}$$

$$t=H=n_e+1 > m_e+1 \Leftrightarrow \gamma \quad \gamma \quad n \quad n \quad \gamma \Leftrightarrow \text{II}$$

$$m_e+1 > n_e+1 = t > H \Leftrightarrow \gamma \quad n \quad \gamma \quad \gamma \quad n \Leftrightarrow \text{III}$$

$$n_e+1 = t > m_e+1 > H \Leftrightarrow \gamma \quad n \quad n \quad n \quad n \Leftrightarrow \text{IV}$$

(with only exception: 2)

4. III \Leftrightarrow Shephard group (Orlik-Solomon [2])

References

1. P. Orlik-Solomon, Unitary reflection groups and cohomology, *Invent. math.* 59 (1980), 77-94.
2. P. Orlik-Solomon, The Hessian map in the invariant theory of reflection groups, *Nagoya Math. J.* 109 (1988), 1-21.
3. G.C. Shephard-J.A. Todd, Finite unitary reflection groups, *Canad. J. Math.* 6 (1954), 274-304.
4. H. Terao, Free arrangements of hyperplanes and unitary reflection groups, *Proc. Japan Acad. Ser. A* 56 (1980), 389-392.
5. H. Terao, On a certain invariant of a finite unitary reflection group, *Proc. Japan Acad. Ser. A* 65 (1989), 320-322.