

ACTIONS OF COUNTABLE DISCRETE GROUPS ON THE FERMIONIC FACTOR

東北大理 齋藤和之(Kazuyuki SAITÔ)

The main theorem of Sullivan-Weiss-Wright shows that when A is the Dixmier algebra, if G_1 and G_2 are arbitrary countably infinite discrete groups and if $\alpha^{G_1}, \alpha^{G_2}$ are, respectively, outer, ergodic actions on A as $*$ -automorphism groups, then, (A, G_1, α^{G_1}) is always weakly equivalent to (A, G_2, α^{G_2}) (see below Definition 2), where the Dixmier algebra is the C^* -algebra of all bounded complex Borel functions on the unit interval $[0,1]$, modulo the ideal of functions with meagre supports (This is the regular completion of the C^* -algebra $C[0,1]$. See [1]).

The idea of the proof of this result is to construct an action of the dyadic group $\bigoplus \mathbb{Z}_2$ directly, by gluing together parts of homeomorphisms in G_1 and, at the same time, split the base space into dyadic pieces which can be mapped into the Cantor set in a natural way (see [11]). This means that the classifications modulo meagre sets are very rough. So it seems reasonable to conjecture that an analogous result is true when $A = \widehat{F}$, the regular completion of the Fermion algebra F .

In this report, we would like to give you a negative answer to this conjecture by a concrete construction of an outer, ergodic action β of each countably infinite discrete group G on \widehat{F} so that β^{G_1} is weakly equivalent to β^{G_2} if, and only if, G_1 is isomorphic to G_2 .

This is a joint work with John D.M. Wright [8], [9] and [10].

We recall that the regular completion \widehat{B} of a separable unital C^* -algebra B is the unique monotone complete C^* -algebra in

which B is canonically embedded in such a way that

(1) \hat{B} is monotone generated by B ,

(2) for each self-adjoint element b in \hat{B} , the set

$$(-\infty, b] \cap B_h = \{x \in B_h \mid x \leq b\}$$

has b as its supremum in \hat{B}_h , where C_h is the set of all self-adjoint elements of a $*$ -algebra C .

In fact, let \mathcal{B}_B be the Borel envelope of B and let \mathcal{M}_B be the set of elements in \mathcal{B}_B such that

$$\{\phi \in \partial_e X_B \mid m(\phi) \neq 0\}$$

is a meagre subset of $\partial_e X_B$, where $\partial_e X_B$ is the Baire space of all pure states of B relative to the weak*-topology. Then $\mathcal{B}_B / \mathcal{M}_B$ meets all the requirements of the properties of \hat{B} . When B is simple and infinite dimensional, then \hat{B} is a wild AW*-factor of Type III. In particular, \hat{F} is a monotone complete, wild factor of Type III. (See [14] and [15]. See also [2]). We call this \hat{F} , temporarily, a Fermionic factor.

The Fermion algebra F may be identified with $\bigotimes M_2(\mathbb{C})$, the infinite tensor product of a countably infinite family of the algebras $M_2(\mathbb{C})$ of all two by two matrices over the complex numbers field \mathbb{C} . ([4] and [12]).

Let G be any countably infinite group. We are able to construct an action β^G of G on F called the Bernoulli shift action. We can show that each β_h (h is in G with $h \neq e$, where e is the neutral element of G) is an outer automorphism of F . (See [8]). The following lemma plays a key role in our discussions :

Lemma 1. Let θ be any $*$ -automorphism of F . Then, there exists a unique extension of θ to a $*$ -automorphism $\hat{\theta}$ of \hat{F} . If θ is an outer automorphism of F , then $\hat{\theta}$ is an outer automorphism of \hat{F} .

Since F is simple, we can apply Theorem 8 in [1]. So, using Lemma 1, we can extend our Bernoulli shift action β of G on F to an outer action of G , $\hat{\beta}$ on \hat{F} . In fact, if g and h are in G , then for each x in F , we have

$$\hat{\beta}_g(\hat{\beta}_h(x)) = \hat{\beta}_g(\beta_h(x)) = \beta_g(\beta_h(x)) = \beta_{gh}(x) = \hat{\beta}_{gh}(x).$$

The uniqueness of the extension of β_{gh} to $\hat{\beta}_{gh}$, now tells us that $\hat{\beta}_{gh} = \hat{\beta}_g \hat{\beta}_h$ for each pair g and h in G . Moreover, $\hat{\beta}_g$ ($g \neq e$) is outer on \hat{F} . We call this action the Bernoulli shift action of G on \hat{F} .

Definition 1. Let M be a monotone complete C^* -algebra. Let $g \rightarrow \gamma_g$ be an action of G on M . This action is said to be ergodic if the only projections in M which are invariant under this action are 0 and 1.

It is natural to ask: When is the Bernoulli shift action on \hat{F} ergodic?

Theorem 1. $\hat{\beta}$ is ergodic on \hat{F} .

An outline of a proof. We shall assume that $\hat{\beta}$ is not ergodic, that is, there exists a non-zero projection p in \hat{F} such that $p \neq 1$ and $\hat{\beta}_g(p) = p$ for all g in G . Since $p \neq 0$, $1/2$ is not

is not an upper bound for $(-\infty, p] \cap (F_0)_h$ (where F_0 is the canonical locally finite dimensional *-subalgebra of F). So, there is an a in $(-\infty, p] \cap (F_0)_h$ such that $a \not\leq 1/2$. Thus we have $(2\|a^+\|)^{-1} < 1$ (where a^+ is the positive portion of a). Since $1 - p \neq 0$, there exists a b in $(-\infty, 1 - p] \cap (F_0)_h$ such that $\|b^+\| > (2\|a^+\|)^{-1}$.

Because our action is asymptotically abelian, there exists g in G such that $\hat{\beta}_g(a)$, $\hat{\beta}_g(a^+)$, b , b^+ all commute with each other and

$$\|\hat{\beta}_g(a^+)b^+\| = \|\hat{\beta}_g(a^+)\| \|b^+\| = \|a^+\| \|b^+\|.$$

Moreover, since p is invariant under $\hat{\beta}$, we can show that

$$0 \leq 2\hat{\beta}_g(a^+)b^+ \leq 1.$$

So $\|a^+\| \|b^+\| = \|\hat{\beta}_g(a^+)b^+\| \leq 1/2$. This is a contradiction. So, this action $\hat{\beta}$ of G on \hat{F} is ergodic.

Next we shall discuss about weak equivalences of the Bernoulli shift actions. Before going into discussions, we shall prepare monotone complete cross products of C*-dynamical systems. Let (A, G, α) be a discrete C*-dynamical system over a monotone complete C*-algebra A , where G is a countably infinite discrete group. Let $A \bar{\otimes} \mathcal{L}(l^2(G))$ be the monotone complete tensor product of A by the Type I factor $\mathcal{L}(l^2(G))$ (see [3] for details. See also [7]). We shall explain briefly about this construction. The elements of the monotone complete tensor product $A \bar{\otimes} \mathcal{L}(l^2(G))$ may be identified with those matrices $(a_{g,h})$ which correspond to elements in $A'' \bar{\otimes} \mathcal{L}(l^2(G))$ and whose entries are in A , where A'' is the second dual of A and $A'' \bar{\otimes} \mathcal{L}(l^2(G))$ is the usual von Neumann tensor product. The definition of "addition", "scalar multiplication" and "involution" of those matrices is straightforward and correspond

to these operations in $A \bar{\otimes} \mathcal{L}(l^2(G))$. "Multiplication" is more subtle, because, when multiplication $(a_{g,h})$ by $(b_{g,h})$, we need to be able to assign a value in A to infinite sums such as

$$" \sum_k a_{g,k} b_{k,h} "$$

The notion of the order convergence in A by Kadison and Pedersen is used to define

$$0\text{-} \sum_k a_{g,k} b_{k,h}$$

in such a way that $A \bar{\otimes} \mathcal{L}(l^2(G))$ is a monotone complete C^* -algebra. Hamana ([3], see also [5] and [13]) defines the monotone complete cross product associated with (A, G, α) to be the subalgebra of $A \bar{\otimes} \mathcal{L}(l^2(G))$ corresponding to the algebra of all matrices such that

$$a_{hg,kg} = \alpha_g^{-1}(a_{h,k})$$

for each g, h and k in G . We write $M(A, G, \alpha)$ to denote this algebra. We know that $M(A, G, \alpha)$ is a monotone closed $*$ -subalgebra of $A \bar{\otimes} \mathcal{L}(l^2(G))$. Let

$$\pi : A \rightarrow M(A, G, \alpha)$$

be defined by

$$\pi(a) = (\delta_{h,k} \alpha_h^{-1}(a))$$

for any a in A . Then π is a $*$ -isomorphism from A onto $D \cap M(A, G, \alpha)$, where D is the diagonal subalgebra of $A \bar{\otimes} \mathcal{L}(l^2(G))$, corresponding to those matrices which vanish off the diagonal. Let

$$E^\# : A \bar{\otimes} \mathcal{L}(l^2(G)) \rightarrow D$$

be defined by $E^\#((a_{g,h})) = (\delta_{g,h} a_{g,h})$. It is straightforward to verify that $E = E^\#|_{M(A, G, \alpha)}$ is a normal conditional expectation from $M(A, G, \alpha)$ onto $\pi(A)$. For each g in G , let

$$U_g = (\delta_{h, gk})_{h, k \text{ in } G}.$$

Then U_g is a unitary in $M(A, G, \alpha)$ such that

$$(1) \quad U_g \pi(a) U_g^* = \pi(\alpha_g(a)) \quad \text{for all } a \text{ in } A \text{ and } g \text{ in } G.$$

Straightforward calculations show that $g \rightarrow U_g$ is a group isomorphism of G into the unitary group $U(M(A, G, \alpha))$ such that

$$(2) \quad E(U_g x U_g^*) = U_g E(x) U_g^* = \alpha_g(E(x)) \quad \text{for all } a \text{ in } A \text{ and } g \text{ in } G.$$

Moreover, it follows that, for any x in $M(A, G, \alpha)$,

$$(3) \quad E(x U_g) = 0 \quad \text{for each } g \text{ in } G, \text{ then } x = 0,$$

and

$$(4) \quad E(U_g) = 0 \quad \text{if } g \neq e.$$

Definition 2. Let (A, G_1, α_1) and (A, G_2, α_2) be C^* -dynamical systems over the same monotone complete C^* -algebra A . If, there exists an isomorphism θ from $M(A, G_1, \alpha_1)$ onto $M(A, G_2, \alpha_2)$ such that θ maps $\pi_1(A)$ onto $\pi_2(A)$, then, (A, G_1, α_1) and (A, G_2, α_2) are said to be weakly equivalent. (See [7]).

The main purpose of the rest of this report is to outline a proof of the following:

Theorem 2. Let G_1 and G_2 be discrete countably infinite groups. For each j , let β^j be the Bernoulli shift action of G_j on \hat{F} . Let (\hat{F}, G_1, β^1) and (\hat{F}, G_2, β^2) be weakly equivalent. Then G_1 is isomorphic to G_2 .

If A is a monotone complete factor and α is an outer action of G on A . Then, $M(A, G, \alpha)$ is a factor. In particular, $M(\hat{F}, G, \hat{\beta})$ is a wild, monotone complete factor of Type III. ([3], see also [5] and [13]).

Let us simplify the notations. Let B be any unital C^* -algebra and let $U(B)$ be the group of all unitaries in B . Let A be a unital $*$ -subalgebra of B . We define an A -normalizing unitary to be any u in $U(B)$ such that $uAu^* = A$. We denote the group of all A -normalizing unitaries in B by $N(B:A)$. Let G be a group. We define a G system to be an ordered quadruple (B, A, U_G, E) where E is a conditional expectation from B onto A and U_G is an isomorphic image of G in $N(B:A)$ such that the following conditions are satisfied:

- (a) Let u be in U_G . If the automorphism Adu is also implemented by some unitary in A , then $u = 1$.
- (b) For each u in U_G and each b in B ,

$$E(ubu^*) = uE(b)u^*.$$
- (c) Let b be in B . If $E(ubv) = 0$ for each pair u and v in U_G , then $b = 0$.
- (d) Whenever u in U_G satisfies that $u \neq 1$, then $E(u) = 0$.

It is easy to check that $(M(\hat{F}, G, \hat{\beta}), \pi(\hat{F}), U_G, E)$ satisfies the above properties because of (1), (2), (3) and (4). Moreover, we can show the following:

Lemma 2. Let (B, A, U_G, E) be any G system and let the centre Z_A of A be one dimensional. Let x be in $B \cap A'$. Then x is a scalar multiple of 1.

Using this lemma and analysing carefully the structure of $N(B:A)$, we have the following:

Theorem 3. Let G_1 and G_2 be discrete groups. Let (B_1, A_1, U_{G_1}, E_1) and (B_2, A_2, U_{G_2}, E_2) be, respectively, a G_1 system and a G_2 system. Let θ be a $*$ -isomorphism of B_2 onto B_1 such that θ maps A_2 onto A_1 . Let A_1 have one dimensional centre. Then U_{G_1} and U_{G_2} are isomorphic.

Theorem 2 is an easy consequence of Theorem 3.

References

- [1] J. Dixmier, Sur certains espaces consideres par M.H. Stone, Summa Math. Brasil., 11(1951), 151-182.
- [2] G.A. Elliott, K. Saitô and J.D.M. Wright, Embedding AW^* -algebras as double commutants in Type I algebras, J. London Math. Soc., 28(1983), 376-384.
- [3] M. Hamana, Tensor products for monotone complete C^* -algebras, I, II, Japan. J. Math., 8(1982), 261-295.
- [4] R.V. Kadison and J.R. Ringrose, Fundamentals of the theory of operator algebras, vol. II, Academic Press(1986).
- [5] K. Saitô, AW^* -algebras with monotone convergence property and examples by Takenouchi and Dyer, Tôhoku Math. J., 31(1979), 31-40.

- [6] K. Saitô and J.D.M. Wright, Outer automorphisms of regular completions, *J. London Math. Soc.*, 27(1983), 150-156.
- [7] K. Saitô and J.D.M. Wright, On tensor products of monotone complete algebras, *Quart. J. Math. Oxford*, 35(1984), 209-221.
- [8] K. Saitô, Dynamic actions on monotone complete factors, *Quart. J. Math. Oxford*, 44(1993), 239-247.
- [9] K. Saitô and J.D.M. Wright, A continuum of discrete group actions on a Fermionic factor, To appear in *Quart. J. Math. Oxford*, 1993.
- [10] K. Saitô and J.D.M. Wright, Ergodic actions on the Fermionic factor, To appear in *Quart. J. Math. Oxford*, 1993.
- [11] D. Sullivan, B. Weiss and J.D.M. Wright, Generic dynamics and monotone complete C^* -algebras, *Trans. Amer. Math. Soc.*, 395(1986), 795-809.
- [12] Z. Takeda, Inductive limit and infinite direct product of operator algebras, *Tôhoku Math. J.*, 7(1955), 67-86.
- [13] O. Takenouchi, A non W^* , AW^* -factor, C^* -algebras and applications to Physics. (Proc. Second Japan-U.S.A. Seminar, Los Angeles (1977)), *Lecture Notes in Math.*, 650(1978), 135-139.
- [14] J.D.M. Wright, Regular σ -completion of C^* -algebras, *J. London Math. Soc.*, 12(1976), 299-309.
- [15] J.D.M. Wright, Wild AW^* -factors and Kaplansky-Rickart C^* -algebras, *J. London Math. Soc.*, 13(1976), 83-89.

Mathematical Institute
Tôhoku University
Sendai, Japan