

# Asymptotic profile of a radially symmetric solution with transition layers for an unbalanced bistable equation

沼津工業高等専門学校 Hiroshi Matsuzawa(Hiroshi Matsuzawa)  
Numazu National College of Technology

## 1 Introduction and Main Results

In this paper, we consider the following boundary value problem:

$$(P_\varepsilon) \begin{cases} -\varepsilon^2 \Delta u = h(|x|)^2(u - a(|x|))(1 - u^2) & \text{in } B_1(0) \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial B_1(0) \end{cases}$$

where  $\varepsilon > 0$  is a small parameter,  $B_1(0)$  is a unit ball in  $\mathbb{R}^N$  centered at the origin and the function  $a$  is a  $C^1$  function on  $[0, 1]$  satisfying  $-1 < a(|x|) < 1$  and  $a'(0) = 0$ . The function  $h$  is a positive  $C^1$  function on  $[0, 1]$  satisfying  $h'(0) = 0$ . We set  $r = |x|$ .

Problem  $(P_\varepsilon)$  appears in various models such as population genetics, chemical reactor theory and phase transition phenomena. See [1] and the references therein. If the function  $h$  satisfies  $h(r) \equiv 1$  and the function  $a$  satisfies  $a(r) \not\equiv 0$ , then this problem  $(P_\varepsilon)$  has been studied in [1], [4] and [7]. In this case, it is shown that there exist radially symmetric solutions with transition layers near the set  $\{x \in B_1(0) | a(|x|) = 0\}$ . If the set  $\{r \in \mathbb{R} | a(r) = 0\}$  contains an interval  $I$ , then the problem to decide the configuration of transition layer on  $I$  is more delicate.

On the other hand, in the case of  $N = 1$ , if the function  $h$  satisfies  $h(r) \not\equiv 1$  and the function  $a$  satisfies  $a(r) \equiv 0$ , then this problem  $(P_\varepsilon)$  has been studied in [8] and [9]. In this case, it is shown that there exist stable solutions with transition layers near prescribed local minimum points of  $h$ .

In this paper, we consider the case where the function  $a$  satisfies  $a(r) \not\equiv 0$  with  $a(r) = 0$  on some interval  $I \subset (0, 1)$ . We show the minimum point of the function  $r^{N-1}h(r)$  on  $I$  has very important role to decide the configuration of transition layer on  $I$  in this case.

We note that in [4], Dancer and Shusen Yan considered a problem similar to ours. They assume that  $N \geq 2$ ,  $h \equiv 1$  and the nonlinear term is  $u(u - a|x|)(1 - u)$  satisfying  $a(r) = 1/2$  on  $I = [l_1, l_2]$  and  $a(r) < 1/2$  for  $l_1 - r > 0$  small and  $a(r) > 1/2$  for  $r - l_2 > 0$  small, then a global minimizer of the corresponding functional has a transition layer near the  $l_1$ , that is, the minimum point of  $r^{N-1}$  on  $I$  (see [4, Theorem 1.3]). In this sense, we can say that our results are natural extension of the results in [4]. We are going to follow throughout the variational

procedure used in [4] with a few modifications prompted by the presence of the function  $h$ .

Here we state the energy functional corresponding to  $(P_\varepsilon)$ :

$$J_\varepsilon(u) = \int_{B_1(0)} \frac{\varepsilon^2}{2} |\nabla u|^2 - F(|x|, u) dx,$$

where  $F(|x|, u) = \int_{-1}^u f(|x|, s) ds$  and  $f(|x|, u) = h(|x|)^2(u - a(|x|))(1 - u^2)$ .

It is easy to see that the following minimization problem has a minimizer

$$\inf\{J_\varepsilon(u) | u \in H^1(B_1(0))\}. \quad (1.1)$$

Let  $A_- = \{x \in B_1(0) | a(|x|) < 0\}$  and  $A_+ = \{x \in B_1(0) | a(|x|) > 0\}$ .

In this paper, we will analyze the profile of the minimizer of (1.1). Our main theorem is the following:

**Theorem 1.1.** *Let  $u_\varepsilon$  be a global minimizer of (1.1). Then  $u_\varepsilon$  is radially symmetric and*

$$u_\varepsilon \rightarrow \begin{cases} 1 & , \text{uniformly on any compact subset of } A_-, \\ -1 & , \text{uniformly on any compact subset of } A_+, \end{cases}$$

as  $\varepsilon \rightarrow 0$ . In particular  $u_\varepsilon$  converges uniformly near the boundary of  $B_1(0)$ , that is, if  $a(r) < 0$  on  $[r_0, 1]$  for some  $r_0 > 0$ ,  $u_\varepsilon \rightarrow 1$  uniformly on  $\overline{B_1(0)} \setminus B_{r_0}(0)$  and if  $a(r) > 0$  on  $[r_0, 1]$  for some  $r_0 > 0$ ,  $u_\varepsilon \rightarrow -1$  uniformly on  $\overline{B_1(0)} \setminus B_{r_0}(0)$ . Moreover, for any  $0 < r_1 \leq r_2 < 1$  with  $a(r_i) = 0$ ,  $i = 1, 2$ ,  $a(r) \neq 0$  for  $r_1 - r > 0$  small and for  $r - r_2 > 0$  small,  $a(r) = 0$  if  $r \in [r_1, r_2]$ , we have:

(i) *If  $a(r) < 0$  for  $r_1 - r > 0$  small and  $a(r) > 0$  for  $r - r_2 > 0$ , then for any small  $\eta > 0$  and for any small  $\theta > 0$ , there exists a positive number  $\varepsilon_0$  which has the following properties: For any  $\varepsilon \in (0, \varepsilon_0]$ , there exist  $t_{\varepsilon,1} < t_{\varepsilon,2}$  such that*

(a)

$$\begin{cases} u_\varepsilon(r) > 1 - \eta & \text{for } r \in [r_1 - \theta, t_{\varepsilon,1}), \\ u_\varepsilon(t_{\varepsilon,1}) = 1 - \eta, \\ u_\varepsilon(t_{\varepsilon,2}) = -1 + \eta, \\ u_\varepsilon(r) < -1 + \eta, & \text{for } r \in (t_{\varepsilon,2}, r_2 + \theta]. \end{cases}$$

(b) *The function  $u_\varepsilon(r)$  is decreasing in  $(t_{\varepsilon,1}, t_{\varepsilon,2})$*

(c) *The inequality  $0 < R_1 \leq \frac{t_{\varepsilon,2} - t_{\varepsilon,1}}{\varepsilon} \leq R_2$  holds, where  $R_1$  and  $R_2$  are two constants independent of  $\varepsilon > 0$ .*

(d) *If  $t_{\varepsilon_j,1}, t_{\varepsilon_j,2} \rightarrow \bar{t}$  for some positive sequence  $\{\varepsilon_j\}$  converging to zero as  $j \rightarrow \infty$ , then  $\bar{t}$  satisfies  $h(\bar{t})\bar{t}^{N-1} = \min_{s \in [r_1, r_2]} h(s)s^{N-1}$ .*

(ii) If  $a(r) > 0$  for  $r_1 - r > 0$  small and  $a(r) < 0$  for  $r - r_2 > 0$ , then for any small  $\eta > 0$  and for any small  $\theta > 0$ , there exists a positive number  $\varepsilon_0$  which has the following properties: For any  $\varepsilon \in (0, \varepsilon_0]$ , there exist  $t_{\varepsilon,1} < t_{\varepsilon,2}$  such that

(a)

$$\begin{cases} u_\varepsilon(r) < -1 + \eta & \text{for } r \in [r_1 - \theta, t_{\varepsilon,1}), \\ u_\varepsilon(t_{\varepsilon,1}) = -1 + \eta, \\ u_\varepsilon(t_{\varepsilon,2}) = 1 - \eta, \\ u_\varepsilon(r) > 1 - \eta, & \text{for } r \in (t_{\varepsilon,2}, r_2 + \theta]. \end{cases}$$

(b) The function  $u_\varepsilon(r)$  is increasing in  $(t_{\varepsilon,1}, t_{\varepsilon,2})$ .

(c) The inequality  $0 < R_1 \leq \frac{t_{\varepsilon,2} - t_{\varepsilon,1}}{\varepsilon} \leq R_2$  holds, where  $R_1$  and  $R_2$  are two constants independent of  $\varepsilon > 0$ .

(d) If  $t_{\varepsilon_j,1}, t_{\varepsilon_j,2} \rightarrow \bar{t}$  for some positive sequence  $\{\varepsilon_j\}$  converging to zero as  $j \rightarrow \infty$ , then  $\bar{t}$  satisfies  $h(\bar{t})\bar{t}^{N-1} = \min_{s \in [r_1, r_2]} h(s)s^{N-1}$ .




Figure 1: The profile of the global minimizer  $u_\varepsilon$ .

**Remarks .** (i) We note that results from (a) to (c) both in cases (i) and (ii) are not related to the presence of the function  $h$ . The effect of presence of function  $h$  appears in the result (d) in (i) and (ii).

(ii) If  $\min_{s \in [r_1, r_2]} s^{N-1}h(s)$  is attained at a unique point  $\bar{t}$ , we can show  $t_{\varepsilon,1}, t_{\varepsilon,2} \rightarrow \bar{t}$  as  $\varepsilon \rightarrow 0$  without taking subsequences.

(iii) If the function  $r^{N-1}h(r)$  is constant on  $[r_1, r_2]$ , it is a very difficult problem to know the location of the point  $\bar{t} \in [r_1, r_2]$ .

This paper is organized as follows. In section 2, we prepare some preliminary results. We will prove Theorems 1.1 in section 3.

## 2 Preliminary Results

In this section we prepare some preliminary results.

Let  $D$  is a bounded domain in  $\mathbb{R}^N$ . Let  $\bar{f}(x, t)$  be a function defined on  $\bar{D} \times \mathbb{R}$  which is bounded on  $\bar{D} \times [-1, 1]$ . Suppose  $\bar{f}$  is continuous on  $t \in \mathbb{R}$  for each  $x \in \bar{D}$  and is measurable in  $D$  for each  $t \in \mathbb{R}$ . We also assume

$$\bar{f}(x, t) > 0 \text{ for } x \in \bar{D}, t < -1; \bar{f}(x, t) < 0, \text{ for } x \in \bar{D}, t > 1. \quad (2.1)$$

Consider the following minimization problem:

$$\inf \left\{ \bar{J}_\varepsilon(u, D) := \int_D \frac{\varepsilon^2}{2} |\nabla u|^2 - \bar{F}(x, u) dx : u - \eta \in H_0^1(D) \right\}, \quad (2.2)$$

where  $\eta \in H^1(D)$  with  $-1 \leq \eta \leq 1$  on  $D$  and

$$\bar{F}(x, t) = \int_{-1}^t \bar{f}(x, s) ds.$$

We can prove next two lemmas by methods similar to [4]. For readers's convenience we prove these lemmas in this section.

**Lemma 2.1.** *Suppose that  $\bar{f}(x, t)$  satisfies (2.1). Let  $u_\varepsilon$  be a minimizer of (2.2). Then  $-1 \leq u_\varepsilon \leq 1$  on  $D$ .*

*Proof.* We prove  $-1 \leq u_\varepsilon$  on  $D$ . Let  $M = \{x : u_\varepsilon(x) < -1\}$ . Define  $\tilde{u}_\varepsilon$  as follows:

$$\tilde{u}_\varepsilon(x) = \begin{cases} u_\varepsilon(x) & \text{if } x \in D \setminus M \\ -1 & \text{if } x \in M. \end{cases}$$

Since  $u_\varepsilon(x) = \eta \geq -1$  on  $\partial D$ , we see  $M$  is compactly contained in  $D$ . Thus  $\tilde{u}_\varepsilon - \eta \in H_0^1(D)$ . If the measure  $m(M)$  of  $M$  is positive, we have  $\bar{J}_\varepsilon(\tilde{u}_\varepsilon, D) < \bar{J}_\varepsilon(u_\varepsilon, D)$ . Because  $u_\varepsilon$  is a minimizer, we see  $m(M) = 0$ , where  $m(A)$  denotes the Lebesgue measure of the set  $A$ . Thus  $u_\varepsilon \geq -1$ . Similarly we can prove that  $u_\varepsilon \leq 1$ .  $\square$

**Lemma 2.2.** *Suppose that  $\bar{f}_1(x, t)$  and  $\bar{f}_2(x, t)$  both satisfy (2.1) and the same regularity assumption on  $\bar{f}$ . Assume that  $\eta_i \in H^1(D)$  satisfy  $-1 \leq \eta_i \leq 1$  on  $D$  for  $i = 1, 2$ . Let  $u_{\varepsilon, i}$  be a corresponding minimizer of (2.2), where  $\bar{f} = \bar{f}_i$  and  $\eta = \eta_i$ ,  $i = 1, 2$ . Suppose that  $\bar{f}_1(x, t) \geq \bar{f}_2(x, t)$  for all  $(x, t) \in \bar{D} \times [-1, 1]$  and  $1 \geq \eta_1 \geq \eta_2 \geq -1$ . Then  $u_{\varepsilon, 1} \geq u_{\varepsilon, 2}$ .*

*Proof.* Let  $M = \{x \in D : u_{\varepsilon, 2} > u_{\varepsilon, 1}\}$ . Define  $\varphi_\varepsilon = (u_{\varepsilon, 2} - u_{\varepsilon, 1})^+$ . Since  $\eta_1 \geq \eta_2$ , we have  $\varphi_\varepsilon \in H_0^1(D)$ . Set  $\bar{F}_i(x, u) = \int_{-1}^u \bar{f}_i(x, s) ds$ . Since  $u_{\varepsilon, i}$  is a minimizer of

$$J_{\varepsilon, i}(u) := \int_D \frac{\varepsilon^2}{2} |\nabla u|^2 - \bar{F}_i(x, u) dx$$

and  $\varphi_\varepsilon = 0$  for  $x \in D \setminus M$ , we have

$$\begin{aligned} 0 &\leq J_{\varepsilon,1}(u_{\varepsilon,1} + \varphi_\varepsilon) - J_{\varepsilon,1}(u_{\varepsilon,1}) \\ &= \int_M \frac{\varepsilon^2}{2} (|\nabla(u_{\varepsilon,1} + \varphi_\varepsilon)|^2 - |\nabla u_{\varepsilon,1}|^2) dx - \int_M \int_{u_{\varepsilon,1}}^{u_{\varepsilon,1} + \varphi_\varepsilon} \bar{f}_1(x, s) ds \\ &\leq \int_M \frac{\varepsilon^2}{2} (|\nabla(u_{\varepsilon,1} + \varphi_\varepsilon)|^2 - |\nabla u_{\varepsilon,1}|^2) dx - \int_M \int_{u_{\varepsilon,1}}^{u_{\varepsilon,1} + \varphi_\varepsilon} \bar{f}_2(x, s) ds \\ &= J_{\varepsilon,2}(u_{\varepsilon,2}) - J_{\varepsilon,2}(u_{\varepsilon,2} - \varphi_\varepsilon) \leq 0. \end{aligned}$$

This implies that  $u_{\varepsilon,1} + \varphi_\varepsilon$  is also a minimizer of  $J_{\varepsilon,1}(u)$ . Let  $L > 0$  be large enough such that  $\bar{f}_1(x, t) + Lt$  is strictly increasing for  $x \in \bar{D}$ ,  $t \in [-1, 1]$ . From

$$-\varepsilon^2 \Delta(u_{\varepsilon,1} + \varphi_\varepsilon) = \bar{f}_1(u_{\varepsilon,1} + \varphi_\varepsilon),$$

we obtain

$$-\varepsilon^2 \Delta \varphi_\varepsilon = \bar{f}_1(u_{\varepsilon,1} + \varphi_\varepsilon) - \bar{f}_1(u_{\varepsilon,1}).$$

Thus

$$-\varepsilon^2 \Delta \varphi_\varepsilon + L\varphi_\varepsilon = \bar{f}_1(u_{\varepsilon,1} + \varphi_\varepsilon) + L(u_{\varepsilon,1} + \varphi_\varepsilon) - (\bar{f}_1(u_{\varepsilon,1}) + Lu_{\varepsilon,1}) > 0$$

in  $D$ . Fix  $z_0 \in M$ . Let  $x_0 \in \partial M$  such that  $|x_0 - z_0| = \text{dist}(z_0, \partial M)$ . Using the Strong maximum principle and Hopf's lemma in  $B_{\text{dist}(z_0, \partial M)}(z_0)$ , we obtain that  $\frac{\partial \varphi_\varepsilon}{\partial \nu}(x_0) < 0$ , where  $\nu = (x_0 - z_0)/|x_0 - z_0|$ . But  $\varphi_\varepsilon(x) = 0$  for  $x \notin M$ . Thus,  $\frac{\partial \varphi_\varepsilon}{\partial \nu}(x_0) = 0$ . This is a contradiction. Thus we obtain  $M = \emptyset$ .  $\square$

### 3 Proof of Main Theorem

In this section we prove Theorem 1.1. The following proposition is the first part of Theorem 1.1.

**Proposition 3.1.** *Let  $u_\varepsilon$  be a global minimizer of the problem (1.1). Then  $u_\varepsilon$  satisfies*

$$u_\varepsilon \rightarrow \begin{cases} 1 & \text{uniformly on any compact subset of } A_- \\ -1 & \text{uniformly on any compact subset of } A_+ \end{cases}$$

as  $\varepsilon \rightarrow 0$ .

*Proof.* Let  $x_0 \in A_-$ . Choose  $\delta > 0$  small so that  $B_\delta(x_0) \subset\subset A$ . Take  $b \in (\max_{z \in \overline{B_\delta(x_0)}} a(z), 1/2)$ . Define  $f_{x_0, \delta, b}(t) = (\min_{z \in B_\delta(x_0)} h(z)^2)(t - b)(1 - t^2)$ . Then for  $x \in \overline{B_\delta(x_0)}$ ,  $t \in [-1, 1]$ , we have  $f(|x|, t) \geq f_{x_0, \delta, b}(t)$ . Let  $u_{\varepsilon, x_0, \delta, b}$  be the minimizer of

$$\inf \left\{ \int_{B_\delta(x_0)} \frac{\varepsilon^2}{2} |\nabla u|^2 - F_{x_0, \delta, b}(u) dx : u + 1 \in H_0^1(B_\delta(x_0)) \right\},$$

where  $F_{x_0, \delta, b}(t) = \int_{-1}^t f_{x_0, \delta, b}(s) ds$ . It follows from Lemmas 2.1 and 2.2 that

$$u_{\varepsilon, x_0, \delta, b}(x) \leq u_\varepsilon(x) \leq 1, \text{ for } x \in B_\delta(x_0).$$

Since  $\int_{-1}^1 f_{x_0, \delta, b}(s) ds > 0$ , it follows from [2, 3] that  $u_{\varepsilon, x_0, \delta, b}(x) \rightarrow 1$  as  $\varepsilon \rightarrow 0$  uniformly in  $B_{\delta/2}(x_0)$ , thus  $u_\varepsilon(x) \rightarrow 1$  as  $\varepsilon \rightarrow 0$  uniformly in  $B_{\delta/2}(x_0)$ .  $\square$

To prove the rest of Theorem 1.1, we need the following proposition and lemma.

**Proposition 3.2.** *Let  $u$  be a local minimizer of the following problem:*

$$\inf \left\{ \int_{B_1(0)} \frac{1}{2} |\nabla u|^2 - G(|x|, u) dx : u \in H^1(B_1(0)) \right\}.$$

Here  $G(r, t) = \int_{-1}^t g(r, s) ds$ ,  $g(r, t)$  is  $C^1$  in  $t \in \mathbb{R}$  for each  $r \geq 0$ ,  $g(r, t)$  and  $g_t(r, t)$  are measurable on  $[0, +\infty)$  for each  $t \in \mathbb{R}$ ,  $g(r, t) < 0$  if  $t < -1$  or  $t > 1$  and  $|g(r, t)| + |g_t(r, t)|$  is bounded on  $[0, k] \times [-2, 2]$  for any  $k > 0$ . Then  $u$  is radial, i.e.,  $u(x) = u(|x|)$ .

*Proof.* See [4, Proposition 2.6].  $\square$

Before we prove Theorem 1.1, we prepare a lemma.

**Lemma 3.3.** *Let  $0 < \eta < 1$  be any fixed constant and  $w$  satisfies*

$$\begin{cases} -w_{zz} = w(1 - w^2) & \text{on } \mathbb{R}, \\ w(0) = -1 + \eta \text{ (resp. } w(0) = 1 - \eta), \\ w(z) \leq -1 + \eta \text{ (resp. } w(z) \geq 1 - \eta) & \text{for } z \leq 0, \\ w \text{ is bounded on } \mathbb{R}. \end{cases}$$

*Then  $w$  is a unique solution of*

$$\begin{cases} -w_{zz} = w(1 - w^2) & \text{on } \mathbb{R}, \\ w(0) = -1 + \eta \text{ (resp. } w(0) = 1 - \eta), \\ w'(z) > 0 \text{ (resp. } w'(z) < 0) & z \in \mathbb{R}, \\ w(z) \rightarrow \pm 1 \text{ (resp. } w(z) \rightarrow \mp 1) & \text{as } z \rightarrow \pm\infty. \end{cases}$$

*Proof.* See for example [6].  $\square$

Now we prove the rest of Theorem 1.1.

*Proof of Theorem 1.1.* For the sake of simplicity, we prove for the case where  $a(r) < 0$  on  $[0, r_1]$ ,  $a(r) = 0$  on  $[r_1, r_2]$  and  $a(r) > 0$  on  $(r_2, 1]$  for some  $0 < r_1 < r_2 < 1$  (see Figure 1 in Section 1).

**Part 1.** First we show that  $u_\varepsilon$  converges uniformly near the boundary of  $B_1(0)$ , that is,  $u_\varepsilon \rightarrow -1$  uniformly on  $\overline{B_1(0)} \setminus B_{r_2+\tau}(0)$  for any small  $\tau > 0$ . We note that

we have  $u_\varepsilon \rightarrow -1$  uniformly on  $\overline{B_{1-\tau}(0)} \setminus B_{r_2+\tau}(0)$  as  $\varepsilon \rightarrow 0$ . Now we claim that  $u_\varepsilon(r) \leq u_\varepsilon(1-\tau) =: T_\varepsilon$  for  $r \in [1-\tau, 1]$ . We define the function  $\tilde{u}_\varepsilon$  as follows:

$$\tilde{u}_\varepsilon(r) = \begin{cases} u_\varepsilon(r) & \text{if } r \in [0, 1-\tau] \\ u_\varepsilon(r) & \text{if } u_\varepsilon(r) < T_\varepsilon \text{ and } r \in [1-\tau, 1], \\ T_\varepsilon & \text{if } u_\varepsilon(r) \geq T_\varepsilon \text{ and } r \in [1-\tau, 1]. \end{cases}$$

We note that  $\tilde{u}_\varepsilon \in H^1(B_1(0))$  and  $-F(r, T_\varepsilon) \leq -F(r, t)$  for  $\varepsilon > 0$  and  $|r-1|$  small and  $t \geq T_\varepsilon$ . Hence we obtain  $J_\varepsilon(\tilde{u}_\varepsilon) < J_\varepsilon(u_\varepsilon)$  and we have a contradiction if we assume that the measure of the set  $\{r \in [0, 1] \mid u_\varepsilon(r) > T_\varepsilon \text{ and } r \in [1-\tau, 1]\}$  is positive. Hence  $-1 < u_\varepsilon(r) \leq T_\varepsilon$  and  $u_\varepsilon \rightarrow -1$  uniformly on  $\overline{B_1(0)} \setminus B_{r_2+\tau}(0)$ .

**Part 2.** Next we remark that, by Proposition 3.2,  $u_\varepsilon$  is radially symmetric and we note that for any  $t_2 > t_1$ ,  $u_\varepsilon$  is a minimizer of the following problem

$$\inf\{J_\varepsilon(u, B_{t_2}(0) \setminus \overline{B_{t_1}(0)}) : u - u_\varepsilon \in H_0^1(B_{t_2}(0) \setminus \overline{B_{t_1}(0)})\},$$

where

$$J_\varepsilon(u, M) = \int_M \frac{\varepsilon^2}{2} |\nabla u|^2 - F(|x|, u) dx$$

for any open set  $M$ . Let  $m_{\varepsilon, t_1, t_2}$  be the minimum value of this minimization problem.

In this part we show that  $u_\varepsilon$  has exactly one layer near the interval  $[r_1, r_2]$ .

**Step 2.1.** First we estimate the energy of transition layer.

Let  $\eta > 0$  and  $\theta > 0$  be small numbers. Since  $u_\varepsilon \rightarrow 1$  uniformly on  $[0, r_1 - \theta]$  and  $u_\varepsilon \rightarrow -1$  uniformly on  $[r_2 + \theta, 1 - \theta]$ , we can find  $\bar{r}_\varepsilon \in (r_1 - \theta, r_2 + \theta)$  such that  $u_\varepsilon(r) \geq 1 - \eta$  if  $r \in [0, \bar{r}_\varepsilon]$ ,  $u_\varepsilon(r) < 1 - \eta$  for  $r - \bar{r}_\varepsilon > 0$  small. Let  $\tilde{r}_\varepsilon > \bar{r}_\varepsilon$  be such that  $u_\varepsilon(r) \leq \eta$  if  $r \in [\tilde{r}_\varepsilon, 1 - \theta]$ ,  $u_\varepsilon(r) > \eta$  for  $\tilde{r}_\varepsilon - r > 0$  small. We may assume that  $\bar{r}_\varepsilon \rightarrow \bar{r} \in [r_1, r_2]$  and  $\tilde{r}_\varepsilon \rightarrow \tilde{r} \in [r_1, r_2]$

We employ the so-called blow-up argument. Let  $v_\varepsilon(t) = u_\varepsilon(\varepsilon t + \bar{r}_\varepsilon)$ . Then

$$-v_\varepsilon'' - \varepsilon \frac{N-1}{\varepsilon t + \bar{r}_\varepsilon} v_\varepsilon' = f(\varepsilon t + \bar{r}_\varepsilon, v_\varepsilon),$$

$-1 \leq v_\varepsilon \leq 1$  and  $v_\varepsilon(0) = 1 - \eta$ . Since  $\bar{r}_\varepsilon \rightarrow \bar{r} \in [r_1, r_2]$ , it is easy to see that  $v_\varepsilon \rightarrow v$  in  $C_{\text{loc}}^1(\mathbb{R})$  and

$$-v'' = h(\bar{r})^2(v - v^3), \quad t \in \mathbb{R}.$$

and  $v(t) \geq 1 - \eta$  for  $t \leq 0$ . If we set  $v(t) = V(h(\bar{r})t)$ , the function  $V(t)$  satisfies

$$\begin{cases} -V'' = V - V^3 & \text{on } \mathbb{R}, \\ V(0) = 1 - \eta, \\ V'(t) \geq 1 - \eta & t \leq 0. \end{cases} \quad (3.1)$$

Hence by Lemma 3.3, the function  $V$  is a unique solution for

$$\begin{cases} -V'' = V - V^3 & \text{on } \mathbb{R}, \\ V(0) = 1 - \eta, \\ V'(t) < 0 & t \leq 0. \\ V(t) \rightarrow \pm 1 & \text{as } t \rightarrow \mp\infty. \end{cases} \quad (3.2)$$

Thus, we can find an  $R > 0$  large, such that  $v(R) = \eta$ . Since  $v_\varepsilon \rightarrow v$  in  $C_{\text{loc}}^1(\mathbb{R})$ , we can find an  $R_\varepsilon \in (R - 1, R + 1)$ , such that  $v'_\varepsilon(r) < 0$  if  $r \in [0, R_\varepsilon]$  and  $v_\varepsilon(R_\varepsilon) = -1 + \eta$ . Hence  $u'_\varepsilon(r) < 0$  if  $r \in [\bar{r}_\varepsilon, \bar{r}_\varepsilon + \varepsilon R_\varepsilon]$  and  $u_\varepsilon(\bar{r}_\varepsilon + \varepsilon R_\varepsilon) = -1 + \eta$ . Then we have

$$\begin{aligned} & J_\varepsilon(u_\varepsilon, B_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}) \\ &= \omega_{N-1}(\bar{r}_\varepsilon^{N-1} + o_\varepsilon(1)) \int_{\bar{r}_\varepsilon}^{\bar{r}_\varepsilon + \varepsilon R_\varepsilon} \left( \frac{\varepsilon^2}{2} |u'_\varepsilon|^2 - F(t, u_\varepsilon) \right) dt \\ &= \omega_{N-1}(\bar{r}_\varepsilon^{N-1} + o_\varepsilon(1)) \varepsilon \int_0^{R_\varepsilon} \left( \frac{1}{2} |v'_\varepsilon|^2 - F(\varepsilon t + \bar{r}_\varepsilon, v_\varepsilon) \right) dt \\ &= \omega_{N-1}(\bar{r}_\varepsilon^{N-1} + o_\varepsilon(1)) (\beta_{h(\bar{r})} + O(\eta) + o_\varepsilon(1)) \varepsilon, \end{aligned} \quad (3.3)$$

where  $\omega_{N-1}$  is the area of the unit sphere in  $\mathbb{R}^N$ ,  $o_\varepsilon(1) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ ,  $\beta_{h(s)}$  is the positive value defined by

$$\begin{aligned} \beta_{h(s)} &= \int_{-\infty}^{+\infty} \left( \frac{1}{2} |w'_{h(s)}(t)|^2 + h(s)^2 \frac{(w_{h(s)}^2 - 1)^2}{4} \right) dt \\ &= h(s) \int_{-\infty}^{+\infty} \frac{1}{2} |V'(t)|^2 + \frac{(V(t)^2 - 1)^2}{4} dt \\ &= h(s) \beta_1 \end{aligned}$$

and  $w_{h(s)}(t) = V(h(s)t)$  for  $s \in [0, 1]$ . We note that although the function  $V$  depends on  $\eta$ , the value

$$\beta_1 = \int_{-\infty}^{+\infty} \frac{1}{2} |V'(t)|^2 + \frac{(V(t)^2 - 1)^2}{4} dt$$

is independent of  $\eta$ .

**Step 2.2.** We claim  $u_\varepsilon$  has exactly one layer near the interval  $[r_1, r_2]$ . To show  $u_\varepsilon$  has exactly one layer near the interval  $[r_1, r_2]$ , it sufficient to prove the following claim:

**Claim.**  $\tilde{r}_\varepsilon = \bar{r}_\varepsilon + \varepsilon R_\varepsilon$ .

Suppose that the claim is not true. Then we can find a  $t_\varepsilon > \bar{r}_\varepsilon + R_\varepsilon \varepsilon$  such that  $u_\varepsilon(r) < -1 + \eta$  if  $r \in (\bar{r}_\varepsilon + R_\varepsilon \varepsilon, t_\varepsilon)$ ,  $u_\varepsilon(t_\varepsilon) = -1 + \eta$ . Thus we can use the blow-up argument again at  $t_\varepsilon$  to deduce that there is a  $\tilde{t}_\varepsilon = t_\varepsilon + \varepsilon \tilde{R}_\varepsilon$  with  $u'_\varepsilon(r) > 0$  if



$r \in (t_\varepsilon, \tilde{t}_\varepsilon)$ ,  $u_\varepsilon(\tilde{t}_\varepsilon) = 1 - \eta$ . We may assume that  $t_\varepsilon, \tilde{t}_\varepsilon \rightarrow \bar{t}$  as  $\varepsilon \rightarrow 0$  for some  $\bar{t} \in [r_2, r_3]$ . Moreover

$$J_\varepsilon(u_\varepsilon, B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{t_\varepsilon}(0)}) = \omega_{N-1}(t_\varepsilon^{N-1} + o_\varepsilon(1))(\beta_{h(\bar{t})} + O(\eta))\varepsilon + o_\varepsilon(1) \quad (3.4)$$

Now we claim  $\tilde{t}_\varepsilon \geq r_1$ . Suppose  $\tilde{t}_\varepsilon < r_1$ .

Let  $F_a(t) = \int_{-1}^t (v-a)(1-v^2)dv$ . Then for any  $t > 0$  small and  $s \in [-1+t, 1-t]$ ,

$$\begin{aligned} & F_a(1-t) - F_a(s) \\ &= F_0(1-t) - F_0(s) + F_a(1-t) - F_0(1-t) - F_a(s) + F_0(s) \\ &= \left[ \frac{(v^2-1)^2}{4} \right]_s^{1-t} - a \int_s^{1-t} (1-v^2)dv \end{aligned} \quad (3.5)$$

Thus it follows from (3.5) that if  $a < 0$  then

$$F_a(1-t) - F_a(s) > 0 \quad (3.6)$$

for  $s \in [-1+t, 1-t]$ . Define

$$\bar{u}_\varepsilon(r) := \begin{cases} 1 - \eta & r \in [\bar{r}_\varepsilon, \bar{r}_\varepsilon + R_\varepsilon \varepsilon] \cup [t_\varepsilon, \tilde{t}_\varepsilon], \\ -u_\varepsilon(r) & r \in [\bar{r}_\varepsilon + R_\varepsilon \varepsilon, t_\varepsilon]. \end{cases}$$

By the assumption that  $\tilde{t}_\varepsilon < r_1$  and using (3.6), we see  $F(r, u_\varepsilon) < F(r, \bar{u}_\varepsilon)$  if  $r \in [\bar{r}_\varepsilon, \tilde{t}_\varepsilon]$ . Hence, we obtain

$$J_\varepsilon(\bar{u}_\varepsilon, B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}) < J_\varepsilon(u_\varepsilon, B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}).$$

Thus we obtain a contradiction. Therefore we have that  $\tilde{t}_\varepsilon \geq r_1$ .

Since  $a(r) \geq 0$  for  $r \in [r_1, 1]$ , we see  $F(r, t) \leq F(r, -1) = 0$  if  $r \in [r_1, 1]$ . Since  $u_\varepsilon(r) \in (-1, -1 + \eta)$  for  $r \in [\bar{r}_\varepsilon + R_\varepsilon \varepsilon, t_\varepsilon]$ , we have

$$\begin{aligned} m_{\varepsilon, \bar{r}_\varepsilon, \tilde{r}_\varepsilon} &= J_\varepsilon(\bar{u}_\varepsilon, B_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}) + J_\varepsilon(\bar{u}_\varepsilon, B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{t_\varepsilon}(0)}) \\ &\quad + J_\varepsilon(\bar{u}_\varepsilon, B_{t_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}(0)}) + J_\varepsilon(\bar{u}_\varepsilon, B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{\tilde{t}_\varepsilon}(0)}) \\ &\geq \omega_{N-1}(\bar{r}_\varepsilon^{N-1} \beta_{h(\bar{r})} \varepsilon + t_\varepsilon^{N-1} \beta_{h(\bar{t})} \varepsilon) + O(\eta \varepsilon) + o(\varepsilon) \\ &\quad + \inf \left\{ - \int_{B_{t_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}(0)}} F(r, w) : -1 \leq w \leq 1 + \eta \right\} \\ &\quad + \inf \left\{ - \int_{B_{\tilde{t}_\varepsilon}(0) \setminus \overline{B_{\tilde{t}_\varepsilon}(0)}} F(r, w) : -1 \leq w \leq 1 \right\} \\ &\geq \omega_{N-1}(\bar{r}_\varepsilon^{N-1} \beta_{h(\bar{r})} \varepsilon + t_\varepsilon^{N-1} \beta_{h(\bar{t})} \varepsilon) + O(\eta \varepsilon) + o(\varepsilon) \end{aligned} \quad (3.7)$$

Now we give an upper bound for  $m_{\varepsilon, \bar{r}_\varepsilon, \tilde{r}_\varepsilon}$ . Let  $R > 0$  be such that  $V(h(\bar{r})R) = \eta$ , where  $V$  is a unique solution to (3.2). Define  $\bar{u}_\varepsilon$  as follows:

$$\bar{u}_\varepsilon(r) := \begin{cases} V(h(\bar{r})\frac{r-\bar{r}_\varepsilon}{\varepsilon}) & r \in [\bar{r}_\varepsilon, \bar{r}_\varepsilon + \varepsilon R] \\ -1 + \eta - \frac{\eta}{\varepsilon}(r - \bar{r}_\varepsilon - \varepsilon R) & r \in [\bar{r}_\varepsilon + \varepsilon R, \bar{r}_\varepsilon + \varepsilon R + \varepsilon] \\ -1 & r \in [\bar{r}_\varepsilon + \varepsilon R + \varepsilon, \tilde{r}_\varepsilon - \varepsilon] \\ -1 + \frac{\eta}{\varepsilon}(r - \tilde{r}_\varepsilon + \varepsilon) & r \in [\tilde{r}_\varepsilon - \varepsilon, \tilde{r}_\varepsilon] \end{cases} \quad (3.8)$$

Now we note that  $|F(r, t)| = O(\eta)$  for  $r \in [\bar{r}_\varepsilon, \tilde{r}_\varepsilon]$  and  $-1 \leq t \leq -1 + \eta$ . Then we have

$$\begin{aligned} m_{\varepsilon, \bar{r}_\varepsilon, \tilde{r}_\varepsilon} &\leq J_\varepsilon(\bar{u}_\varepsilon, B_{\bar{r}_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}) \\ &\leq J_\varepsilon(\bar{u}_\varepsilon, B_{\bar{r}_\varepsilon + R\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon}(0)}) + J_\varepsilon(\bar{u}_\varepsilon, B_{\bar{r}_\varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon - \varepsilon}(0)}) \\ &\quad + J_\varepsilon(\bar{u}_\varepsilon, B_{\bar{r}_\varepsilon - \varepsilon}(0) \setminus \overline{B_{\bar{r}_\varepsilon + \varepsilon R}(0)}) \\ &\leq \omega_{N-1} \bar{r}_\varepsilon^{N-1} (\beta_{h(\bar{r})} + O(\eta)) \varepsilon + o(\varepsilon) + O(\varepsilon \eta) + o(\varepsilon) \\ &= \omega_{N-1} \bar{r}_\varepsilon^{N-1} \beta_{h(\bar{r})} + O(\eta \varepsilon) + o(\varepsilon) \end{aligned} \quad (3.9)$$

By (3.7) and (3.9), we have

$$\omega_{N-1} (\bar{r}_\varepsilon^{N-1} \beta_{h(\bar{r})} + t_\varepsilon^{N-1} \beta_{h(\bar{t})}) \varepsilon \leq \omega_{N-1} \bar{r}_\varepsilon^{N-1} \beta_{h(\bar{r})} \varepsilon + O(\varepsilon \eta) + o(\varepsilon)$$

This is a contradiction. So we can conclude  $\tilde{r}_\varepsilon = \bar{r}_\varepsilon + \varepsilon R_\varepsilon$ .

**Part 3.** It remains to prove that if  $\bar{r}_{\varepsilon_j} \rightarrow \bar{r}$  for some positive sequence  $\{\varepsilon_j\}$  converging to zero as  $j \rightarrow \infty$  then  $\bar{r}$  satisfies

$$\bar{r}^{N-1} h(\bar{r}) = \min_{s \in [r_1, r_2]} s^{N-1} h(s).$$

**Step 3.1.** First we note that from Part 1, the function  $u_\varepsilon$  satisfies  $-1 \leq u_\varepsilon \leq -1 + \eta$  for  $r \in [\bar{r}_\varepsilon + \varepsilon R_\varepsilon, 1]$  in this case.

**Step 3.2.** Set  $H(s) = s^{N-1} h(s)$ . Assume that the result is not true. Then there exists a subsequence of  $\{\bar{r}_\varepsilon\}$  (denoted by  $\bar{r}_\varepsilon$ ) such that  $\bar{r}_\varepsilon \rightarrow r' \in [r_1, r_2]$  and  $H(r') > \min_{s \in [r_1, r_2]} H(s)$ . Then we can find a point  $\bar{t} \in (r_1, r_2)$  such that  $H(r') > H(\bar{t})$ .

Next we give a lower estimate for  $J_\varepsilon(u_\varepsilon)$ . We have

$$\begin{aligned} J_\varepsilon(u_\varepsilon) &= J_\varepsilon(u_\varepsilon, B_{\bar{r}_\varepsilon}(0)) + J_\varepsilon(u_\varepsilon, B_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}(0) \setminus B_{\bar{r}_\varepsilon}(0)) \\ &\quad + J_\varepsilon(u_\varepsilon, B_1(0) \setminus \overline{B_{\bar{r}_\varepsilon + R_\varepsilon \varepsilon}(0)}). \end{aligned} \quad (3.10)$$

First we note that  $1 - \eta \leq u_\varepsilon(r) \leq 1$  for  $r \leq \bar{r}_\varepsilon$  and for sufficiently small  $\eta > 0$ ,  $-F(r, u) \geq -F(r, 1)$  ( $u \in [1 - \eta, 1]$ ). We also remark that since  $a(r) < 0$  for  $r < r_1$  and  $a(r) = 0$  for  $r_1 \leq r \leq r_2$  and  $a(r) > 0$  for  $r > r_2$ , we have  $-F(r, 1) < 0$  for

$r < r_1$  and  $-F(r, 1) = 0$  for  $r_1 \leq r \leq r_2$  and  $-F(r, 1) > 0$  for  $r > r_2$ . Hence we have  $-\int_{r_1}^{\bar{r}_\varepsilon} r^{N-1} F(r, 1) dr \geq 0$  and we obtain the following estimate

$$\begin{aligned} J_\varepsilon(u_\varepsilon, B_{\bar{r}_\varepsilon}(0)) &\geq -\int_0^{\bar{r}_\varepsilon} r^{N-1} F(r, u_\varepsilon) dr \\ &\geq -\int_0^{\bar{r}_\varepsilon} r^{N-1} F(r, 1) dr \\ &= -\int_0^{r_1} r^{N-1} F(r, 1) dr - \int_{r_1}^{\bar{r}_\varepsilon} r^{N-1} F(r, 1) dr \\ &\geq -\int_0^{r_1} r^{N-1} F(r, 1) dr =: A. \end{aligned}$$

We also obtain

$$J_\varepsilon(u_\varepsilon, B_{\bar{r}_\varepsilon + R_\varepsilon}(0) \setminus B_{\bar{r}_\varepsilon}(0)) \geq \omega_{N-1} H(r') \beta_1 \varepsilon + O(\eta \varepsilon) + o(\varepsilon). \quad (3.11)$$

by methods similar to proof of (3.3).

Since  $-1 \leq u_\varepsilon(r) \leq -1 + \eta$  for  $r \geq \bar{r}_\varepsilon + \varepsilon R_\varepsilon$  and for sufficiently small  $\eta > 0$ ,  $-F(r, u) \geq -F(r, -1) = 0$  ( $u \in [-1, -1 + \eta]$ ), we obtain the following estimate:

$$\begin{aligned} J_\varepsilon(u_\varepsilon, B_1(0) \setminus B_{\bar{r}_\varepsilon + R_\varepsilon}(0)) &\geq -\int_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}^1 r^{N-1} F(r, u_\varepsilon) dr \\ &\geq -\int_{\bar{r}_\varepsilon + \varepsilon R_\varepsilon}^1 r^{N-1} F(r, -1) dr = 0. \end{aligned} \quad (3.12)$$

Thus we obtain

$$J(u_\varepsilon) \geq A + \omega_{N-1} H(r') \beta_1 \varepsilon + O(\eta \varepsilon) + o(\varepsilon). \quad (3.13)$$


Next we give an upper bound for  $J_\varepsilon(u_\varepsilon)$ . Consider the following function  $\bar{w}_\varepsilon$ :

$$\bar{w}_\varepsilon(r) := \begin{cases} 1 & r \in [0, \bar{t} - \varepsilon] \\ 1 - \frac{\eta}{\varepsilon}(r - \bar{t} + \varepsilon) & r \in [\bar{t} - \varepsilon, \bar{t}] \\ V\left(h(\bar{t}) \frac{r - \bar{t}}{\varepsilon}\right) & r \in [\bar{t}, \bar{t} + \varepsilon R'] \\ -1 - \frac{\eta}{\varepsilon}(r - \bar{t} - \varepsilon R' - \varepsilon) & r \in [\bar{t} + \varepsilon R', \bar{t} + \varepsilon R' + \varepsilon] \\ -1 & r \in [\bar{t} + \varepsilon R' + \varepsilon, 1], \end{cases}$$

where  $R' > 0$  is the number satisfying  $V(h(\bar{t})R') = -1 + \eta$ . Then we can see

$$J_\varepsilon(u_\varepsilon) \leq J_\varepsilon(\bar{w}_\varepsilon) \leq A + \omega_{N-1} H(\bar{t}) \beta_1 \varepsilon + O(\eta \varepsilon) + o(\varepsilon). \quad (3.14)$$

By (3.13) and (3.14) we have a contradiction. The proof of Theorem 1.1 is completed. In the more complicated case, we can show by similar method (see Remark below).  $\square$



⊗ 2:

**Remark .** We briefly show in more complicated case, that is, when  $a$  is the function as in Figure 2. More precisely we set  $I_1 := [r_1, r_2]$  and  $I_2 := [r_3, r_4]$  and we assume  $a > 0$  on  $[0, r_1] \cup (r_4, 1]$  and  $a < 0$  on  $(r_3, r_4)$ .

Let  $\eta > 0$  and  $\theta > 0$  be small numbers. As in Part 1, we can find pairs of numbers  $(\bar{r}_{1,\varepsilon}, \bar{r}_{2,\varepsilon})$  and  $(R_{1,\varepsilon}, R_{2,\varepsilon})$  satisfying  $\bar{r}_{1,\varepsilon} \in (r_1 - \theta, r_2 + \theta)$ ,  $\bar{r}_{2,\varepsilon} \in (r_3 - \theta, r_4 + \theta)$ ,  $\sup_\varepsilon |R_{1,\varepsilon}| < \infty$ ,  $\sup_\varepsilon |R_{2,\varepsilon}| < \infty$  and

$$\begin{cases} u_\varepsilon(r) < -1 + \eta & \text{for } 0 < r < \bar{r}_{1,\varepsilon} \\ u_\varepsilon(\bar{r}_{1,\varepsilon}) = -1 + \eta \\ u_\varepsilon(\bar{r}_{1,\varepsilon} + \varepsilon R_{1,\varepsilon}) = 1 - \eta \\ u_\varepsilon(r) > 1 - \eta & \text{for } \bar{r}_{1,\varepsilon} + \varepsilon R_{1,\varepsilon} < r < \bar{r}_{2,\varepsilon} \\ u_\varepsilon(\bar{r}_{2,\varepsilon}) = 1 - \eta \\ u_\varepsilon(\bar{r}_{2,\varepsilon} + \varepsilon R_{2,\varepsilon}) = -1 + \eta \\ u_\varepsilon(r) < -1 + \eta & \text{for } \bar{r}_{2,\varepsilon} + \varepsilon R_{2,\varepsilon} < r < 1 \end{cases}$$

We assume  $\bar{r}_{1,\varepsilon_j} \rightarrow \bar{r}_1 \in I_1$  and  $\bar{r}_{2,\varepsilon_j} \rightarrow \bar{r}_2 \in I_2$  for some sequence  $\{\varepsilon_j\}$  which converges to 0 as  $j \rightarrow \infty$ . In this case it is easy to show that the energy of global minimizer  $J(u_\varepsilon)$  is estimated as follows:

$$J_{\varepsilon_j}(u_{\varepsilon_j}) \geq J_{\varepsilon_j}(u_{\varepsilon_j}, B_{r_2-\varepsilon}(0)) + \varepsilon_j \omega_{N-1} H(\bar{r}_2) \beta_1 + B + O(\varepsilon_j \eta) + o(\varepsilon_j), \quad (3.15)$$

where  $B = - \int_{r_2}^{r_3} r^{N-1} F(r, 1) dr$ .

Let us assume the result does not hold. Then  $H(\bar{r}_1) > \min_{s \in I_1} H(s)$  or  $H(\bar{r}_2) > \min_{s \in I_2} H(s)$  hold. We assume  $H(\bar{r}_1) = \min_{s \in I_1} H(s)$  and  $H(\bar{r}_2) > \min_{s \in I_2} H(s)$ . We also assume  $r_1 = \bar{r}_1$ . We note that if  $H(\bar{r}_1) > \min_{s \in I_1} H(s)$  or  $\bar{r}_1 \in \text{int} I_1$ , the proof is more easy.

Let us take  $\tilde{r}_2 \in \text{int}I_2$  such that  $H(\bar{r}_2) > H(\tilde{r}_2) > \min_{s \in I_2} H(s)$  and consider the following function:

$$\tilde{u}_\varepsilon(r) := \begin{cases} u_\varepsilon(r) & \text{on } [0, r_2 - \varepsilon) \\ 1 + \frac{\eta}{\varepsilon}(r - r_2) & \text{on } [r_2 - \varepsilon, r_2] \\ 1 & \text{on } [r_2, \tilde{r}_2 - \varepsilon] \\ 1 - \frac{\eta}{\varepsilon}(r - \tilde{r}_2 + \varepsilon) & \text{on } [\tilde{r}_2 - \varepsilon, \tilde{r}_2] \\ V\left(h(\tilde{r}_2)\frac{r - \tilde{r}_2}{\varepsilon}\right) & \text{on } [\tilde{r}_2, \tilde{r}_2 + \varepsilon R''] \\ -1 - \frac{\eta}{\varepsilon}(r - \tilde{r}_2 - \varepsilon R'' - \varepsilon) & \text{on } [\tilde{r}_2 + \varepsilon R'', \tilde{r}_2 + \varepsilon R'' + \varepsilon] \\ -1 & \text{on } [\tilde{r}_2 + \varepsilon R'' + \varepsilon, 1], \end{cases}$$

where  $V$  is the unique solution of (3.2) and  $R''$  is the unique value such that  $V(h(\tilde{r}_2)R'') = -1 + \eta$ .

Since  $u_\varepsilon$  is global minimizer, we can estimate the energy of  $J_\varepsilon(\tilde{u}_\varepsilon)$  as follows:

$$J_\varepsilon(u_\varepsilon) \leq J_\varepsilon(\tilde{u}_\varepsilon) \leq J_\varepsilon(u_\varepsilon, B_{r_2 - \varepsilon}(0)) + \varepsilon \omega_{N-1} H(\tilde{r}_2) \beta_1 + B + O(\varepsilon \eta) + o(\varepsilon). \quad (3.16)$$

Then we have a contradiction from (3.15) and (3.16) by taking  $\varepsilon = \varepsilon_j$  and sufficiently large  $j$ .

## References

- [1] S. B. Angenent, J. Mallet-Paret, and L. A. Peletier, Stable transition layers in a semilinear boundary value problem, *J. Differential Equations*, **67** (1987), 212-242.
- [2] Ph. Clément and L. A. Peletier, On a nonlinear eigenvalue problem occurring in population genetics, *Proc. Royal Soc. Edinburg*, **100A**(1985), 85-101.
- [3] Ph. Clément and G. Sweers, Existence of multiplicity results for a semilinear eigenvalue problem, *Ann. Scuola Norm. Sup. Pisa*, **14**(1987), 97-121
- [4] E. N. Dancer, S. Yan, Construction of various type of solutions for an elliptic problem, *Calculus of Variations and Partial Differential Equations* **20**(2004), 93-118.
- [5] D. Gilbarg and N.S. Trudinger, "Elliptic partial differential equations of second order", Springer-Verlag, Berlin, second edition 1983.
- [6] H. Matsuzawa, Stable transition layers in a balanced bistable equation with degeneracy, *Nonlinear Analysis* **58** (2004), 45-67.

- [7] A. S. do. Nascimento, Stable transition layers in a semilinear diffusion equation with spatial inhomogeneities in  $N$ - dimensional domains, J. Differential Equations, **190** (2003), 16-38.
- [8] K. Nakashima, Multi-layered stationary solutions for a spatially inhomogeneous Allen-Cahn equation, J. Differential Equations, **191** (2003), 234-276.
- [9] K. Nakashima, K. Tanaka, Clustering layers and boundary layers in spatially inhomogeneous phase transition problems, Ann. Inst. H. Poincaré Anal. Non Linéaire **20** (2003), 107-143.