

Brauer indecomposability of Scott modules and local subgroups

Hiroki Ishioka
 Department of Mathematics,
 Tokyo University of Science

1. INTRODUCTION

Let p be a prime number and k an algebraically closed field of characteristic p . For a p -subgroup Q of a finite group G and a kG -module M , the Brauer quotient $M(Q)$ of M with respect to Q is naturally a $kN_G(Q)$ -module. A kG -module M is said to be *Brauer indecomposable* if $M(Q)$ is indecomposable or zero as a $kC_G(Q)$ -module for any p -subgroup Q of G ([6]). Brauer indecomposability of p -permutation modules is important for constructing stable equivalences of Morita type between blocks of finite groups (see [2]).

There is a connection between Brauer indecomposability of p -permutation kG -modules and fusion systems, as shown in [6]. The main result in [6] is the following.

Theorem 1 ([6, Theorem 1.1]). *Let P be a p -subgroup of G and M an indecomposable p -permutation kG -module with vertex P . If M is Brauer indecomposable, then $\mathcal{F}_P(G)$ is a saturated fusion system.*

In the case that P is abelian and M is the Scott kG -module $S(G, P)$, it is known that the converse of the above theorem holds.

Theorem 2 ([6, Theorem 1.2]). *Let P be an abelian p -subgroup of G . If $\mathcal{F}_P(G)$ is saturated, then $S(G, P)$ is Brauer indecomposable.*

In general, the above theorem does not hold in the case that P is non-abelian. However, there are some cases in which the Scott kG -module $S(G, P)$ is Brauer indecomposable for non-abelian P (see [5, 7]). Moreover, it was shown that there are some relationships between Brauer indecomposability of Scott modules and fusion systems ([3, 5]). In particular, we proved the following theorem in [3].

Theorem 3 ([3, Theorem 1.3]). *Let G be a finite group and P a p -subgroup of G . Suppose that $M = S(G, P)$ and that $\mathcal{F}_P(G)$ is saturated. Then the following are equivalent.*

- (i) *M is Brauer indecomposable.*
- (ii) *$\text{Res}_{QC_G(Q)}^{N_G(Q)} S(N_G(Q), N_P(Q))$ is indecomposable for each fully normalized subgroup Q of P .*

If these conditions are satisfied, then $M(Q) \cong S(N_G(Q), N_P(Q))$ for each fully normalized subgroup $Q \leq P$.

The above theorem gives a criterion to determine whether the Scott module $S(G, P)$ is Brauer indecomposable.

We investigate the possibility of providing applications of the above theorem. In this paper, we will prove the following result.

Theorem 4. *Let G be a finite group and P a p -subgroup of G . Suppose that $\mathcal{F} := \mathcal{F}_P(G)$ is a saturated fusion system. Consider the following two conditions:*

- (i) $S(N_G(Q), N_P(Q))$ is Brauer indecomposable for each fully \mathcal{F} -normalized subgroup $Q \leq P$.
- (ii) $S(G, P)$ is Brauer indecomposable.

Then (i) implies (ii), and the converse holds if $\mathcal{F} = \mathcal{F}_P(N_G(P))$.

The above theorem shows that there exists some relationship between G and its local subgroups in terms of the Brauer indecomposability of Scott modules, and will be a useful tool for the study of the Brauer indecomposability of Scott modules.

2. PRELIMINARIES

2.1. Scott modules. First, We recall the definition of Scott modules and some of its properties:

Definition 5. For a subgroup H of G , the Scott kG -module $S(G, H)$ with respect to H is the unique indecomposable summand M of $\text{Ind}_H^G k_H$ such that $k_G \mid M$.

If P is a Sylow p -subgroup of H , then $S(G, H)$ is isomorphic to $S(G, P)$. By definition, the Scott kG -module $S(G, P)$ is a p -permutation kG -module.

By Green's indecomposability criterion, the following result holds.

Lemma 6. *Let H be a subgroup of G such that $|G : H| = p^a$ (for some $a \geq 0$). Then $\text{Ind}_H^G k_H$ is indecomposable. In particular, we have that*

$$S(G, H) \cong \text{Ind}_H^G k_H.$$

The following theorem gives us information of restrictions of Scott modules.

Theorem 7 ([4, Theorem 1.7]). *Let P be a p -subgroup of H . Let Q be a maximal element of $P \cap_G H = \{gP \cap H \mid g \in G\}$. Then $S(H, Q)$ is a direct summand of $\text{Res}_H^G S(G, P)$.*

2.2. Brauer quotients. Let M be a kG -module and H a subgroup of G . We denote by M^H the set of H -fixed elements in M . For subgroups L of H , we denote by Tr_L^H the trace map $\text{Tr}_L^H : M^L \rightarrow M^H$. Brauer quotients are defined as follows.

Definition 8. Let M be a kG -module. For a p -subgroup Q of G , the Brauer quotient of M with respect to Q is the k -vector space

$$M(Q) := M^Q / \left(\sum_{R < Q} \text{Tr}_R^Q(M^R) \right).$$

This k -vector space has a natural structure of $kN_G(Q)$ -module.

Brauer quotients have the following well-known properties.

Proposition 9. *Let P be a p -subgroup of G and $M = S(G, P)$. Then $M(P) \cong S(N_G(P), P)$.*

Proposition 10. *Let M be an indecomposable p -permutation kG -module with vertex P . Let Q be a p -subgroup of G . Then $Q \leq_G P$ if and only if $M(Q) \neq 0$.*

2.3. Fusion systems. For subgroups Q, R of G , we denote by $\text{Hom}_G(Q, R)$ the set of all group homomorphisms from Q to R which are induced by conjugation in G . For a p -subgroup P of G , the *fusion system* $\mathcal{F}_P(G)$ of G over P is the category whose objects are the subgroups of P and whose morphism set from Q to R is $\text{Hom}_G(Q, R)$. We refer the reader to [1] for background involving fusion systems.

Definition 11. Let P be a p -subgroup of G

- (i) A subgroup Q of P is said to be *fully normalized* in $\mathcal{F}_P(G)$ if $|N_P({}^xQ)| \leq |N_P(Q)|$ for all $x \in G$ such that ${}^xQ \leq P$.
- (ii) A subgroup Q of P is said to be *fully automized* in $\mathcal{F}_P(G)$ if $p \nmid |N_G(Q) : N_P(Q)C_G(Q)|$.
- (iii) A subgroup Q of P is said to be *receptive* in $\mathcal{F}_P(G)$ if it has the following property: for each $R \leq P$ and $\varphi \in \text{Iso}_{\mathcal{F}_P(G)}(R, Q)$, if we set

$$N_\varphi := \{g \in N_P(Q) \mid \exists h \in N_P(R), c_g \circ \varphi = \varphi \circ c_h\},$$

then there is $\bar{\varphi} \in \text{Hom}_{\mathcal{F}_P(G)}(N_\varphi, P)$ such that $\bar{\varphi}|_{R=} = \varphi$.

Saturated fusion systems are defined as follows.

Definition 12. Let P be a p -subgroup of G . The fusion system $\mathcal{F}_P(G)$ is *saturated* if the following two conditions are satisfied:

- (i) P is fully normalized in $\mathcal{F}_P(G)$.
- (ii) For each subgroup Q of P , if Q is fully normalized in $\mathcal{F}_P(G)$, then Q is receptive in $\mathcal{F}_P(G)$.

For example, if P is a Sylow p -subgroup of G , then $\mathcal{F}_P(G)$ is saturated.

3. PROOF OF THEOREM 4

In this section, we give a proof of Theorem 4.

For a saturated fusion system \mathcal{F} over p -group P and a subgroup Q of P , the normalizer fusion system $N_{\mathcal{F}}(Q)$ of Q is defined and is a fusion system over $N_P(Q)$ (see [1, II, §2]). We note that if $\mathcal{F} = \mathcal{F}_P(G)$, then $N_{\mathcal{F}}(Q) = \mathcal{F}_{N_P(Q)}(N_G(Q))$.

Proof of Theorem 4. Suppose that (i) holds. Let Q be a fully \mathcal{F} -normalized subgroup of P . Then $S(N_G(Q), N_P(Q))(Q)$ is indecomposable, and we have that

$$S(N_G(Q), N_P(Q)) \cong S(N_G(Q), N_P(Q))(Q).$$

Therefore, $S(G, P)$ is Brauer indecomposable by Theorem 3.

Next, suppose that (ii) and $\mathcal{F} = \mathcal{F}_P(N_G(P))$ hold. Then any subgroup Q of P is fully \mathcal{F} -normalized. Let Q be any subgroup of P . Then $\mathcal{F}_{N_P(Q)}(N_G(Q)) = N_{\mathcal{F}}(Q)$ is saturated by [1, II, Theorem 2.1]. Let R be a fully $N_{\mathcal{F}}(Q)$ -normalized subgroup of $N_P(Q)$. It is sufficient to show that $S(N_{N_G(Q)}(R), N_{N_P(Q)}(R))$ is indecomposable as $kC_{N_G(Q)}(R)$ -module by Theorem 3.

Since QR is fully \mathcal{F} -normalized, $S(N_G(QR), N_P(QR))$ is indecomposable as $kC_G(QR)$ -module, and hence is also indecomposable as $kC_{N_G(Q)}(R)$ -module. Therefore, it is sufficient to show that

$$\text{Res}_{N_{N_G(Q)}(R)}^{N_G(QR)} S(N_G(QR), N_P(QR)) \cong S(N_{N_G(Q)}(R), N_{N_P(Q)}(R)),$$

and if we show that $N_{N_P(Q)}(R)$ is a maximal element of $N_P(QR) \cap_{N_G(QR)} N_{N_G(Q)}(R)$, then the isomorphism holds by Theorem 7 and the indecomposability of $S(N_G(QR), N_P(QR))$ as a $N_{N_G(Q)}(R)$ -module.

Let g be an element of $N_G(QR)$ such that $N_{N_P(Q)}(R) \leq {}^g N_P(QR) \cap N_{N_G(Q)}(R)$. Then we have $Q^g \leq (QR)^g = QR \leq P$ and hence there is $h \in N_G(P)$ such that $gh^{-1} \in C_G(Q)$ since $\mathcal{F} = \mathcal{F}_P(N_G(P))$. We have that

$$\begin{aligned} |N_{N_P(Q)}(R)| &\leq |{}^g N_P(QR) \cap N_{N_G(Q)}(R)| \\ &= |{}^g P \cap N_G(QR) \cap N_G(Q) \cap N_G(R)| \\ &= |{}^g P \cap N_G(Q) \cap N_G(R)| \\ &= |P \cap N_G(Q^g) \cap N_G(R^g)| \\ &= |N_{N_P(Q^g)}(R^g)| \\ &= |N_{N_P(Q^h)}(R^g)| \\ &= |N_{N_P(Q)^h}(R^g)| \\ &= |N_{N_P(Q)}(R^{gh^{-1}})^h| \\ &= |N_{N_P(Q)}(R^{gh^{-1}})|. \end{aligned}$$

On the other hand, since

$$\begin{aligned} R^{gh^{-1}} &\leq N_{N_P(Q)}(R)^{gh^{-1}} \\ &\leq ({}^g N_P(QR) \cap N_{N_G(Q)}(R))^{gh^{-1}} \\ &\leq ({}^g P \cap N_G(Q))^{gh^{-1}} \\ &= P^{h^{-1}} \cap N_G(Q^{gh^{-1}}) \\ &= P \cap N_G(Q) \\ &= N_P(Q) \end{aligned}$$

and $gh^{-1} \in C_G(Q) \leq N_G(Q)$, the conjugation map $(\)^{gh^{-1}}: R \rightarrow R^{gh^{-1}}$ is an isomorphism in $N_{\mathcal{F}}(Q)$. Since R is fully $N_{\mathcal{F}}(Q)$ -normalized, we have that $|N_{N_P(Q)}(R^{gh^{-1}})| \leq |N_{N_P(Q)}(R)|$. Therefore, $N_{N_P(Q)}(R) = {}^g N_P(QR) \cap N_{N_G(Q)}(R)$, and $N_{N_P(Q)}(R)$ is maximal in $N_P(QR) \cap_{N_G(QR)} N_{N_G(Q)}(R)$, as desired. \square

REFERENCES

- [1] M. Aschbacher, R. Kessar, and B. Oliver, *Fusion systems in algebra and topology*, London Mathematical Society Lecture Note Series, vol. 391, Cambridge University Press, Cambridge, 2011.
- [2] M. Broué, *Equivalences of blocks of group algebras*, Finite-dimensional algebras and related topics (Ottawa, ON, 1992), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 424, Kluwer Acad. Publ., Dordrecht, 1994, pp. 1–26.
- [3] H. Ishioka and N. Kunugi, *Brauer indecomposability of Scott modules*, J. Algebra **470** (2017), 441–449.
- [4] H. Kawai, *On indecomposable modules and blocks*, Osaka J. Math. **23** (1986), 201–205.
- [5] R. Kessar, S. Koshitani, and M. Linckelmann, *On the Brauer indecomposability of Scott modules*, Q. J. Math. **66** (2015), 895–903.
- [6] R. Kessar, N. Kunugi, and N. Mitsuhashi, *On saturated fusion systems and Brauer indecomposability of Scott modules*, J. Algebra **340** (2011), 90–103.

- [7] Í. Tuvay, *On Brauer indecomposability of Scott modules of Park-type groups*, J. Group Theory **17** (2014), 1071–1079.

DEPARTMENT OF MATHEMATICS
TOKYO UNIVERSITY OF SCIENCE
1-3 KAGURAZAKA, SHINJUKU-KU, TOKYO 162-8601
JAPAN

E-mail address: 114701@ed.tus.ac.jp

東京理科大学・理学研究科 石岡 大樹