

On sharpness of the Yang-Yau inequality for the genus two case

Toshihiro Shoda *

Faculty of Education,
Saga University

Abstract

This is a survey article for our recent work related to the sharp upper bound of the Yang-Yau inequality for the genus two case.

1 Introduction

Let M be an orientable closed surface. For a Riemannian metric $ds^2 = g_{ij}dx^i dx^j$ on M , the Laplacian is defined as

$$\Delta_{ds^2} = \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x^i} \left(\sqrt{|g|} g^{ij} \frac{\partial}{\partial x^j} \right) \quad ((g^{ij}) = (g_{kl})^{-1}, |g| = \det(g_{ij})).$$

Let

$$\Lambda(ds^2) := \lambda_1(ds^2) \cdot \text{area}(ds^2),$$

where $\lambda_1(ds^2)$ is the first positive eigenvalue of Δ_{ds^2} and $\text{area}(ds^2)$ is the area of M with respect to ds^2 . The following are well-known results related to the upper bound of $\Lambda(ds^2)$.

Theorem 1.1. (i) (Hersch [Her]) *For any metric ds^2 on the sphere S^2 , $\Lambda(ds^2) \leq 8\pi$ holds.*
(ii) (Nadirashvili [N]) *For any metric ds^2 on the torus, $\Lambda(ds^2) \leq 8\pi^2/\sqrt{3}$*

*The author was partly supported by JSPS Grant-in-Aid for Scientific Research (C) 16K05134. This work is jointed with Shin Nayatani, Nagoya University.

holds.

(iii) (Yang-Yau [YY]) *If γ is the genus of M , then for any metric ds^2 on M , we have*

$$\Lambda(ds^2) \leq 8\pi \cdot \left\lfloor \frac{\gamma + 3}{2} \right\rfloor. \quad (1)$$

The inequality of (i) is sharp since the equality is attained for the standard metric on S^2 , and thus the inequality (1) is sharp for $\gamma = 0$. On the other hand, by (ii), the inequality (1) is not sharp when $\gamma = 1$.

For $\gamma = 2$, the inequality (1) implies $\Lambda(ds^2) \leq 16\pi$, and Jakobson-Levitin-Nadirashvili-Nigam-Polterovich [JLNNP] conjectured the following:

Conjecture (Jakobson et al. [JLNNP]). Let B be the closed Riemann surface of genus two defined by

$$B = \{(z, w) \in \mathbb{C}^2 \mid w^2 = z(z^4 + 1)\} \cup \{(\infty, \infty)\},$$

called the *Bolza surface*. Then, there exists a singular metric ds_B^2 on B such that $\lambda_1(ds_B^2) = 2$, that is, $\Lambda(ds_B^2) = 16\pi$ holds.

Note that ds_B^2 can be approximated by smooth metrics, which remarked in [JLNNP], and hence the inequality (1) is sharp for $\gamma = 2$.

To see the conjecture, they considered the following deformation of B . For $0 < \theta < \pi/2$, let B_θ be the Riemann surface of genus two defined by

$$B_\theta = \{(z, w) \in \mathbb{C}^2 \mid w^2 = z(z^4 + 2 \cos 2\theta \cdot z^2 + 1)\} \cup \{(\infty, \infty)\}.$$

Note that $B_{\pi/4} = B$. Let ds_θ^2 denote the pull-back of the standard metric on $S^2 = \overline{\mathbb{C}}$ by the meromorphic function $g_\theta: B_\theta \ni (z, w) \mapsto z \in \overline{\mathbb{C}}$. Then $ds_{\pi/4}^2$ is the desired singular metric ds_B^2 in the conjecture. They used numerical arguments, and thus their result has been treated as conjecture as above. For this open problem, our recent paper [NS] gives an affirmative answer.

Main Theorem (Nayatani and Shoda, 2019). There exists $\theta_1 \approx 0.65$ so that for $\theta_1 \leq \theta \leq \pi/2 - \theta_1$, we have $\lambda_1(ds_{B_\theta}^2) = 2$ and therefore $\Lambda(ds_\theta^2) = 16\pi$.

2 Index and nullity of a holomorphic map

In this section, we first review the arguments in [EK] and [MR], and translate the conjecture into their situations.

Let $g: M \rightarrow \overline{\mathbb{C}}$ be a non-constant meromorphic function on a compact Riemann surface M , that is, a holomorphic map from M to S^2 . Let ds_g^2 be the pull-back metric of the standard metric on $\overline{\mathbb{C}} = S^2$. Note that ds_g^2 is a singular metric, and we can consider the Laplacian Δ_g with respect to ds_g^2 and its eigenvalues in a standard way. The *index* of g (resp. the *nullity* of g) is defined as the number of eigenvalues of Δ_g less than 2 counted with multiplicities (resp. the multiplicity of eigenvalue 2 of Δ_g). Let $\text{ind}(g)$ (resp. $\text{nul}(g)$) be the index of g (resp. the nullity of g).

We now consider the image of g in $S^2 \subset \mathbb{R}^3$. For a constant vector $a \in \mathbb{R}^3$, it is well-known that $\langle a, g \rangle$ gives an eigenfunction of Δ_g of eigenvalue 2, where $\langle *, * \rangle$ is the standard inner product on \mathbb{R}^3 . Hence we have $\text{nul}(g) \geq 3$. Eigenfunctions of Δ_g of eigenvalue 2 other than $\langle a, g \rangle$ -type are called *extra eigenfunctions*. Ejiri-Kotani [EK] and Montiel-Ros [MR] obtained the procedure to determine extra eigenfunctions, which omitted in this report.

Since every constant function is a single eigenfunction of Δ_g of eigenvalue 0, we have $\text{ind}(g) \geq 1$. The conjecture states that, for $g = g_B$, the second least eigenvalue of Δ_{g_B} should equal 2, and so it is equivalent to $\text{ind}(g_B) = 1$.

3 Outline of proof

In this section, we observe behaviors of eigenvalues of Δ_{g_θ} less than 2 as $\theta: 0 \rightarrow \pi/2$.

By the Montiel-Ros result [MR], we have $\text{Ind}(g_\theta) = 3$ for θ which is sufficiently near to 0. Thus there exists a single eigenvalue 0 and there exist two positive eigenvalues of Δ_{g_θ} less than 2. We next see

Lemma 3.1. *Set*

$$A = \int_0^\infty \frac{dt}{\sqrt{t(t^4 + 2 \cos 2\theta \cdot t^2 + 1)}}, \quad B = \int_0^\infty \frac{dt}{\sqrt{t(t^4 - 2 \cos 2\theta \cdot t^2 + 1)}},$$

$$C = \int_0^\infty \frac{t^3 dt}{\sqrt{t(t^4 + 2 \cos 2\theta \cdot t^2 + 1)}^3}, \quad D = \int_0^\infty \frac{t^3 dt}{\sqrt{t(t^4 - 2 \cos 2\theta \cdot t^2 + 1)}^3}.$$

Let $\theta_1 (\approx 0.65)$ be the unique solution of

$$A(B^2 + 16D^2 \sin^2 2\theta) + 8(AD + BC)(B \cos 2\theta - 4D \sin^2 2\theta) = 0, \quad (2)$$




Figure 1: A rough sketch of behaviors of eigenvalues of Δ_{g_θ} less than 2

and set $\theta_2 = \pi/2 - \theta_1 (\approx 0.91)$. Then there exist two extra eigenfunctions for $\theta = \theta_1, \theta_2$. Therefore we have

$$\text{Nul}(g_\theta) = \begin{cases} 5, & \theta = \theta_1, \theta_2, \\ 3, & \theta \neq \theta_1, \theta_2. \end{cases}$$

By Lemma 3.1, we can find two eigenvalues 2 of $\Delta_{g_{\theta_i}}$ ($i = 1, 2$) other than three eigenvalues 2 given by $\langle a, g \rangle$ -type. By symmetries of the extra eigenfunctions, we can show that their two eigenvalues are monotonically increasing (resp. decreasing) when they pass through $\theta = \theta_1$ (resp. $\theta = \theta_2$). Hence we can see the behaviors of eigenvalues of Δ_{g_θ} less than 2 as $\theta: 0 \rightarrow \pi/2$ (see Figure 1).

Therefore, we have the following theorem which implies our main result:

Theorem 3.2.

$$\text{ind}(g_\theta) = \begin{cases} 3, & 0 < \theta < \theta_1, \\ 1, & \theta_1 \leq \theta \leq \theta_2, \\ 3, & \theta_2 < \theta < \pi/2. \end{cases}$$

4 Appendix

In this section, we will prove that there exist no solutions of (2) for $\pi/4 \leq \theta < \pi/2$, which omitted in the original paper. In fact, the equation seems to be even more complicated than indicated, since the numbers A, B, C, D

depend in a non-trivial way on θ . So it would be helpful for the readers to give arguments for the claim even if it consists of long calculations.

We first rewrite A, B, C, D by the complete elliptic integrals. Recall that, for $0 < k < 1$, Let

$$K(k) = \int_0^{\frac{\pi}{2}} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}, \quad E(k) = \int_0^{\frac{\pi}{2}} \sqrt{1 - k^2 \sin^2 \theta} d\theta.$$

$K(k)$ is called the *complete elliptic integral of first kind*, and $E(k)$ the *complete elliptic integral of second kind*.

By 222 in [BF], we have

$$\begin{aligned} A &= \frac{1}{2} \int_0^\infty \frac{dt}{\sqrt{t(t^4 + 2\cos(2\theta)t^2 + 1)}} + \frac{1}{2} \int_0^\infty \frac{t dt}{\sqrt{t(t^4 + 2\cos(2\theta)t^2 + 1)}} \\ &\stackrel{x=\sqrt{t}-1/\sqrt{t}}{=} \int_{-\infty}^\infty \frac{dx}{\sqrt{x^4 + 4x^2 + 2 + 2\cos 2\theta}} = 2 \int_0^\infty \frac{dx}{\sqrt{x^4 + 4x^2 + 2 + 2\cos 2\theta}} \\ &= \frac{2}{\sqrt{2(1 + \sin \theta)}} K\left(\sqrt{\frac{2 \sin \theta}{1 + \sin \theta}}\right), \\ B &= 2 \int_0^\infty \frac{dx}{\sqrt{x^4 + 4x^2 + 2 - 2\cos 2\theta}} = \frac{2}{\sqrt{2(1 + \cos \theta)}} K\left(\sqrt{\frac{2 \cos \theta}{1 + \cos \theta}}\right), \\ C &= 2 \int_0^\infty \frac{dx}{\sqrt{x^4 + 4x^2 + 2 + 2\cos 2\theta}} \\ &= \frac{1}{4\sqrt{2(1 + \sin \theta)} \sin^2 \theta (1 - \sin \theta)} \\ &\quad \times \left(E\left(\sqrt{\frac{2 \sin \theta}{1 + \sin \theta}}\right) - (1 - \sin \theta) K\left(\sqrt{\frac{2 \sin \theta}{1 + \sin \theta}}\right) \right), \\ D &= 2 \int_0^\infty \frac{dx}{\sqrt{x^4 + 4x^2 + 2 - 2\cos 2\theta}} \\ &= \frac{1}{4\sqrt{2(1 + \cos \theta)} \cos^2 \theta (1 - \cos \theta)} \\ &\quad \times \left(E\left(\sqrt{\frac{2 \cos \theta}{1 + \cos \theta}}\right) - (1 - \cos \theta) K\left(\sqrt{\frac{2 \cos \theta}{1 + \cos \theta}}\right) \right). \end{aligned}$$

(2) is equivalent to

$$\cos 2\theta(AB^2 \cos 2\theta + 8ABD + 8B^2C) + \sin^2 2\theta(AB^2 - 16AD^2 - 32BCD) = 0.$$

Thus, for $\pi/4 \leq \theta < \pi/2$, it is sufficient to show

$$AB^2 \cos 2\theta + 8ABD + 8B^2C > 0, \quad AB^2 - 16AD^2 - 32BCD < 0,$$

since this claim implies that the left hand side of (2) is negative for $\pi/4 \leq \theta < \pi/2$.

For simplicity, we set $k^2 = 2 \sin \theta / (1 + \sin \theta)$, $l^2 = 2 \cos \theta / (1 + \cos \theta)$, and we obtain

$$\begin{aligned} \sin \theta &= \frac{k^2}{2 - k^2}, & 1 - \sin \theta &= \frac{2(1 - k^2)}{2 - k^2}, & \cos \theta &= \frac{l^2}{2 - l^2}, \\ 1 - \cos \theta &= \frac{2(1 - l^2)}{2 - l^2}, & k^4 l^4 &= 16(1 - k^2)(1 - l^2). \end{aligned}$$

4.1 Proof of $AB^2 \cos 2\theta + 8ABD + 8B^2C > 0$

The inequality $AB^2 \cos 2\theta + 8ABD + 8B^2C > 0$ follows from the next lemma:

Lemma 4.1. *For $\pi/4 \leq \theta < \pi/2$, we have $A \cos 2\theta + 8C > 0$.*

Proof. We first obtain

$$\begin{aligned} A \cos 2\theta + 8C &= \frac{2}{\sqrt{2(1 + \sin \theta)} \sin^2 \theta (1 - \sin \theta)} \\ &\quad \times (E(k) + (\cos 2\theta \sin^2 \theta - 1)(1 - \sin \theta)K(k)). \end{aligned}$$

Next, we find

$$\frac{\pi}{4} \leq \theta < \frac{\pi}{2} \iff 2(\sqrt{2} - 1) \leq k^2 < 1,$$

and the inequality $A \cos 2\theta + 8C > 0$ is equivalent to

$$E(k) + \frac{4(-k^8 + 2k^6 - 10k^4 + 16k^2 - 8)}{(2 - k^2)^5} \cdot (1 - k^2)K(k) > 0.$$

By **710.02**, **710.04** in [BF], we find

$$\frac{d}{dk} E(k) = \frac{E(k) - K(k)}{k}, \quad \frac{d}{dk} ((1 - k^2)K(k)) = \frac{dE(k)}{dk} - kK(k).$$

Thus we have

$$\begin{aligned}
& \frac{d}{dk} \left(E(k) + \frac{4(-k^8 + 2k^6 - 10k^4 + 16k^2 - 8)}{(2 - k^2)^5} \cdot (1 - k^2)K(k) \right) \\
&= \frac{E(k) - K(k)}{k} - \frac{8k(k^8 + 4k^6 + 18k^4 - 24k^2 + 8)}{(2 - k^2)^6} \cdot (1 - k^2)K(k) \\
&\quad + \frac{4(-k^8 + 2k^6 - 10k^4 + 16k^2 - 8)}{(2 - k^2)^5} \left(\frac{E(k) - K(k)}{k} - kK(k) \right) \\
&= \frac{k(k^8 - 6k^6 + 32k^4 - 40k^2 + 16)}{(2 - k^2)^5} (K(k) - E(k)) \\
&\quad + \frac{4k^3(k^4 + 4k^2 - 4)(k^4 + 6k^2 - 6)}{(2 - k^2)^6} K(k) \\
&\left(= \frac{k(k^8 + 6k^4(1 - k^2) + k^4 + (5k^2 - 4)^2)}{(2 - k^2)^5} (K(k) - E(k)) \right. \\
&\quad \left. + \frac{4k^3(k^4 + 4k^2 - 4)(k^4 + 6k^2 - 6)}{(2 - k^2)^6} K(k) \right) \\
&= \frac{k}{(2 - k^2)^6} ((2 - k^2)(k^8 - 6k^6 + 32k^4 - 40k^2 + 16)(K(k) - E(k)) \\
&\quad + 4k^2(k^4 + 4k^2 - 4)(k^4 + 6k^2 - 6)K(k)).
\end{aligned}$$

Hence, for $\sqrt{15} - 3 \leq k^2 < 1$, this is automatically positive. For $2(\sqrt{2} - 1) \leq k^2 < \sqrt{15} - 3$, we obtain

$$\begin{aligned}
& (2 - k^2)(k^8 - 6k^6 + 32k^4 - 40k^2 + 16)(K(k) - E(k)) \\
&\quad + 4k^2(k^4 + 4k^2 - 4)(k^4 + 6k^2 - 6)K(k) \\
&\geq \lim_{k^2 \rightarrow 2(\sqrt{2}-1)} (2 - k^2)(k^8 - 6k^6 + 32k^4 - 40k^2 + 16)(K(k) - E(k)) \\
&\quad - 0.1 \cdot K \left(\sqrt{\sqrt{15} - 3} \right) \\
&> (7424 - 5248\sqrt{2})(K(\sqrt{0.82}) - E(\sqrt{0.82})) - 0.1 \cdot K(\sqrt{0.88}) \\
&> 2.2(2.305232 - 1.164798) - 0.1 \cdot 2.492635 = 2.25969 > 0
\end{aligned}$$

(see pp. 324 in [BF]). In fact, by setting $f(x) = (2 - x)(x^4 - 6x^3 + 32x^2 - 40x + 16)$, $f(x)$ is monotonically increasing for $2(\sqrt{2}-1) \leq x < 1$. By **710.05** in [BF], we have

$$\frac{d}{dk}(K(k) - E(k)) = \frac{kE(k)}{1 - k^2} > 0,$$

and thus $K(k) - E(k)$ is monotonically increasing.

Hence, for $2(\sqrt{2} - 1) \leq k^2 < 1$, we find

$$\frac{d}{dk} \left(E(k) + \frac{4(-k^8 + 2k^6 - 10k^4 + 16k^2 - 8)}{(2 - k^2)^5} \cdot (1 - k^2)K(k) \right) > 0.$$

It follows that, for $2(\sqrt{2} - 1) \leq k^2 < 1$,

$$\begin{aligned} & E(k) + \frac{4(-k^8 + 2k^6 - 10k^4 + 16k^2 - 8)}{(2 - k^2)^5} \cdot (1 - k^2)K(k) \\ & \geq E\left(\sqrt{2(\sqrt{2} - 1)}\right) - \frac{\sqrt{2} - 1}{\sqrt{2}} K\left(\sqrt{2(\sqrt{2} - 1)}\right) \\ & > E(\sqrt{0.82}) - 0.3K(\sqrt{0.83}) = 1.164798 - 0.3 \cdot 2.331409 = 0.465375 > 0. \end{aligned}$$

(see again pp.324 in [BF]). \square

4.2 Proof of $AB^2 - 16AD^2 - 32BCD < 0$

Since

$$\begin{aligned} & AB^2 - 16AD^2 - 32BCD \\ &= \begin{cases} A(B - \frac{192}{25}D)(B + \frac{25}{12}D) + BD(\frac{1679}{300}A - 32C) & (\pi/4 \leq \theta \leq 5\pi/16) \\ A(B - 10D)(B + \frac{8}{5}D) + BD(\frac{42}{5}A - 32C) & (5\pi/16 \leq \theta \leq 3\pi/8) \\ A(B - 16D)(B + D) + BD(15A - 32C) & (3\pi/8 \leq \theta < \pi/2), \end{cases} \end{aligned}$$

it is sufficient to prove

$$25B - 192D < 0 \quad (\pi/4 \leq \theta \leq 5\pi/16), \quad (3)$$

$$1679A - 9600C < 0 \quad (\pi/4 \leq \theta \leq 5\pi/16), \quad (4)$$

$$B - 10D < 0 \quad (5\pi/16 \leq \theta \leq 3\pi/8), \quad (5)$$

$$21A - 80C < 0 \quad (5\pi/16 \leq \theta \leq 3\pi/8), \quad (6)$$

$$B - 16D < 0 \quad (3\pi/8 \leq \theta < \pi/2), \quad (7)$$

$$15A - 32C < 0 \quad (3\pi/8 \leq \theta < \pi/2) \quad (8)$$

to show $AB^2 - 16AD^2 - 32BCD < 0$.

Proof of (3) Straightforward calculations yield

$$\begin{aligned} 25B - 192D &= \frac{2}{\sqrt{2(1 + \cos \theta)} \cos^2 \theta (1 - \cos \theta)} \\ &\times (25 \cos^2 \theta (1 - \cos \theta) K(l) - 24(E(l) - (1 - \cos \theta) K(l))), \end{aligned}$$

$$\begin{aligned} & 25 \cos^2 \theta (1 - \cos \theta) K(l) - 24(E(l) - (1 - \cos \theta)K(l)) \\ &= \frac{2}{(2 - l^2)^3} ((49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3 E(l)). \end{aligned}$$

Thus, we will prove

$$(49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3 E(l) < 0$$

for

$$0.714298 \approx \frac{2 \cos \frac{5}{16}\pi}{1 + \cos \frac{5}{16}\pi} \leq l^2 \leq \frac{2 \cos \frac{\pi}{4}}{1 + \cos \frac{\pi}{4}} \approx 0.828427.$$

We first obtain

$$\begin{aligned} & \frac{d}{dl} ((49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3 E(l)) \\ &= (196l^3 - 192l)(1 - l^2)K(l) + (49l^4 - 96l^2 + 96) \left(\frac{E - K}{l} - lK \right) \\ &\quad + 72l(2 - l^2)^2 E(l) - 12(2 - l^2)^3 \cdot \frac{E - K}{l} \\ &= l(-(257l^4 - 507l^2 + 336)K(l) + (84l^4 - 311l^2 + 336)E(l)). \end{aligned}$$

For $0.71 \leq l^2 \leq 0.83$, $K(l)$ is positive and monotonically increasing, and $257l^4 - 507l^2 + 336$, $84l^4 - 311l^2 + 336$, $E(l)$ are positive and monotonically decreasing. We now divide the interval $0.71 \leq l^2 \leq 0.83$ into $0.71 \leq l^2 \leq 0.81$ and $0.81 \leq l^2 \leq 0.83$, and estimate separately. For $0.71 \leq l^2 \leq 0.81$, we find

$$\begin{aligned} & -(257l^4 - 507l^2 + 336)K(l) + (84l^4 - 311l^2 + 336)E(l) \\ & \leq -(257 \cdot 0.81^2 - 507 \cdot 0.81 + 336)K(\sqrt{0.71}) \\ & \quad + (84 \cdot 0.71^2 - 311 \cdot 0.71 + 336)E(\sqrt{0.71}) \\ & \approx -1.723 < 0. \end{aligned}$$

For $0.81 \leq l^2 \leq 0.83$,

$$\begin{aligned} & -(257l^4 - 507l^2 + 336)K(l) + (84l^4 - 311l^2 + 336)E(l) \\ & \leq -(257 \cdot 0.83^2 - 507 \cdot 0.83 + 336)K(\sqrt{0.81}) \\ & \quad + (84 \cdot 0.81^2 - 311 \cdot 0.81 + 336)E(\sqrt{0.81}) \\ & \approx -47.2487 < 0. \end{aligned}$$

Hence, for $0.71 \leq l^2 \leq 0.83$, we have

$$\frac{d}{dl}((49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3E(l)) < 0.$$

Thus

$$(49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3E(l)$$

is monotonically decreasing. Since $K(l)$ is positive and monotonically increasing, and $E(l)$ is positive and monotonically decreasing, we have

$$\begin{aligned} & (49l^4 - 96l^2 + 96)(1 - l^2)K(l) - 12(2 - l^2)^3E(l) \\ & \leq (49 \cdot 0.714^2 - 96 \cdot 0.714 + 96)(1 - 0.714)K(\sqrt{0.714}) \\ & \quad - 12(2 - 0.714)^3E(\sqrt{0.714}) \\ & \approx -0.0337104 < 0. \end{aligned}$$

Therefore the claim follows.

Proof of (4) We first find

$$\begin{aligned} 1679A - 9600C &= \frac{2}{\sqrt{2(1 + \sin \theta)} \sin^2 \theta (1 - \sin \theta)} \\ &\times (1679 \sin^2 \theta (1 - \sin \theta)K(k) - 1200(E(k) - (1 - \sin \theta)K(k))), \\ 1679 \sin^2 \theta (1 - \sin \theta)K(k) - 1200(E(k) - (1 - \sin \theta)K(k)) \\ &= \frac{2}{(2 - k^2)^3} ((2879k^4 - 4800k^2 + 4800)(1 - k^2)K(k) - 600(2 - k^2)^3E(k)). \end{aligned}$$

Note that

$$0.828427 \approx \frac{2 \sin \frac{\pi}{4}}{1 + \sin \frac{\pi}{4}} \leq k^2 < 1.$$

Direct calculation yields

$$\begin{aligned} & \frac{d}{dk}((2879k^4 - 4800k^2 + 4800)(1 - k^2)K(k) - 600(2 - k^2)^3E(k)) \\ &= (4 \cdot 2879k^3 - 9600k)(1 - k^2)K(k) \\ & \quad + (2879k^4 - 4800k^2 + 4800) \left(\frac{E - K}{k} - kK \right) \\ & \quad + 3600k(2 - k^2)^2E(k) - 600(2 - k^2)^3 \cdot \frac{E - K}{k} \\ &= k(-(14995k^4 - 26637k^2 + 16800)K(k) \\ & \quad + (4200k^4 - 15121k^2 + 16800)E(k)). \end{aligned}$$

We can check that, for $0.82 \leq k^2 < 1$, $14995x^2 - 26637x + 16800$ is a positive function which attains the minimum at $x = 26637/(2 \cdot 14995)$ and the maximum at $x = 1$, and $4200x^2 - 15121x + 16800$ is positive and monotonically decreasing. Hence we have

$$\begin{aligned} & - (14995k^4 - 26637k^2 + 16800)K(k) + (4200k^4 - 15121k^2 + 16800)E(k) \\ & \leq - \left(14995 \cdot \left(\frac{26637}{2 \cdot 14995} \right)^2 - 26637 \cdot \frac{26637}{2 \cdot 14995} + 16800 \right) K(\sqrt{0.82}) \\ & \quad + (4200 \cdot 0.82^2 - 15121 \cdot 0.82 + 16800)E(\sqrt{0.82}) \\ & \approx -3042.79 < 0. \end{aligned}$$

Thus, $(2879k^4 - 4800k^2 + 4800)(1 - k^2)K(k) - 600(2 - k^2)^3E(k)$ is monotonically decreasing. We finally obtain

$$\begin{aligned} & (2879k^4 - 4800k^2 + 4800)(1 - k^2)K(k) - 600(2 - k^2)^3E(k) \\ & \leq (2879 \cdot 0.8284^2 - 4800 \cdot 0.8284 + 4800)(1 - 0.8284)K(\sqrt{0.8284}) \\ & \quad - 600(2 - 0.8284)^3E(\sqrt{0.8284}) \\ & \leq (2879 \cdot 0.8284^2 - 4800 \cdot 0.8284 + 4800)(1 - 0.8284)K(\sqrt{0.8284}) \\ & \quad - 600(2 - 0.8284)^3E(\sqrt{0.8284}) \\ & \approx -0.371741 < 0. \end{aligned}$$

Therefore the claim is proved.

Proof of (5). We first have

$$\begin{aligned} B - 10D &= \frac{1}{2\sqrt{2(1 + \cos \theta)} \cos^2 \theta (1 - \cos \theta)} \\ &\quad \times (2 \cos^2 \theta (1 - \cos \theta)K(l) - 5(E(l) - (1 - \cos \theta)K(l))), \\ 2 \cos^2 \theta (1 - \cos \theta)K(l) - 5(E(l) - (1 - \cos \theta)K(l)) \\ &= \frac{1}{(2 - l^2)^3} (2(7l^4 - 20l^2 + 20)(1 - l^2)K(l) - 5(2 - l^2)^3E(l)), \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dl} (2(7l^4 - 20l^2 + 20)(1 - l^2)K(l) - 5(2 - l^2)^3E(l)) \\ = 2(28l^3 - 40l)(1 - l^2)K(l) + 2(7l^4 - 20l^2 + 20) \left(\frac{E - K}{l} - lK \right) \end{aligned}$$

$$\begin{aligned}
& + 30l(2 - l^2)^2 E(l) - 5(2 - l^2)^3 \cdot \frac{E - K}{l} \\
& = l(-(75l^4 - 192l^2 + 140)K(l) + (35l^4 - 136l^2 + 140)E(l)).
\end{aligned}$$

Note that

$$0.553537 \approx \frac{2 \cos \frac{3}{8}\pi}{1 + \cos \frac{3}{8}\pi} \leq l^2 \leq \frac{2 \cos \frac{5}{16}\pi}{1 + \cos \frac{5}{16}\pi} \approx 0.714298,$$

and $75l^4 - 192l^2 + 140$, $35l^4 - 136l^2 + 140$ are positive and monotonically decreasing for $0.55 \leq l^2 \leq 0.715$.

We now divide the interval $0.55 \leq l^2 \leq 0.715$ among $0.55 \leq l^2 \leq 0.58$, $0.58 \leq l^2 \leq 0.62$, $0.62 \leq l^2 \leq 0.66$, $0.66 \leq l^2 \leq 0.715$, and estimate separately. For $0.55 \leq l^2 \leq 0.58$, we find

$$\begin{aligned}
& -(75l^4 - 192l^2 + 140)K(l) + (35l^4 - 136l^2 + 140)E(l) \\
& \leq -(75 \cdot 0.58^2 - 192 \cdot 0.58 + 140)K(\sqrt{0.55}) \\
& \quad + (35 \cdot 0.55^2 - 136 \cdot 0.55 + 140)E(\sqrt{0.55}) \\
& \approx -1.87479 < 0.
\end{aligned}$$

For $0.58 \leq l^2 \leq 0.62$, we obtain

$$\begin{aligned}
& -(75l^4 - 192l^2 + 140)K(l) + (35l^4 - 136l^2 + 140)E(l) \\
& \leq -(75 \cdot 0.62^2 - 192 \cdot 0.62 + 140)K(\sqrt{0.58}) \\
& \quad + (35 \cdot 0.58^2 - 136 \cdot 0.58 + 140)E(\sqrt{0.58}) \\
& \approx -0.589051 < 0.
\end{aligned}$$

For $0.62 \leq l^2 \leq 0.66$,

$$\begin{aligned}
& -(75l^4 - 192l^2 + 140)K(l) + (35l^4 - 136l^2 + 140)E(l) \\
& \leq -(75 \cdot 0.66^2 - 192 \cdot 0.66 + 140)K(\sqrt{0.62}) \\
& \quad + (35 \cdot 0.62^2 - 136 \cdot 0.62 + 140)E(\sqrt{0.62}) \\
& \approx -1.5945 < 0.
\end{aligned}$$

For $0.66 \leq l^2 \leq 0.715$,

$$\begin{aligned}
& -(75l^4 - 192l^2 + 140)K(l) + (35l^4 - 136l^2 + 140)E(l) \\
& \leq -(75 \cdot 0.715^2 - 192 \cdot 0.715 + 140)K(\sqrt{0.66}) \\
& \quad + (35 \cdot 0.66^2 - 136 \cdot 0.66 + 140)E(\sqrt{0.66}) \\
& \approx -0.115848 < 0.
\end{aligned}$$

Hence, $2(7l^4 - 20l^2 + 20)(1 - l^2)K(l) - 5(2 - l^2)^3E(l)$ is monotonically decreasing. We finally estimate

$$\begin{aligned} & 2(7l^4 - 20l^2 + 20)(1 - l^2)K(l) - 5(2 - l^2)^3E(l) \\ & \leq 2(7 \cdot 0.55^2 - 20 \cdot 0.55 + 20)(1 - 0.55)K(\sqrt{0.55}) - 5(2 - 0.55)^3E(\sqrt{0.55}) \\ & \approx -1.19735 < 0, \end{aligned}$$

and the proof is completed.

Proof of (6)

Since $80/21 \approx 3.80952 > 3.8 = 19/5$, it is sufficient to prove $5A - 19C < 0$ for $5\pi/16 \leq \theta < \pi/2$. We first find

$$\begin{aligned} 5A - 19C &= \frac{1}{4\sqrt{2(1 + \sin \theta)} \sin^2 \theta (1 - \sin \theta)} \\ &\quad \times (40 \sin^2 \theta (1 - \sin \theta)K(k) - 19(E(k) - (1 - \sin \theta)K(k))), \\ &40 \sin^2 \theta (1 - \sin \theta)K(k) - 19(E(k) - (1 - \sin \theta)K(k)) \\ &= \frac{1}{(2 - k^2)^3} (2(59k^4 - 76k^2 + 76)(1 - k^2)K(k) - 19(2 - k^2)^3E(k)), \end{aligned}$$

and

$$\begin{aligned} &\frac{d}{dk} (2(59k^4 - 76k^2 + 76)(1 - k^2)K(k) - 19(2 - k^2)^3E(k)) \\ &= 2(236k^3 - 152k)(1 - k^2)K(k) + 2(59k^4 - 76k^2 + 76) \left(\frac{E - K}{k} - kK \right) \\ &\quad + 114k(2 - k^2)^2E(k) - 19(2 - k^2)^3 \cdot \frac{E - K}{k} \\ &= k(-7(87k^4 - 132k^2 + 76)K(k) + (133k^4 - 452k^2 + 532)E(k)). \end{aligned}$$

Note that

$$0.907981 \approx \frac{2 \sin \frac{5}{16}\pi}{1 + \sin \frac{5}{16}\pi} \leq k^2 < 1,$$

and, for $0.9 \leq x < 1$, $87x^2 - 132x + 76$ is positive and monotonically increasing, $133x^2 - 452x + 532$ is positive and monotonically decreasing. Hence, we obtain

$$\begin{aligned} &-7(87k^4 - 132k^2 + 76)K(k) + (133k^4 - 452k^2 + 532)E(k) \\ &\leq -7(87 \cdot 0.9^2 - 132 \cdot 0.9 + 76)K(\sqrt{0.9}) \end{aligned}$$

$$\begin{aligned}
& + (133 \cdot 0.9^2 - 452 \cdot 0.9 + 532)E(\sqrt{0.9}) \\
& \approx -242.015 < 0.
\end{aligned}$$

Thus, $2(59k^4 - 76k^2 + 76)(1 - k^2)K(k) - 19(2 - k^2)^3E(k)$ is monotonically decreasing. We finally estimate

$$\begin{aligned}
& 2(59k^4 - 76k^2 + 76)(1 - k^2)K(k) - 19(2 - k^2)^3E(k) \\
& \leq 2(59 \cdot 0.907^2 - 76 \cdot 0.907 + 76)(1 - 0.907)K(\sqrt{0.907}) \\
& \quad - 19(2 - 0.907)^3E(\sqrt{0.907}) \\
& \approx -0.247829 < 0,
\end{aligned}$$

and therefore the claim follows.

Proof of (7) Straightforward calculation yields

$$\begin{aligned}
B - 16D = & \frac{2}{\sqrt{2(1 + \cos \theta)} \cos^2 \theta (1 - \cos \theta)} \\
& \times (-2E(l) + (\cos^2 \theta + 2)(1 - \cos \theta)K(l)).
\end{aligned}$$

Note that

$$\frac{\pi}{4} \leq \theta < \frac{\pi}{2} \iff 0 < l^2 \leq 2(\sqrt{2} - 1),$$

and the inequality $B - 16D < 0$ is equivalent to

$$-E(l) + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \cdot (1 - l^2)K(l) < 0.$$

By **710.02**, **710.04** in [BF], we have

$$\begin{aligned}
& \frac{d}{dl} \left(-E(l) + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \cdot (1 - l^2)K(l) \right) \\
& = \frac{K(l) - E(l)}{l} + \frac{2l(3l^4 - 4l^2 + 8)}{(2 - l^2)^4} \cdot (1 - l^2)K(l) \\
& \quad + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \left(\frac{E(l) - K(l)}{l} - lK(l) \right) \\
& = \frac{l^2(l^4 - 3l^2 + 4)}{(2 - l^2)^3} \cdot \frac{E(l) - K(l)}{l} - \frac{3l^7}{(2 - l^2)^4}K(l) \\
& = \frac{l^2((l^2 - \frac{3}{2})^2 + \frac{7}{4})}{(2 - l^2)^3} \cdot \frac{E(l) - K(l)}{l} - \frac{3l^7}{(2 - l^2)^4}K(l) < 0.
\end{aligned}$$

Hence,

$$-E(l) + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \cdot (1 - l^2)K(l)$$

is monotonically decreasing. We finally estimate

$$\begin{aligned} & -E(l) + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \cdot (1 - l^2)K(l) \\ & < \lim_{l \rightarrow 0} \left(-E(l) + \frac{3l^4 - 8l^2 + 8}{(2 - l^2)^3} \cdot (1 - l^2)K(l) \right) = -E(0) + K(0) \\ & = -\frac{\pi}{2} + \frac{\pi}{2} = 0. \end{aligned}$$

Therefore it completes the proof.

Proof of (8) Note that

$$\frac{3}{8}\pi \leq \theta < \frac{\pi}{2} \iff (2 + \sqrt{2}) \left(2\sqrt{2 + \sqrt{2}} - 2 - \sqrt{2} \right) \leq k^2 < 1.$$

We next obtain

$$15A - 32C = \frac{2((15\sin^2\theta + 4)(1 - \sin\theta)K(k(\theta)) - 4E(k(\theta)))}{\sqrt{2(1 + \sin\theta)} \sin^2\theta(1 - \sin\theta)}.$$

Thus it is sufficient to prove

$$\begin{aligned} & (15\sin^2\theta + 4)(1 - \sin\theta)K(k(\theta)) - 4E(k(\theta)) \\ & = \frac{2}{(2 - k^2)^3} ((19k^4 - 16k^2 + 16)(1 - k^2)K(k) - 2(2 - k^2)^3 E(k)) < 0. \end{aligned}$$

To do this, direct calculations yield

$$\begin{aligned} \frac{d}{dk}(19k^4 - 16k^2 + 16) &= 2k(38k^2 - 16), \\ \frac{d}{dk}(1 - k^2)K(k) &= \frac{E(k)}{k} - \frac{1 + k^2}{k}K(k), \\ \frac{d}{dk}(2 - k^2)^3 &= -6k(2 - k^2)^2, \quad \frac{d}{dk}E(k) = \frac{E(k) - K(k)}{k}. \end{aligned}$$

Hence, we have

$$\begin{aligned} \frac{d}{dk} & ((19k^4 - 16k^2 + 16)(1 - k^2)K(k) - 2(2 - k^2)^3E(k)) \\ &= k(-97k^4 + 117k^2 - 56)K(k) + k(14k^4 - 41k^2 + 56)E(k) \\ &= k((14k^4 - 41k^2 + 56)\underbrace{(-K(k) + E(k))}_{<0} + k^2(76 - 83k^2)K(k)). \end{aligned}$$

We now observe that

$$(2 + \sqrt{2}) \left(2\sqrt{2 + \sqrt{2}} - 2 - \sqrt{2} \right) \approx 0.9604 \dots$$

Set $x = k^2$, and, for $0.96 < x < 1$, we find

$$14x^2 - 41x + 56 > 0, \quad 76 - 83x < 0.$$

Thus

$$\frac{d}{dk} ((19k^4 - 16k^2 + 16)(1 - k^2)K(k) - 2(2 - k^2)^3E(k)) < 0.$$

It follows that $(19k^4 - 16k^2 + 16)(1 - k^2)K(k) - 2(2 - k^2)^3E(k)$ is monotonically decreasing. We finally estimate

$$\begin{aligned} & (19k^4 - 16k^2 + 16)(1 - k^2)K(k) - 2(2 - k^2)^3E(k) \\ &= 0.726016 \cdot K(\sqrt{0.96}) - 2.24973 \cdot E(\sqrt{0.96}) \\ &= 0.726016 \cdot 3.016112 - 2.24973 \cdot 1.050502 = -0.173598 < 0. \end{aligned}$$

Therefore, for $3\pi/8 \leq \theta < \pi/2$, $15A - 32C < 0$ holds.

References

- [BF] P. F. Byrd and M. D. Friedman, Handbook of elliptic integrals for engineers and scientists. Second edition, revised, Die Grundlehren der mathematischen Wissenschaften Band 67, Springer-Verlag, New York-Heidelberg, 1971.
- [EK] N. Ejiri and M. Kotani, Index and flat ends of minimal surfaces, Tokyo J. Math. **16** (1993), no. 1, 37–48.

- [Her] J. Hersch, Quatre propriétés isopérimétriques de membranes sphériques homogènes, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A1645–A1648.
- [JLNNP] D. Jakobson, M. Levitin, N. Nadirashvili, N. Nigam, I. Polterovich, How large can the first eigenvalue be on a surface of genus two?, Int. Math. Res. Not. (2005), no. 63, 3967–3985.
- [MR] S. Montiel and A. Ros, Schrödinger operators associated to a holomorphic map, in: Global differential geometry and global analysis (Berlin, 1990), Lecture Notes in Math. **1481**, Springer, Berlin, 1991, pp. 147–174.
- [N] N. Nadirashvili, Berger’s isoperimetric problem and minimal immersions of surfaces, Geom. Funct. Anal. **6** (1996), no. 5, 877–897.
- [NS] S. Nayatani and T. Shoda, Metrics on a closed surface of genus two which maximize the first eigenvalue of the Laplacian, C. R. Math. Acad. Sci. Paris **357** (2019), no. 1, 84–98.
- [YY] P. C. Yang and S. T. Yau, Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), no. 1, 55–63.

Toshihiro Shoda,
 Faculty of Education,
 Saga University,
 Honjo-machi 1-chome, Saga-city, Saga, 840-8502, Japan.
 tshoda@cc.saga-u.ac.jp