# Integral algebra

# 近畿大学・工学部 田中 広志 Hiroshi Tanaka Faculty of Engineering, Kindai University

#### Abstract

A differential ring is a ring equipped with a derivation. In this paper, we introduce integral rings. An integral ring is a ring equipped with a derivation and an integration. We study the basic properties of integral rings.

### 1 Introduction

Throughout this paper, all rings will mean commutative rings of characteristic zero with identity. In this paper we will introduce integral rings. A differential ring is a ring equipped with a derivation. An integral ring is a ring equipped with a derivation and an integration. Differential rings and differential fields were studied in [1], [2] and [5]. However an integral ring is not studied. So we will study the basic properties of an integral ring.

## 2 Differential rings

In this section we introduce differential rings. We follow the notation and definition of differential rings from [4], [3] and [6].

<sup>2020</sup> Mathematics Subject Classification. 03C60.

Key Words and Phrases. Integral fields and differential fields.

**Definition 1.** A derivation on a ring R is an additive homomorphism  $\delta$ :  $R \to R$  such that  $\delta(xy) = x\delta(y) + y\delta(x)$ . A differential ring is a ring equipped with a derivation.

By the definition of  $\delta$ , for all  $x \in R$  and  $n \in \mathbb{N}$ , we have  $\delta(x^n) = nx^{n-1}\delta(x)$ .

**Definition 2.** We let  $C_R$  denote the kernel of the derivation  $\delta$ . We call  $C_R$  the *constants* of R.

 $C_R$  is a subring of R. In particular, if R is a field, then so is  $C_R$ .

**Definition 3.** Given a differential ring R, the differential polynomial ring  $R\{X\}$  in one differential indeterminate X is the ring  $R\{X\}$  :=  $R[X, \delta(X), \delta^2(X), \ldots]$ . By extending  $\delta$  to all of  $R\{X\}$  in the obvious manner, we have that  $R\{X\}$  is a differential ring. For nonzero  $f \in R\{X\}$  we define  $\operatorname{ord}(f)$ , the order of f, to be the largest n, if any, such that  $\delta^n(X)$  appears in f with non-zero coefficient.

**Definition 4.** We say that an ideal  $I \subseteq R$  is a differential ideal if for all  $f \in I$ , we have  $\delta(f) \in I$ .

**Definition 5.** We say that a field K is a differentially closed field if K is a differential field such that if  $f, g \in K\{X\} \setminus \{0\}$  and  $\operatorname{ord}(f) > \operatorname{ord}(g)$ , then there exists  $x \in K$  such that f(x) = 0 and  $g(x) \neq 0$ .

Any differentially closed field is algebraically closed.

Fact 6 ([4, Lemma 2.2]). Every differential field k has an extension K which is differentially closed.

# 3 Integral rings

**Definition 7.** A integration on a differential ring R is a function  $\rho: R \to R$  such that the followings hold.

1. For any  $x \in R$ , we have  $\delta(\rho(x)) = x$ .

2. For any  $x \in R$ , there exists  $\alpha \in R$  such that  $\rho(\delta(x)) = x + \alpha$  and  $\delta(\alpha) = 0$ .

A integral ring is a differential ring equipped with an integration.

By the definition of an integration  $\rho$ , it follows that  $\delta$  is surjective and  $\rho$  is injective. Since  $\delta(x^n) = nx^{n-1}\delta(x)$  for any  $x \in R$  and  $n \in \mathbb{N}$ , there exists  $\alpha \in C_R$  such that  $\rho(x^n\delta(x)) = \frac{1}{n+1}x^{n+1} + \alpha$ .

In this paper we let integration be a function, but we may let integration be a relation.

**Remark 8.** There exists a differential ring which is not an integral ring. For example, let  $\mathbb{R}\left\{\frac{1}{X}\right\}$  be a differential ring, where  $\delta\left(\frac{1}{X}\right) = -\frac{1}{X^2}$ . Then,  $\mathbb{R}\left\{\frac{1}{X}\right\}$  is not an integral ring. By Fact 6, there exists a differentially closed field  $K \supseteq \mathbb{R}\left\{\frac{1}{X}\right\}$ . Then, there exists  $x \in K$  such that  $\delta(x) = \frac{1}{X}$ .

**Lemma 9.** Let  $x, y \in R$ . The followings hold.

- 1. There exists  $\alpha \in C_R$  such that  $\rho(x+y) = \rho(x) + \rho(y) + \alpha$ .
- 2. For any  $\alpha \in C_R$ , there exists  $\beta \in C_R$  such that  $\rho(\alpha x) = \alpha \rho(x) + \beta$ .
- 3. There exists  $\alpha \in C_R$  such that  $\rho(\delta(x)y) = xy \rho(x\delta(y)) + \alpha$ .

**Definition 10.** Suppose that R is an integral ring. The strong constant ring of R is the ring  $C'_R := \{x \in R : \exists n \in \mathbb{N}, \delta^n(x) = 0\}$ 

The constant ring  $C_R$  of R is not an integral ring.

**Lemma 11.** The strong constant ring  $C'_R$  of R is an integral ring.

**Definition 12.** We say that a differential ideal  $I \subseteq R$  is a *integral ideal* if for all  $f \in I$ , we have  $\rho(f) \in I$ .

If  $I \subseteq R$  is a differential ideal, then naturally R/I is a differential quotient ring of R. However, if  $I \subseteq R$  be a integral ideal, then R/I is not an integral quotient ring of R.

**Lemma 13.** Let  $I \subsetneq R$  be an integral ideal. Then there exists a maximal integral ideal  $J \supseteq I$ .

## References

- [1] E. R. Kolchin, Galois Theory of Differential Fields, Amer. J. Math 75, 1953, 753-824.
- [2] E. R. Kolchin, Constrained Extensions of Differential Fields, Advances in Math. 12, 1974, 141-170.
- [3] D. Marker, Model theory: an introduction, Graduate Texts in Mathematics 217, Springer, 2002.
- [4] D. Marker, *Model Theory of Differential Fields*, in Model Theory of Fields Second Edition, Lecture Notes in Logic **5**, Springer, 2005.
- [5] A. Seidenberg, Some Basic Theorems in Differential Algebra, Trans. Amer. Math. Soc. 73, 1952, 174-190.
- [6] C. Wood, *Differentially Closed Fields*, in Model Theory and Algebraic Geometry, Lecture Notes in Math. **1696**, Springer 1998.