

Convex minimization problems on geodesic spaces
and improvement of approximating sequences
測地距離空間における凸最小化問題と近似列の改良

東邦大学・理学研究科 日高雄太
Yuta Hidaka

Department of Information Science
Toho University

東邦大学・理学部 木村泰紀
Yasunori Kimura

Department of Information Science
Toho University

Abstract

In 2021, Hidaka and Kimura proved a Δ -convergence theorem in CAT(1) spaces. In this paper, we modified the coefficient condition and obtain another Δ -convergence theorem.

1 Introduction

In [1], the authors proved the following theorem.

Theorem 1 ([1]). *Let X be an admissible complete CAT(1) space. Let $f : X \rightarrow]-\infty, \infty]$ be a proper convex lower semicontinuous function and suppose that $\operatorname{argmin} f \neq \emptyset$. Let $\{\beta_n\}$ and $\{\gamma_n\}$ be real sequences in $[0, 1]$ such that $\sum_{n=1}^{\infty} \beta_n = \infty$ and that both $\{\beta_n\}$ and $\{\gamma_n\}$ converge to 0. For an initial point $x_1 \in X$, generate a sequence $\{x_n\}$ as follows:*

$$\begin{aligned} y_n &= J_f x_n, \\ \alpha_n &\in [\min\{\beta_n, d(x_n, y_n) - \gamma_n\}, 1] \cap [0, 1], \\ x_{n+1} &= (1 - \alpha_n)x_n \oplus \alpha_n y_n. \end{aligned}$$

Suppose that one of the following conditions holds:

- $\inf_{n \in \mathbb{N}} \alpha_n > 0$;
- $\sum_{n=1}^{\infty} \alpha_n < \infty$.

Then, $x_n \xrightarrow{\Delta} x_0 \in \operatorname{argmin} f$.

In this paper, we try to remove the condition that $\inf_{n \in \mathbb{N}} \alpha_n > 0$ in Theorem 1. In order to this, we modify the coefficient condition.

2 Preliminaries

Let λ be a positive real number. A metric space X is said to be λ -geodesic if for each $x, y \in X$ with $d(x, y) < \lambda$, there exists a mapping $c : [0, l] \rightarrow X$ such that $c(0) = x, c(l) = y$, and

$$d(c(t_1), c(t_2)) = |t_1 - t_2|$$

for all $t_1, t_2 \in [0, l]$, where $l = d(x, y)$. The mapping c is called a geodesic from x to y . If a geodesic c from x to y is unique, the geodesic segment $[x, y]$ is defined by

$$[x, y] = \{c(t) : 0 \leq t \leq l\}.$$

Let X be a uniquely geodesic space, and $x, y, z \in X$. Let $x, y, z \in X$ with $d(x, y) + d(y, z) + d(z, x) < 2\pi$. The set $\Delta = \Delta(x, y, z)$ is defined by $\Delta = [x, y] \cup [y, z] \cup [z, x]$. We take $\bar{x}, \bar{y}, \bar{z} \in \mathbb{S}^2$ such that $d(x, y) = d_{\mathbb{S}^2}(\bar{x}, \bar{y}), d(y, z) = d_{\mathbb{S}^2}(\bar{y}, \bar{z}), d(z, x) = d_{\mathbb{S}^2}(\bar{z}, \bar{x})$. The set $\bar{\Delta} = \bar{\Delta}(\bar{x}, \bar{y}, \bar{z})$ is defined by $\bar{\Delta} = [\bar{x}, \bar{y}] \cup [\bar{y}, \bar{z}] \cup [\bar{z}, \bar{x}]$. X is called a CAT(1) space, if for all Δ and $p, q \in \Delta$ with $\bar{p}, \bar{q} \in \bar{\Delta}$,

$$d(p, q) \leq d_{\mathbb{S}^2}(\bar{p}, \bar{q}).$$

We say that a CAT(1) space X is admissible if

$$d(w, w') < \frac{\pi}{2}$$

for all $w, w' \in X$.

The definition of the resolvent of f is as follows [4]:

$$J_f x = \operatorname{argmin}_{y \in X} \{f(y) + \tan d(y, x) \sin d(y, x)\},$$

where $f : X \rightarrow]-\infty, \infty]$ is a proper lower semicontinuous function. We denote by $\operatorname{argmin}_X f$ the set of all $u \in X$ such that $f(u) = \inf f(X)$. For a bounded sequence $\{x_n\} \subset X$, the asymptotic center $\mathcal{A}(\{x_n\})$ of $\{x_n\}$ is defined by

$$\mathcal{A}(\{x_n\}) = \left\{ u \in X \mid \limsup_{n \rightarrow \infty} d(u, x_n) = \inf_{y \in X} \limsup_{n \rightarrow \infty} d(y, x_n) \right\}.$$

A sequence $\{x_n\}$ is said to be Δ -convergent to a point $p \in X$ if

$$\mathcal{A}(\{x_{n_i}\}) = \{p\}$$

for each subsequence $\{x_{n_i}\}$ of $\{x_n\}$.

We know the following lemmas.

Lemma 1 ([4]). *Let X be an admissible complete CAT(1) space, f a proper convex lower semicontinuous functions of X into $]-\infty, \infty]$. Let $J_{\eta f}$ be the resolvent of ηf for all $\eta > 0$ and $C_{\eta, z}$ the real number given by*

$$C_{\eta, z} = \cos d(J_{\eta f} z, z)$$

for all $\eta > 0$ and $z \in X$. Then

$$\begin{aligned} & (\lambda C_{\lambda, x}^2 (1 + C_{\mu, y}^2) C_{\mu, y} + \mu C_{\mu, y}^2 (1 + C_{\lambda, x}^2) C_{\lambda, x}) \cos d(J_{\lambda f} x, J_{\mu f} y) \\ & \geq \lambda C_{\lambda, x}^2 (1 + C_{\mu, y}^2) \cos d(J_{\lambda f} x, y) + \mu C_{\mu, y}^2 (1 + C_{\lambda, x}^2) \cos d(J_{\mu f} y, x) \end{aligned}$$

for all $x, y \in X$ and $\lambda, \mu > 0$. Further,

$$\frac{\pi}{2} \left(\frac{1}{C_{\lambda, x}^2} + 1 \right) (C_{\lambda, x} \cos d(u, J_{\lambda f} x) - \cos d(u, x)) \geq \lambda (f(J_{\lambda f} x) - f(u))$$

and

$$(1) \quad \cos d(J_{\lambda f} x, x) \cos d(u, J_{\lambda f} x) \geq \cos d(u, x)$$

for all $x \in X, u \in \operatorname{argmin}_X f$ and $\lambda > 0$.

Let X be a metric space such that $d(v, v') < \pi/2$ for all $v, v' \in X$, T a mapping of X into itself, and C_z the real number given by

$$C_z = \cos d(Tz, z)$$

for all $z \in X$. The mapping T is said to be vicinal [5] if

$$\begin{aligned} & (C_x^2 (1 + C_y^2) + C_y^2 (1 + C_x^2)) \cos d(Tx, Ty) \\ & \geq C_x^2 (1 + C_y^2) \cos d(Tx, y) + C_y^2 (1 + C_x^2) \cos d(Ty, x) \end{aligned}$$

for all $x, y \in X$.

Lemma 2 ([5]). *Let X be a metric space such that $d(v, v') < \pi/2$ for all $v, v' \in X$, T a vicinal mapping of X into itself, p an element of X , and $\{x_n\}$ a sequence in X such that $\mathcal{A}(\{x_n\}) = \{p\}$ and $d(Tx_n, x_n) \rightarrow 0$. Then p is a fixed point of T .*

Lemma 3 ([3]). *Let X be a complete CAT(1) space such that $d(v, v') < \pi/2$ for all $v, v' \in X$. Let f be a proper lower semicontinuous convex function of X into $]-\infty, \infty]$, J_f the resolvent of f . Then $F(J_f) = \operatorname{argmin}_X f$.*

3 Main result

The following theorem is the main result of this paper.

Theorem 2. *Let X be an admissible complete CAT(1) space. Let $f : X \rightarrow]-\infty, \infty]$ be a proper convex lower semicontinuous function such that $\operatorname{argmin} f \neq \emptyset$. Let $\{\gamma_n\}$ be a real sequence in $[0, 1]$ converging to 0. For an initial point $x_1 \in X$ such that $f(x_1) < \infty$, generate a sequence $\{x_n\}$ as follows:*

$$\begin{aligned} y_n &= J_f x_n, \\ \alpha_n &\in \left[\frac{1}{2}d(x_n, y_n) - \gamma_n, 1 \right] \cap [0, 1], \\ x_{n+1} &= (1 - \alpha_n)x_n \oplus \alpha_n y_n. \end{aligned}$$

Then, $x_n \xrightarrow{\Delta} x_0 \in \operatorname{argmin} f$.

Proof. Take $\{d(x_{n_i}, y_{n_i})\} \subset \{d(x_n, y_n)\}$. There exists $\{\alpha_{n_{i_j}}\} \subset \{\alpha_{n_i}\}$ such that $\alpha_{n_{i_j}} \rightarrow \alpha_0 \in [0, 1]$. If $\alpha_0 = 0$, then

$$\frac{1}{2}d(x_{n_{i_j}}, y_{n_{i_j}}) \leq \alpha_{n_{i_j}} + \gamma_{n_{i_j}} \rightarrow 0.$$

We get $d(x_{n_{i_j}}, y_{n_{i_j}}) \rightarrow 0$.

If $\alpha_0 \in]0, 1]$, for $u \in \operatorname{argmin} f$, using (1) in Lemma 1, we have

$$\begin{aligned} \cos d(u, x_{n+1}) &\geq (1 - \alpha_n) \cos d(u, x_n) + \alpha_n \cos d(u, y_n) \\ &\geq (1 - \alpha_n) \cos d(u, x_n) + \alpha_n \frac{\cos d(u, x_n)}{\cos d(y_n, x_n)} \\ &= \cos d(u, x_n) + \alpha_n \cos d(u, x_n) \left(\frac{1}{\cos d(y_n, x_n)} - 1 \right). \end{aligned}$$

It implies

$$\frac{\cos d(u, x_{n+1})}{\cos d(u, x_n)} - 1 \geq \alpha_n \left(\frac{1}{\cos d(y_n, x_n)} - 1 \right).$$

We know that $d(u, x_n) \rightarrow [0, \pi/2[$. In fact, since

$$\cos d(u, x_{n+1}) \geq (1 - \alpha_n) \cos d(u, x_n) + \alpha_n \cos d(u, y_n) \geq \cos d(u, x_n),$$

we have

$$d(u, x_{n+1}) \leq d(u, x_n).$$

Thus $d(u, x_n) \rightarrow [0, \pi/2[$. Hence we get,

$$0 \leq \alpha_{n_{i_j}} \left(\frac{1}{\cos d(y_{n_{i_j}}, x_{n_{i_j}})} - 1 \right) \leq \frac{\cos d(u, x_{n_{i_j}+1})}{\cos d(u, x_{n_{i_j}})} - 1 \rightarrow 0.$$

We get $d(x_{n_{i_j}}, y_{n_{i_j}}) \rightarrow 0$. It means that $d(x_n, y_n) \rightarrow 0$.

Let $\{x_{n_i}\} \subset \{x_n\}$ with $w = \mathcal{A}(\{x_{n_i}\})$. There exists $\{x_{n_{i_j}}\} \subset \{x_{n_i}\}$ such that $x_{n_{i_j}} \xrightarrow{\Delta} z$. In fact, since

$$d(u, x_{n+1}) \leq d(u, x_n) \leq d(u, x_1) < \frac{\pi}{2},$$

we have $\limsup_{n \rightarrow \infty} d(u, x_n) < \pi/2$. Using Lemma 2 and Lemma 3, we have $z \in F(J_f) = \operatorname{argmin} f$. We put $v = \mathcal{A}(\{x_n\})$. Then,

$$\begin{aligned} \lim_{n \rightarrow \infty} d(x_n, z) &= \lim_{i \rightarrow \infty} d(x_{n_i}, z) \\ &= \lim_{j \rightarrow \infty} d(x_{n_{i_j}}, z) \\ &\leq \limsup_{j \rightarrow \infty} d(x_{n_{i_j}}, w) \\ &\leq \limsup_{i \rightarrow \infty} d(x_{n_i}, w) \\ &\leq \limsup_{i \rightarrow \infty} d(x_{n_i}, v) \\ &\leq \limsup_{n \rightarrow \infty} d(x_n, v). \end{aligned}$$

We get $v = w = z$. Hence, $x_n \xrightarrow{\Delta} v = z \in \operatorname{argmin} f$. □

Acknowledgement. This work was partially supported by JSPS KAKENHI Grant Number JP21K03316.

References

- [1] Y. Hidaka and Y. Kimura, *An improvement of the coefficient condition for a convergence theorem in a complete geodesic space*, RIMS kôkyûroku, no. 2194 (2021), pp. 13–17.
- [2] Y. Kimura, *Further improvement of a coefficient condition for a weakly convergent iterative scheme*, Nonlinear Anal. **71** (2009), e2023–e2027.
- [3] Y. Kimura and F. Kohsaka, *Spherical nonspreadingness of resolvents of convex functions in geodesic spaces*, J. Fixed Point Theory Appl. **18** (2016), 93–115.
- [4] Y. Kimura and F. Kohsaka, *Two modified proximal point algorithms in geodesic spaces with curvature bounded above*, Rend. Circ. Mat. Palermo, II. Ser **68** (2019), 83–104.
- [5] F. Kohsaka, *Existence and approximation of fixed points of vicinal mappings in geodesic spaces*, Pure Appl. Funct. Anal (2018), 91–106.