

NOTE ON THE SOLVABILITY OF THE FULLY FOURTH ORDER NONLINEAR BOUNDARY VALUE PROBLEMS

TOKYO UNIVERSITY OF INFORMATION SCIENCES
TOSHIKAZU WATANABE

1. INTRODUCTION

Many researchers have considered the following differential equation;

$$(1.1) \quad \begin{cases} u''''(t) = f(t, u(t)), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = u'''(1) = 0, \end{cases}$$

where f is a continuous mapping from $[0, 1] \times \mathbb{R}$ into \mathbb{R} , we denote by \mathbb{R} the set of all real numbers; see [9, 10, 14, 15, 21, 22, 23, 25, 26]. Equation (1.1) can be used to model the deformations of an Cantilever equation. In mechanics, the problem is called Cantilever beam equation, and in the equation, the physical meaning of the derivatives of the deformation function $u(t)$ is as follows: u'''' is the load density stiffness, u''' is the shear force stiffness, u'' is the bending moment stiffness, and u' is the slope ;see [9, 10]. In some practice, only its positive solution is significant.

Meanwhile, fractional differential equations have been studied by many researchers; see [3, 5, 6, 7, 8, 11, 12, 17, 18, 19, 24]. However, to the best of our knowledge, there are no results for the boundary value problem represented by (1.4) for $3 < \alpha \leq 4$, which we consider in the present paper. We use the method of order reduction and contraction principle [27, 28] to prove the existence and uniqueness of solutions. In [20], we consider the boundary value problem for fractional order differential equation

$$(1.2) \quad \begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & t \in [0, 1] \\ u(0) = u'(0) = u''(1) = u'''(1) = 0, \end{cases}$$

where D_{0+}^{α} is the Riemann-Liouville derivative of order α , $3 < \alpha \leq 4$, and f is a continuous function of $[0, 1] \times \mathbb{R}$ into \mathbb{R} .

Several authors consider the following fourth order boundary value problems for the fully nonlinear with boundary conditions see; [27, 28, 29, 30, 31, 32, 33].

$$(1.3) \quad \begin{cases} u''''(t) = f(t, u(t), u'(t), u''(t), u'''(t)), & 0 < t < 1, \\ u(0) = u'(0) = u''(1) = u'''(1) = 0, \end{cases}$$

Recently in [27], using an order reduction and contraction method, authors give an existence and uniqueness of solution. In [28] authors using a contraction method and iteration method they give an existence and uniqueness of solutions of boundary value problems and also examples of non-linear function as a function f in (1.3).

2010 *Mathematics Subject Classification.* Primary 46A40, 47B50, 47H10.

Key words and phrases. Fixed point theorem, fractional boundary value problem.

In this paper we propose the following differential equation of fractional order α , $3 < \alpha \leq 4$ with the two point boundary condition involving the form (1.2).

$$(1.4) \quad \begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t), D_{0+}^{\alpha-3} u(t), D_{0+} D_{0+}^{\alpha-3} u(t), D_{0+} D_{0+} D_{0+}^{\alpha-3} u(t)), \\ 0 < t < 1, \\ u(0) = D_{0+}^{\alpha-3} u(0) = D_{0+}^{\alpha-2} u(1) = D_{0+}^{\alpha-1} u(1) = 0. \end{cases}$$

In particular, using a order reduction and contraction method, we give a unique solution comparing method in [27, 28], we also give a suitable Lipschitz constant through a study.

Let $\alpha > 0$. The Riemann-Liouville fractional integral of order α of u , denoted $I_{0+}^{\alpha} u$, is defined by

$$I_{0+}^{\alpha} u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} u(s) ds,$$

provided the right-hand side exists. The Riemann-Liouville fractional derivative of order α of a function u of $(0, \infty)$ into \mathbb{R} is given by

$$D_{0+}^{\alpha} u(t) = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t (t-s)^{n-\alpha-1} u(s) ds,$$

where $n = [\alpha] + 1$ ($[\alpha]$ denotes the integer part of α) and $\Gamma(\alpha)$ denotes the gamma function; see [11, 18]. Note that for $\alpha > \beta > 0$, we have

$$D_{0+}^{\alpha} t^{\beta} = \frac{\Gamma(\beta+1)}{\Gamma(\beta-\alpha+1)} t^{\beta-\alpha}.$$

A function $u \in C[0, 1]$ is called a solution of problem (1.4) if $D^{\alpha} u \in C[0, 1]$, $D^{\alpha-3} u \in L^1[0, 1]$, $D^{\alpha-2} u \in L^1[0, 1]$, $D^{\alpha-1} u \in L^1[0, 1]$, u satisfies the boundary conditions and equality in (1.4) a.e. on $[0, 1]$.

2. LEMMAS

Let h be a continuous mapping $[0, 1]$ into \mathbb{R} , we consider the following fractional differential boundary value problems defined by

$$(2.1) \quad \begin{cases} D_{0+}^{\alpha} u(t) = h(t), & 0 < t < 1, \\ u(0) = u(1) = 0, D_{0+} D_{0+}^{\alpha-3} u(0) = D_{0+} D_{0+}^{\alpha-3} u(1) = 0, \end{cases}$$

where $3 < \alpha \leq 4$. In this section, we show the unique solution to the boundary value problem represented by (2.1). A mapping u of $[0, 1]$ into \mathbb{R} is a solution of that boundary value problems if u is continuous on $[0, 1]$ and u satisfies (2.1).

The following lemma can be found in [6]; see also [11]. We denoted by $C(0, 1)$ the set of all continuous mappings of $(0, 1)$ into \mathbb{R} and by $L(0, 1)$ the set of all Lebesgue integrable mappings of $[0, 1]$ into \mathbb{R} .

Lemma 1. *Let $\alpha > 0$. If $u(t) \in C(0, 1) \cap L(0, 1)$ satisfying $D_{0+}^{\alpha} u(t) \in C(0, 1) \cap L(0, 1)$, then there exist constants $C_1, C_2, \dots, C_n \in \mathbb{R}$ such that*

$$I_{0+}^{\alpha} D_{0+}^{\alpha} u(t) = u(t) + C_1 t^{\alpha-1} + C_2 t^{\alpha-2} + \dots + C_n t^{\alpha-n},$$

where $n = [\alpha] + 1$ and $I_{0+}^{\alpha} u$ is the Riemann-Liouville fractional integral of order α of a function u is defined by

$$I_{0+}^{\alpha} u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} u(s) ds.$$

Using Lemma 1, we obtain the following.

Lemma 2. *Let $g \in C^n[0, 1]$ be given. Then the unique solution to problem $D^\alpha y(t) = g(t)$ together with the boundary conditions in (1.4) is*

$$u(t) = \int_0^t G(t, s)g(s)ds,$$

where

$$(2.2) \quad G(t, s) = \begin{cases} \frac{1}{\Gamma(\alpha)} ((t-s)^{\alpha-1} - t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) & \text{if } 0 \leq s \leq t \leq 1, \\ \frac{1}{\Gamma(\alpha)} (-t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) & \text{if } 0 \leq t \leq s < 1. \end{cases}$$

Proof. In order to have $G(t, s)$, by Lemma 1, we have

$$\begin{aligned} u(t) &= c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + c_3 t^{\alpha-3} + c_4 t^{\alpha-4} + I_+^\alpha g(t) \\ &= c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + c_3 t^{\alpha-3} + c_4 t^{\alpha-4} + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds \end{aligned}$$

By the boundary condition in (1.4), since $u(0) = 0$, we have $c_4 = 0$.

$$D_{0+}^{\alpha-3} \left(\int_0^t (t-s)^{\alpha-1} g(s) ds \right) = \frac{\Gamma(\alpha)}{2} \int_0^t (t-s)^2 g(s) ds.$$

Since $D_{0+}^{\alpha-3} u(0) = 0$, we have $c_3 = 0$. Since

$$D_{0+}^{\alpha-2} \left(\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds \right) = \int_0^t (t-s) g(s) ds.$$

and $D_{0+}^{\alpha-2} u(1) = 0$, we have

$$0 = c_1 \Gamma(\alpha) + c_2 \Gamma(\alpha-1) + \int_0^1 (1-s) g(s) ds.$$

Also we have

$$D_{0+}^{\alpha-1} \left(\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds \right) = \int_0^t g(s) ds.$$

Then we have

$$c_1 = -\frac{1}{\Gamma(\alpha)} \int_0^1 g(s) ds.$$

Since

$$\begin{aligned} 0 &= c_1 \Gamma(\alpha) + c_2 \Gamma(\alpha-1) + \int_0^1 (1-s) g(s) ds \\ &= -\int_0^1 g(s) ds + c_2 \Gamma(\alpha-1) + \int_0^1 (1-s) g(s) ds, \end{aligned}$$

we have

$$\begin{aligned} c_2 &= -\frac{1}{\Gamma(\alpha-1)} \left(\int_0^1 (1-s) g(s) ds - \int_0^1 g(s) ds \right) \\ &= \frac{(\alpha-1)}{\Gamma(\alpha)} \left(\int_0^1 s g(s) ds \right). \end{aligned}$$

Hence we have

$$\begin{aligned}
u(t) &= c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + D_{0+}^{-\alpha} g(t) \\
&= \frac{1}{\Gamma(\alpha)} \int_0^1 -g(s) ds t^{\alpha-1} + \frac{(\alpha-1)}{\Gamma(\alpha)} \left(\int_0^1 s g(s) ds \right) t^{\alpha-2} \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} g(s) ds \\
&= \frac{1}{\Gamma(\alpha)} \left(\int_0^t ((t-s)^{\alpha-1} - t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) g(s) ds \right. \\
&\quad \left. + \int_t^1 (-t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) g(s) ds \right).
\end{aligned}$$

Then the green function $G(t, s)$ is defined by

$$G(t, s) = \begin{cases} \frac{1}{\Gamma(\alpha)} ((t-s)^{\alpha-1} - t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) & \text{if } 0 \leq s \leq t, \\ \frac{1}{\Gamma(\alpha)} (-t^{\alpha-1} + (\alpha-1)st^{\alpha-2}) & \text{if } t \leq s < 1. \end{cases}$$

□

Next in order to estimate the order reduction, we consider the followings.

$$\int_0^t G(t, s) h(s) ds = [-G(t, s)v(s)]_0^t + \int_0^t G_1(t, s)v(s) ds,$$

and

$$\int_t^1 G(t, s) h(s) ds = [-G(t, s)v(s)]_t^1 + \int_t^1 G_1(t, s)v(s) ds,$$

where $v(t) = \int_t^1 h(s) ds$ and $G_1(t, s) = \frac{\partial G}{\partial s}(t, s)$;

$$(2.3) \quad G_1(t, s) = \begin{cases} \frac{1}{\Gamma(\alpha-1)} (-(t-s)^{\alpha-2} + t^{\alpha-2}) & \text{if } 0 \leq s \leq t \leq 1 \\ \frac{1}{\Gamma(\alpha-1)} t^{\alpha-2} & \text{if } 0 \leq t \leq s \leq 1. \end{cases}$$

Moreover

$$\int_0^1 G(t, s) h(s) ds = \int_0^1 G_1(t, s)v(s) ds.$$

In fact since $v(t) = \int_1^t h(s) ds$, $v(1) = 0$,

$$\begin{aligned}
&[-G(t, s)v(s)]_0^1 \\
&= -\frac{1}{\Gamma(\alpha)} \left([(t-s)^{\alpha-1}v(s)]_0^t + [-t^{\alpha-1} + (\alpha-1)st^{\alpha-2}v(s)]_0^1 \right) \\
&= \frac{1}{\Gamma(\alpha)} (t^{\alpha-1}) v(0) \\
&\quad + \frac{1}{\Gamma(\alpha)} ((-t^{\alpha-1} + (\alpha-1)t^{\alpha-2}) v(1) - t^{\alpha-1}v(0)) = 0.
\end{aligned}$$

We also have

$$D_{0+}^{\alpha-3} \left(\int_0^1 G_1(t, s)v(s) ds \right) = \int_0^1 G_2(t, s)v(s) ds,$$

where

$$(2.4) \quad G_2(t, s) = \begin{cases} s & (0 \leq s \leq t \leq 1), \\ t & (0 \leq t \leq s \leq 1). \end{cases}$$

$$D_{0+}D_{0+}^{\alpha-3} \left(\int_0^1 G(t, s)h(s)ds \right) = D_{0+} \left(\int_0^1 G_2(t, s)v(s)ds \right) = \int_0^1 G_3(t, s)v(s)ds.$$

where

$$(2.5) \quad G_3(t, s) = \begin{cases} 0 & (0 \leq s \leq t \leq 1), \\ 1 & (0 \leq t \leq s \leq 1). \end{cases}$$

3. ESTIMATES OF INTEGRAL EQUATIONS

In this section in order to have the solution of boundary value problem (1.2), we tranfered it to interal equation and estimate its value. We consider the method of order reduction and Banach contarction to the integral equations.

Now we use the method of order reduction to transform (1.4) to a nonlinear integral equation; see [27]. To do this, let

$$(3.1) \quad T_1v(t) = I_{0+}^{\alpha-3}T_2v(t) = \int_0^1 G_1(t, s)v(s)ds,$$

$$(3.2) \quad T_2v(t) = \int_0^1 G_2(t, s)v(s)ds, \quad T_3v(t) = \int_0^1 G_3(t, s)v(s)ds,$$

where $G_1(t, s)$, $G_2(t, s)$ and $G_3(t, s)$ are given by (2.3), (2.4) and (2.5). From the above formulas, it follows that

$$D_{0+}D_{0+}D_{0+}^{\alpha-3}T_1v(t) = D_{0+}D_{0+}T_2v(t) = D_{0+}T_3v(t) = -v(t)$$

Note that since

$$T_1v(t) = \int_0^1 G_1(t, s)v(s)ds = \int_0^1 G(t, s)f(s)ds,$$

we have

$$T_1v(0) = T_1v(1) = 0.$$

Moreover by definition,

$$T_2v(1) = \int_0^1 G_2(1, s)v(s)ds = 0, \quad T_3v(1) = \int_0^1 G_3(1, s)v(s)ds = 0.$$

Boundary value problem (1.4) can be converted into a terminal value problem

$$D_{0+}v(t) = -f(t, T_1v(t), T_2v(t), T_3v(t), -v(t)), \quad v(1) = 0.$$

From the above formulas, it follows that

$$D_{0+}v(t) = f(t, T_1v(t), T_2v(t), T_3v(t), -v(t))$$

where

$$D_{0+}T_3v(t) = -v(t), \quad D_{0+}T_2v(t) = T_3v(t), \quad D_{0+}^{\alpha-3}T_1v(t) = T_2v(t).$$

Then we have the following lemmas.

Lemma 3. *Let $3 < \alpha \leq 4$. The boundary problem (1.4) is equivalent to the following integral equations forms;*

$$\begin{cases} v(t) = \int_t^1 f(s, T_1v(s), T_2v(s), T_3v(s), -v(s))ds, \\ T_1v(t) = \int_0^1 G_1(t, s)v(s)ds, \\ T_2v(t) = \int_0^1 G_2(t, s)v(s)ds, \\ T_3v(t) = \int_0^1 G_3(t, s)v(s)ds, \end{cases}$$

where $G_1(t, s)$, $G_2(t, s)$ and $G_3(t, s)$ are given by (2.3), (2.4) and (2.5).

Next we define an operator A from $C[0, 1]$ into $C[0, 1]$ by

$$(3.3) \quad Av(t) = \int_t^1 f(s, T_1v(s), T_2v(s), T_3v(s), v(s))ds,$$

where $v \in C[0, 1]$. Also let

$$(3.4) \quad (T_1v)(t) = \int_0^1 G_1(t, s)v(s)ds, \quad (T_2v)(t) = \int_0^1 G_2(t, s)v(s)ds,$$

$$(3.5) \quad (T_3v)(t) = \int_0^1 v(s)ds, \quad (T_4v)(t) = v(t).$$

Then the solution of boundary value problem (1.4) is equivalent to the fixed point of A on $C[0, 1]$. Take $u_0(t) = 1 - t$. By (3.4), we get

$$(3.6) \quad \begin{aligned} & \int_t^1 (T_1u_0)(t)dt \\ &= \frac{1}{\Gamma(\alpha)} \left(\frac{1}{\alpha(\alpha+1)} \frac{(1-t^{\alpha+1})}{1-t} - \frac{1}{2\alpha} \frac{(1-t^\alpha)}{1-t} + \frac{1}{2\alpha} \frac{(1-t^{\alpha-2})}{1-t} \right) u_0(t) \end{aligned}$$

In fact take $u_0(t) = 1 - t$. By (3.4), we get

$$\begin{aligned} (T_1u_0)(t) &= \int_0^t \frac{1}{\Gamma(\alpha-1)} (-(t-s)^{\alpha-2}(1-s) + t^{\alpha-2}(1-s)) ds \\ &+ \int_t^1 \frac{1}{\Gamma(\alpha-1)} (t^{\alpha-2} - t^{\alpha-2}s) ds \\ &= \int_0^t \frac{1}{\Gamma(\alpha-1)} (-(t-s)^{\alpha-1} + (t-s)^{\alpha-2}t - (t-s)^{\alpha-2}) \\ &+ \int_0^1 \frac{1}{\Gamma(\alpha-1)} (t^{\alpha-2} - t^{\alpha-2}s) ds \\ &= \frac{-1}{\alpha\Gamma(\alpha-1)} t^\alpha + \frac{1}{(\alpha-1)\Gamma(\alpha-1)} t^\alpha - \frac{1}{(\alpha-1)\Gamma(\alpha-1)} t^{\alpha-1} \\ &+ \frac{1}{\Gamma(\alpha-1)} \left(t^{\alpha-2} - \frac{1}{2} t^{\alpha-2} \right) \\ &= \frac{1}{\Gamma(\alpha+1)} t^\alpha - \frac{1}{\Gamma(\alpha)} t^{\alpha-1} + \frac{1}{2\Gamma(\alpha-1)} t^{\alpha-2} \end{aligned}$$

and

$$\begin{aligned}
& \int_t^1 (T_1 u_0)(s) ds \\
&= \int_t^1 \left(\frac{1}{\Gamma(\alpha+1)} s^\alpha - \frac{1}{\Gamma(\alpha)} s^{\alpha-1} + \frac{1}{2\Gamma(\alpha-1)} s^{\alpha-2} \right) dt \\
&= \frac{1}{\Gamma(\alpha+2)} (1-t^{\alpha+1}) - \frac{1}{\Gamma(\alpha+1)} (1-t^\alpha) + \frac{1}{2\Gamma(\alpha)} (1-t^{\alpha-1}) \\
&= \left(\frac{1}{\Gamma(\alpha+2)} \frac{1-t^{\alpha+1}}{1-t} - \frac{1}{\Gamma(\alpha+1)} \frac{1-t^\alpha}{1-t} + \frac{1}{2\Gamma(\alpha)} \frac{1-t^{\alpha-1}}{1-t} \right) u_0(t).
\end{aligned}$$

Since $1-t^3 < 1-t^\alpha < 1-t^4$, we have $\left(\frac{1-t^{\alpha+1}}{1-t}\right) < \frac{1-t^5}{1-t} = 1+t+t^2+t^3+t^4 \leq 5$. Thus we have

$$\int_t^1 (T_1 u_0)(t) dt \leq C_1 u_0(t)$$

where

$$(3.7) \quad C_1 = \left(\frac{5}{\Gamma(\alpha+2)} - \frac{3}{\Gamma(\alpha+1)} + \frac{3}{2\Gamma(\alpha)} \right).$$

We also have

$$\begin{aligned}
(T_2 u_0)(t) &= \int_0^t s(1-s) ds + \int_t^1 t(1-s) ds = \frac{t}{2} - \frac{t^2}{2} + \frac{t^3}{6} \\
\int_t^1 (T_2 u_0)(t) dt &= \left(\frac{1}{4}(1+t+t^2) - \frac{1}{6}(1+t+t^2+t^3) + \frac{1}{24}(1+t+t^2+t^3+t^4) \right) (1-t) \\
&\leq \frac{1}{6} u_0(t)
\end{aligned}$$

$$(T_3 u_0)(t) = \int_0^1 \frac{1}{2} (1-s)^2 ds = \frac{1}{2} (-t^2 + 3t - 1)$$

$$\int_t^1 (T_3 u_0)(s) ds = \int_t^1 \frac{1}{2} (1-s)^2 ds = \frac{1}{6} (1-t)^3 \leq \frac{1}{6} u_0(t).$$

Thus we have

$$(3.8) \quad \int_t^1 (T_1 u_0)(t) dt \leq M_1 u_0(t), \int_t^1 (T_2 u_0)(t) dt \leq M_2 u_0(t), \int_t^1 (T_3 u_0)(s) ds \leq M_3 u_0(t),$$

where

$$(3.9) \quad M_1 = \left(\frac{5}{\Gamma(\alpha+2)} - \frac{3}{\Gamma(\alpha+1)} + \frac{3}{2\Gamma(\alpha)} \right), M_2 = \frac{1}{6}, M_3 = \frac{1}{6}.$$

Next in order to use the method in [28], first we give the setting. For each number $M > 0$ we denote

$$D_M = \{(t, u, w, y, z) | 0 \leq t \leq 1, \|u\| \leq M_1 M, \|y\| \leq M_2 M, \|v\| \leq M_3 M, \|z\| \leq M\}$$

and by $B[O, M]$ we also denote a closed ball centered at O with the radius M in the space of continuous functions $C[0, 1]$ with the norm $\|\varphi\| = \max_{0 \leq t \leq 1} |\varphi(t)|$. In this case, by [28, Theorem 2.2], we have the following lemma;

Lemma 4. *Assume that there exist numbers $M, C_i > 0$ where $(i = 1, 2, 3, 4)$, such that*

$$(3.10) \quad |f(t, u, w, y, z)| \leq M,$$

$$(3.11) \quad |f(t, x_1, x_2, x_3, x_4) - f(t, y_1, y_2, y_3, y_4)| \leq \sum_{i=1}^4 C_i |x_i - y_i|$$

for any $(t, u, w, y, z), (t, x_1, x_2, x_3, x_4), (t, y_1, y_2, y_3, y_4) \in D_M$. Then the operator A defined by (3.3), where v_φ, u_φ are the solutions of the problems (1.4), maps the closed ball $B[O, M]$ into itself. Moreover, if

$$(3.12) \quad q := M_1 C_1 + M_2 C_2 + M_3 C_3 + C_4 < 1$$

then the operator A is contractive operator in $B[O, M]$.

Proof. We only need to prove that A defined by (3.3) has a unique fixed point in $B[O, M]$. We introduce a linear operator T on $C[0, 1]$ as

$$(3.13) \quad (Tu)(t) = \int_t^1 (M_1(T_1u)(s) + M_2(T_2u)(s) + M_3(T_3u)(s) + M_4(T_4u)(s)) ds.$$

As a first step, we show that for all $v \in C[0, 1]$ with $v(t) > 0$ ($t \in [0, 1]$), there exists $N = N(v)$ such that

$$(3.14) \quad (Tv)(t) \leq Nu_0(t), t \in [0, 1].$$

In fact, we take $N = M_1\|T_1v\| + M_2\|T_2v\| + M_3\|T_3v\| + M_4\|v\|$. Then by (3.13), we obtain the result. Moreover, it follows from (3.8) that

$$(3.15) \quad (Tu_0)(t) \leq qu_0(t), t \in [0, 1],$$

where

$$q = M_1 C_1 + M_2 C_2 + M_3 C_3 + C_4 < 1.$$

For any given $v_0 \in B[O, M]$, let

$$v_n(t) = (Av_{n-1})(t), w_n(t) = |v_n(t) - v_{n-1}(t)|, n = 1, 2, \dots$$

Since A is a operator on $B(O, M)$, for $t \in [0, 1]$, we have

$$\begin{aligned} w_{n+1}(t) &= |v_{n+1}(t) - v_n(t)| = |(Av_n)(t) - (Av_{n-1})(t)| \\ &\leq \int_t^1 |f(s, (T_1v_n)(s), (T_2v_n)(s), (T_3v_n)(s), -v_n(s)) \\ &\quad - f(s, (T_1v_{n-1})(s), (T_2v_{n-1})(s), (T_3v_{n-1})(s), -v_{n-1}(s))| ds \\ &\leq \int_t^1 (M_1 T_1(|v_n - v_{n-1}|)(s) + M_2 T_2(|v_n - v_{n-1}|)(s) \\ &\quad + M_3 T_3(|v_n - v_{n-1}|)(s) + M_4 |v_n(s) - v_{n-1}(s)|) ds = (T|v_n - v_{n-1}|)(t) = (Tw_n)(t). \end{aligned}$$

By (3.14), (3.15), and the method of induction, there exists $N = N(w_1)$ such that

$$w_{n+1}(t) \leq (Tw_n)(t) \leq \dots \leq (T^n w_1)(t) \leq N(T^{n-1} u_0)(t) \leq NM^{n-1} u_0(t), t \in [0, 1].$$

Thus for all $m, n \in N$ and $t \in [0, 1]$,

$$\begin{aligned} |v_{n+m+1}(t) - v_n(t)| &= |v_{n+m+1}(t) - v_{n+m}(t) + \cdots + v_{n+1}(t) - v_n(t)| \\ &\leq w_{n+m+1}(t) + \cdots + w_{n+1}(t) \\ &\leq NM^{n+m-1}u_0(t) + \cdots + Nq^{n-1}u_0(t) \\ &= \frac{NM^{n-1}(1 - q^{m+1})}{1 - q}u_0(t) < \frac{Nq^{n-1}}{1 - q}. \end{aligned}$$

This shows that $\{v_n\}$ is a uniform Cauchy sequence in $B[O, M]$ and since $C[0, 1]$ is complete and $B[O, M]$ is a clozed subspace of $C[0, 1]$, $B[O, M]$ is also complete. Then there exists $v^* \in B[O, M]$ such that $\lim_{n \rightarrow \infty} v_n = v^*$. Moreover, v^* is a fixed point of A that follows from the continuity of A . Next we show that A has at most one fixed point. Suppose that there are two elements $x, y \in B[O, M]$ with $x = Ax$ and $y = Ay$. By (3.14), there exists N such that $(T(|x - y|))(t) \leq Nu_0(t)$, $t \in [0, 1]$. Then for $n \in N$, we have

$$\begin{aligned} |x(t) - y(t)| &= |(A^n x)(t) - (A^n y)(t)| \leq (T^n(|x - y|))(t) \\ &\leq N(T^{n-1}u_0)(t) \leq Nq^{n-1}u_0(t), t \in [0, 1]. \end{aligned}$$

Consequently, we assert that $x = y$. This means that A has at most one fixed point. This completes the proof. \square

Then we have the following theorem.

Theorem 5. *Under the assumption of lemma 4, the boundary value problem (1.4) has a unique solution there hold the estimates*

$$(3.16) \quad \|u\| \leq M_1 M, \|D_{0+}^{\alpha-3}u\| \leq M_2 M, \|D_{0+}^{\alpha-2}u\| \leq M_3 M, \|D_{0+}^{\alpha-1}u\| \leq M.$$

Proof. The proof is similar to that of [28, Theorem 2.2]. \square

Next for the positive solution case, we have the following, For each number $M > 0$ we denote

$$D_M^+ = \left\{ (t, u, y, v, z) \left| \begin{array}{l} 0 \leq t \leq 1, 0 \leq u \leq M_1 M, 0 \leq y \leq M_2 M, \\ 0 \leq v \leq M_3 M, -M \leq z \leq 0 \end{array} \right. \right\}$$

and

$$S_M = \{\varphi \in C[0, 1] \mid 0 < \varphi(t) \leq M \text{ for any } t \in [0, 1]\}.$$

Then as the special case of Theorem 5, we have the following theorem.

Theorem 6. *Suppose that in D_M^+ the function f is such that*

$$(3.17) \quad 0 \leq f(x, u, y, v, z) \leq M,$$

and satisfies the Lipschitz condition (3.11) and condtion (3.12). Then, the problem (1.4) has a unique nonnegative solution.

4. EXAMPLES

For the examples in this section, as the constant value we apply the following. If $\alpha = 3.1$, then $M_1 = \frac{5}{\Gamma(3.1+2)} - \frac{3}{\Gamma(3.1+1)} + \frac{3}{2\Gamma(3.1)} = 0.421205$, $M_2 = \frac{1}{6} = 0.166667$, $M_3 = \frac{1}{6} = 0.166667$, and $M_4 = 1$.

Example 7. Next we consider the examples of fractional order given by a non-linear f .

$$\begin{cases} D_{0+}^{3.1}u(t) = -\frac{1}{3}D_{0+}^{0.1}u(t)D_{0+}^{2.1}u(t) + u(t)(D_{0+}^{1.1}u(t))^2 + \frac{1}{2}e^{-u(t)}, 0 < t < 1, \\ u(0) = D^{0.1}u(0) = D_{0+}^{1.1}u(1) = D_{0+}^{2.1}u(1) = 0, \end{cases}$$

In this example

$$f(x, u, y, v, z) = -\frac{1}{3}yz + uv^2 + \frac{1}{2}e^{-u},$$

which maps $[0, 1] \times R_+^3 \times R^- \rightarrow R_+$.

If we choose $M = 1$, then we have $\frac{1}{3}M_2MM_4M + M_1MM_3M^2 + \frac{1}{2}e^{-M_1M} \leq M$.

Therefore, if $\alpha = 3.1$, the Lipschitz coefficients in Lemma ?? are $C_1 = |f_u| = |v^2 - \frac{1}{2}e^{-u}| \leq |(\frac{1}{6}M)^2 - \frac{1}{2}e^{-M_1M}| \approx 0.30035$, $C_2 = |f_y| = \frac{1}{3}z = \frac{1}{3}M_4M = \frac{1}{3}$, $C_3 = |f_v| = 2uv \leq \frac{1}{3}M_1M^2 \approx 0.140402$, $C_4 = |f_z| = \frac{1}{3}y = \frac{1}{3}M_2M \approx .0555556$. Then we have

$$q := (C_1 * M_1 + C_2 * M_2 + C_3 * M_3 + C_4 * M_4) \approx 0.26102 < 1.$$

Then the conditions of Lemma 4 and Theorem 5 are satisfied.

Next we consider the examples of fractional order given by a non-linear f . They are versions in [28].

Example 8.

$$\begin{cases} D_{0+}^{3.1}u(t) = \frac{u(t)}{6}(u(t) + D_{0+}^{0.1}u(t) + D_{0+}^{1.1}u(t) - D_{0+}^{2.1}u(t)) + 1, 0 < t < 1, \\ u(0) = D^{0.1}u(0) = D_{0+}^{1.1}u(1) = D_{0+}^{2.1}u(1) = 0, \end{cases}$$

In this example

$$f(x, u, y, v, z) = \frac{u}{6}(u + y + v - z) + 1,$$

which maps $[0, 1] \times R_+^3 \times R^- \rightarrow R_+$. We can choose $M = 2$, and therefore, if $\alpha = 3.1$, the Lipschitz coefficients in Lemma ?? are satisfied. By the definition of f , since

$$|f_u| = \frac{1}{3} * u < \frac{2}{3} * M_1 \approx 0.280803,$$

$$f_y = f_v = f_z = \frac{2}{6} * M_1 \approx 0.140402,$$

we have

$$\begin{aligned} q &:= M_1C_1 + M_2C_2 + M_3C_3 + M_4C_4 \\ &= M_1 * \frac{2}{3}M_1 + M_2 * \frac{1}{3}M_1 + \frac{1}{2} * \frac{1}{3}M_1 + \frac{1}{3}M_1 \\ &\approx 0.305478 < 1. \end{aligned}$$

Thus the conditions of Lemma 2 and Theorem 2 are satisfied.

Example 9.

$$\begin{cases} D_{0+}^{3.1}u(t) = \frac{u(x) + D_{0+}^{1.1}u(x)}{1 + u(x)^2 + D_{0+}^{2.1}u(x)^2 + D_{0+}^{1.1}u(x)^2 + D_{0+}^{0.1}u(x)^2} + e^{-u^2}, 0 < x < 1, \\ u(0) = D^{0.1}u(0) = D_{0+}^{1.1}u(1) = D_{0+}^{2.1}u(1) = 0, \end{cases}$$

In this example

$$f(x, u, y, v, z) = \frac{u + y}{1 + u^2 + y^2 + v^2 + z^2} + e^{-u^2},$$

which maps $[0, 1] \times R_+^3 \times R^- \rightarrow R_+$. Analogously we can choose $M = 1$ and Following argument, the function $f(x, u, y, v, z)$ satisfies the condition of Lemma 4 and Theorem 5. Hence, the problem has a unique solution. In fact if we choose $\alpha = 3.1$, then the Lipschitz coefficients in Lemma 4 are given by the following;

$$\begin{aligned} |f_u| &= \left| \frac{(1 + u^2 + y^2 + v^2 + z^2) - 2u(u + y)}{(1 + u^2 + y^2 + v^2 + z^2)^2} - 2ue^{-u^2} \right| \\ &= \left| \frac{1}{(1 + u^2 + y^2 + v^2 + z^2)} - \frac{2u(u + y)}{(1 + u^2 + y^2 + v^2 + z^2)^2} - 2ue^{-u^2} \right| \\ &= \left| \frac{1}{(1 + (M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)} - \frac{2M_1(M_1 + M_2)M^2}{(1 + (M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)^2} - 2M_1Me^{-(M_1M)^2} \right|, \end{aligned}$$

$$\begin{aligned} |f_y| &= \left| \frac{1}{(1 + u^2 + y^2 + v^2 + z^2)} - \frac{2y(u + y)}{(1 + u^2 + y^2 + v^2 + z^2)^2} \right| \\ &= \left| \frac{1}{((1 + (M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)} - \frac{2M_2(M_1 + M_2)M^2}{(1 + (1 + M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)^2} \right|, \end{aligned}$$

$$|f_v| = \left| -\frac{-2v(u + y)}{(1 + u^2 + y^2 + v^2 + z^2)^2} \right| = \left| \frac{2M_3(M_1 + M_2)M^2}{(1 + (1 + M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)^2} \right|,$$

$$|f_z| = \left| -\frac{-2z(u + y)}{(1 + u^2 + y^2 + v^2 + z^2)^2} \right| = \left| \frac{2M_4(M_1 + M_2)M^2}{(1 + (1 + M_1^2 + M_2^2 + M_3^2 + M_4^2)M^2)^2} \right|.$$

As we take $M = 2$, we have

$$|f_u| \approx 0.4430161, |f_y| \approx 0.360078, |f_v| \approx 0.0877564, |f_z| \approx 0.526538,$$

$$q := M_1 * |f_u| + M_2 * |f_y| + M_3 * |f_v| + M_4 * |f_z| \approx 0.787778 < 1.$$

REFERENCES

- [1] R. P. Agarwal and B. Ahmad, *Existence theory for anti-periodic boundary value problems of fractional differential equations and inclusions*, Comput. Math. Appl. 62 (2011), 1200–1214.
- [2] E. Alves, T. F. Ma, M. L. Pelicer, *Monotone positive solutions for a fourth order equation with nonlinear boundary conditions*, Nonlinear Anal., 71 (2009) 3834–3841.
- [3] C. Bai, *Triple positive solutions for a boundary value problem of nonlinear fractional differential equation*, Electron. J. Qual. Theory Differ. Equ., 24 (2008), 1–10.
- [4] Z. Bai and H. Lü, *Positive solutions for boundary value problem of nonlinear fractional differential equation*, J. Math. Anal. Appl., 311 (2005), 495–505.
- [5] G. Chai, *Existence results of positive solutions for boundary value problems of fractional differential equations*, Bound. Value Probl., 2013, 2013:109.
- [6] K. Diethelm and N. J. Ford, *Analysis of fractional differential equations*, J. Math. Anal. Appl., 265 (2002), 229–248.
- [7] C. P. Gupta, *Existence and uniqueness results for the bending of an elastic beam equation at resonance*, J. Math. Anal. Appl., 135 (1988), 208–225.
- [8] C. P. Gupta, *A nonlinear boundary value problem associated with the static equilibrium of an elastic beam supported by sliding clamps*, Int. J. Math. Math. Sci., 12 (1989), 697–711.

- [9] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, In North-Holland Mathematics Studies, 204, Elsevier Science B.V., Amsterdam, 2006.
- [10] S. Liang and J. Zhang, *Positive solutions for boundary value problems of nonlinear fractional differential equation*, *Nonlinear Anal.*, 71 (2009), 5545–5550.
- [11] R. Liu and R. Ma, *Existence of positive solutions for an elastic beam equation with nonlinear boundary conditions*, *J. Appl. Math.*, 2014 (2014), Article ID 972135.
- [12] R. Ma, *Multiple positive solutions for a semipositone fourth-order boundary value problem*, *Hiroshima Math. J.*, 33 (2003), 217–227.
- [13] R. Ma, *Existence of positive solutions of a fourth-order boundary value problem*, *Appl. Math. Comput.*, 168 (2005), 1219–1231.
- [14] T. F. Ma and J. da Silva, *Iterative solutions for a beam equation with nonlinear boundary conditions of third order*, *Appl. Math. Comput.*, 159 (2004), 11–18.
- [15] N. Nyamoradi and M. Javidi, *Existence of multiple positive solutions for fractional differential inclusions with m -point boundary conditions and two fractional orders*, *Electron. J. Differential Equations*, 187 (2012), 1–26.
- [16] S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Translated from the 1987 Russian original, Gordon and Breach Science Publishers, Switzerland, 1993.
- [17] M. Toyoda and T. Watanabe, *Existence and uniqueness theorem for fractional order differential equations with boundary conditions and two fractional order*, *J. Nonlinear Convex Anal.*, 17 (2016), 267–273.
- [18] M. Toyoda and T. Watanabe, *Note on solutions of boundary value problems involving a fractional differential equation*, *Linear and Nonlinear Analysis Volume 3*, Number 3, 2017, 449–455.
- [19] R. A. Usmani, *A uniqueness theorem for a boundary value problems*, *Proc. Amer. Math. Soc.*, 77 (1979), 329–335.
- [20] R. A. Usmani, *Finite difference methods for computing eigenvalues of fourth order boundary value problems*, *Int. J. Math. Math. Sci.*, 9 (1986), 137–143.
- [21] J. R. L. Webb, G. Infante and D. Franco, *Positive solutions of nonlinear fourth-order boundary-value problems with local and non-local boundary conditions*, *Proc. Roy. Soc. Edinburgh Sect. A*, 138 (2008), 427–446.
- [22] X. Xu, D. Jiang and C. Yuan, *Multiple positive solutions for the boundary value problem of a nonlinear fractional differential equation*, *Nonlinear Anal.*, 71 (2009), 4676–4688.
- [23] B. Yang, *Positive solutions for a fourth order boundary value problem*, *Electron. J. Qual. Theory Differ. Equ.*, 3 (2005), 1–17.
- [24] Y. Yang, *Fourth-order two-point boundary value problems*, *Proc. Amer. Math. Soc.*, 104 (1988), 175–180.
- [25] Y. Zoua, Y. Cuib, *Uniqueness result for the cantilever beam equation with fully nonlinear term*, *J. Nonlinear Sci. Appl.*, 10 (2017), 4734–4740.
- [26] D. Quang A., N. Quyb, *Existence results and iterative method for solving the cantilever beam equation with fully nonlinear term*, *Nonlinear Anal:Real World Applications*, 36 (2017) 56–68.
- [27] Y. Li, *Existence of positive solutions for the cantilever beam equations with fully nonlinear terms*, *Nonlinear Anal. RWA* 27 (2016) 221–237.
- [28] E.R. Kaufmann, N. Kosmatov, *Elastic beam problem with higher order derivatives*, *Nonlinear Anal. Real World Appl.* 8 (2007) 811–821.
- [29] Y. Guo, Y. Fei, Y. Liang, *Positive solutions for nonlocal fourth-order boundary value problems with all order derivatives*, in: *Boundary Value Problems 2012*?29, 2012, 12 pp.
- [30] F. Minhos, T. Gyulov, A.I. Santos, *Lower and upper solutions for a fully nonlinear beam equation*, *Nonlinear Anal.* 71 (2009) 281–292.
- [31] A. Cabada, J. Fialho, M. Minhos, *Extremal solutions to fourth order discontinuous functional boundary value problems*, *Math. Nachr.* 286 (2013) 1744–1751.

(Toshikazu Watanabe) MEIJI UNIVERSITY
 Email address: wa-toshi@mti.biglobe.ne.jp