

EXISTENTIALLY CLOSED FIELDS WITH DIFFERENCE/DIFFERENTIAL OPERATORS

KAI INO

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MANCHESTER

ABSTRACT. We survey separably differentially closed fields and separable differential closure in comparison with differentially closed fields and differential closure. Also we observe several topics around the theory of separably differentially closed fields.

CONTENTS

1. Differentially closed fields and differential closure	1
2. Separably differentially closed fields and separable differential closure	3
3. Differential large fields with stable theory	6
4. Difference-differential fields	6
References	7

1. DIFFERENTIALLY CLOSED FIELDS AND DIFFERENTIAL CLOSURE

In this section, we describe classical results for differentially closed fields and differential closure in characteristic zero. We fix a differential field (K, δ) is of characteristic zero where a derivation δ on K is an additive homomorphism satisfying Leibniz rule: $\delta(ab) = a\delta(b) + b\delta(a)$. We denote a differential polynomial ring over (K, δ) in a differential indeterminate x by $K\{x\}$.

A. Robinson [40], in 1959, introduced the notion of differentially closed fields as analogous to algebraically closed fields of characteristic zero; every finite system of algebraic differential equations and inequations over K with a solution in an differential extension of K already has solution in K . One can show that every differential fields of characteristic zero has a differentially algebraic extension that is differentially closed. He axiomatized them and showed that the theory of differentially closed fields is complete, model complete and decidable. Through the notion of existential closedness, We may also define a differential field (K, δ) being *differentially closed* if it is existentially closed in every differential field extension. (Recall that a field K is algebraically closed if it is existentially closed in every field extension.)

L. Blum [1] simplified Robinson's axiom. One can observe the shape of the prime differential ideals for an ingredient for this; Let P is a nonzero prime differential

ideal of $K\{x\}$. For any irreducible f of minimal rank in P , we have $P = [f] : s_f^\infty$ where a *separant* of f , s_f , is obtained by taking the partial derivative of f with respect to its highest order variable, and $[f]$ denote the differential ideal generated by f , and for an ideal I of $K\{x\}$ and $s \in K\{x\}$, the saturated ideal of I over s is defined as $I : s^\infty = \{h \in K\{x\} : s^m h \in I \text{ for some } m \geq 0\}$. Conversely, if $f \in K\{x\}$ is irreducible, then $[f] : s_f^\infty$ is a prime differential ideal and f is of minimal rank in it. In conclusion, Blum [1, §III] showed that a differential field (K, δ) is differentially closed if and only if for any nonzero $f, g \in K\{x\}$ and $\text{ord } g < \text{ord } f$, there exists $a \in K$ such that $f(a) = 0$ and $g(a) \neq 0$ where this scheme of conditions is expressed in a first-order fashion and denoted by DCF_0 .

E.R. Kolchin [25] introduced the notion called constrained elements and constrained extension since every differential closed fields has a nontrivial differentially algebraic extension unlike the property of algebraically closed fields; let $\alpha = (\alpha_i)_{i \in I}$ be a family of elements from a differential field extension of K , and then α is said to be *constrained* over K if there exists $g \in K\{\bar{x}\}$ with $g(\bar{\alpha}) \neq 0$ such that for every differential specialisation β of α over K if $g(\beta) \neq 0$ then β is a generic differential specialisation. Kolchin [25] defined a differential field extension of K to be a constrained extension if every finite family of elements from it is constrained over K , and K having no proper constrained extension is called *constrainedly closed*. (Remark that this Kolchin's definition is for Characteristic zero is slightly different from the case for positive characteristic. See next Section). Note that every constrained extension is differentially algebraic, but the converse is not generally true. In conclusion, he showed that K is differentially closed if and only if K is constrainedly closed.

D. Pierce and A. Pillay [37] reworked the axiom of differentially closed fields in a geometric setting in terms of prolongations of affine algebraic varieties; namely, given an algebraic variety V over K there exists an algebraic bundle $\pi : \tau V \rightarrow V$ over K called that prolongation of V that has the following characteristic property: for any differential field extension (L, δ) of (K, δ) if $a \in V(L)$ then $(a, \delta a) \in \tau V(L)$. In the case when V is affine, say $K[V] = K[x_1, \dots, x_n]/I$ with I a radical ideal, the following equations define the first prolongation of V , τV ,

$$f(\bar{x}) = 0 \quad \text{and} \quad \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\bar{x}) \cdot y_i + f^\delta(\bar{x}) = 0,$$

as f varies in generators of I , the defining ideal of V .

Their geometric axiom has been adapted to other settings; for instance, fields of characteristic zero with several commuting derivations [30], fields of positive characteristic with derivation of Frobenius [15, 26], difference-differential fields of characteristic zero [4], or with several commuting derivations [29, 16].

In the end, we give a differential version of the Basis Theorem in terms of radical differential ideals; namely, every radical differential ideal in $K\{\bar{x}\}$ is finitely generated. To sum up these works by A. Robinson, L. Blum, E.R. Kolchin, D. Pierce and A. Pillay, we have the following:

Fact 1.1. *Let (K, δ) be a differential field of characteristic zero. Then, the following are equivalent:*

- (1) (K, δ) is differentially closed.

- (2) (K, δ) is existentially closed in every differentially algebraic extension.
- (3) (K, δ) is constrainedly closed.
- (4) for each n and every prime differential ideal P of $K\{x_1, \dots, x_n\}$, if $g \in K\{x_1, \dots, x_n\} \setminus P$ then there is $\bar{a} \in K^n$ such that $f(\bar{a}) = 0$ for all $f \in P$ and $g(\bar{a}) \neq 0$.
- (5) for any nonzero $f, g \in K\{x\}$ and $\text{ord } g < \text{ord } f$, there exists $a \in K$ such that $f(a) = 0$ and $g(a) \neq 0$.
- (6) let V and W be K -irreducible affine varieties over K with $W \subseteq \tau V$ and $\pi|_W : W \rightarrow V$ dominant. If O_V and O_W are nonempty Zariski-open subsets over K of V and W , respectively, then there is a K -rational point $a \in O_V$ such that $(a, \delta a) \in O_W$.

Differential closure is defined as analogous to algebraic closure; $\bar{K} \supseteq K$ is a *differential closure* of K if \bar{K} is differentially closed and for any differentially closed $L \supseteq K$, there exists a differential K -embedding $f : \bar{K} \rightarrow L$. L. Blum observed a prime model extension in DCF_0 is exactly differential closure, and she also showed that the theory DCF_0 is ω -stable. In general, in ω -stable theory, a prime model extension exists [35] and it is unique [43]. However, a differential closure does not have exactly the same properties as an algebraic closure. S. Shelah [44], E. R. Kolchin [25] and M. Rosenlicht [41] independently showed that a differential closure is not minimal in general, that is, there exists a differential closed field L such that $K \subseteq L \subset \bar{K}$. For instance, the differential closure of \mathbb{Q} is not minimal.

Strong minimality of the solution set of algebraic differential equations has been studied with many successful applications to functional transcendence results. A differential equation over (K, δ) , $f(x) = 0$ with $\text{ord } f = n$, is strongly minimal if f is absolutely irreducible and for all element a with $f(a) = 0$ and any differential field extension (L, δ) of (K, δ) , we have

$$\text{tr.deg}(L\langle a \rangle/L) = 0 \text{ or } n.$$

For instance, the differential equation satisfied by the j -function [14], Painlevé equations with generic coefficients [36], equations satisfied by Γ -automorphic functions on upper half-plane for Γ a Fuchsian group of first kind [5], to name a few (see also [12]), are all strongly minimal. Also Zilber's Trichotomy in DCF_0 for strongly minimal sets is one of the reason for strong minimality to have deep applications; for instance, differential Galois theory [38], the proof of Modell-Lang conjecture for function fields [18], (Manin-Manford conjecture [19] in the context of difference field) among others.

2. SEPARABLY DIFFERENTIALLY CLOSED FIELDS AND SEPARABLE DIFFERENTIAL CLOSURE

In this section, we describe separably differentially closed fields and separable differential closure. We fix a differential field (K, δ) is of arbitrary characteristic p .

As differentially closed fields of characteristic zero and their model-theoretic properties provide us a suitable universal domain to study for the solution sets of algebraic differential equations, differential Galois theory, differential algebraic group, differential algebraic geometry and its application to number theory, we expect that separably differentially closed fields have the right universal domain for the case of those in positive characteristic. (Remark that C. Wood established

the foundation of the theory of differentially closed fields of positive characteristic, and the differential closures over differential perfect fields where we recall that K is differentially perfect if either its characteristic is zero or the p -th power of K coincides with the constants of K , that is, $[C_K : K^p] = 1$.

Separably differentially closed fields were introduced by the author and O. León Sánchez in [20, 21] as a differential analogous to separably closed fields; (K, δ) is said to be *separably differentially closed* if it is existentially closed in every differential field extension that is separable as a field. In other words, if (L, δ) is a differential extension of (K, δ) and L/K is separable, then (K, δ) is existentially closed in (L, δ) .

In the spirit of L.Blum, we characterised them by analysing the shape of separable prime differential ideals; if P is a nonzero separable prime differential ideal of $K\{x\}$, then $P = [f] : s_f^\infty$ for any irreducible f of minimal rank in P . Conversely, if $f \in K\{x\}$ is irreducible with $s_f \neq 0$, then $[f] : s_f^\infty$ is a separable prime differential ideal and f is of minimal rank in it. In conclusion, we showed that (K, δ) is separably differentially closed if and only if for any nonzero $f, g \in K\{x\}$ with $s_f \neq 0$ and $\text{ord } g < \text{ord } f$, there exists $a \in K$ such that $f(a) = 0$ and $g(a) \neq 0$, where this scheme of conditions is expressed in a first-order fashion and denoted by SDCF.

Kolchin's constrained elements [24] (in arbitrary characteristic) are given as follows; let $\bar{\alpha}$ be an n -tuple from a differential field extension of K . We say that $\bar{\alpha}$ is *constrained* over K if $K\langle\bar{\alpha}\rangle/K$ is separable and there exists $g \in K\{\bar{x}\}$ with $g(\bar{\alpha}) \neq 0$ such that for every differential specialisation $\bar{\beta}$ of $\bar{\alpha}$ over K , with $K\langle\bar{\beta}\rangle/K$ separable, if $g(\bar{\beta}) \neq 0$ then $\bar{\beta}$ is a generic differential specialisation. Then our definition [20, 21] of being *constrained closed* (in arbitrary characteristic) is that for every finite tuple \bar{a} from a differential field extension, if \bar{a} is constrained over K then each entry of \bar{a} is in K . We take this definition which is different from the case of characteristic zero because $K\langle\bar{\alpha}\rangle/K$ is not generally constrained over K for every $\bar{\alpha}$ constrained over K (4. [21]). But this definition is reduced to the Kolchin's definition [25] if K is of characteristic zero.

In the manner of D. Pierce and A. Pillay, we characterised SDCF in a geometric setting by adding separability conditions; namely, given K -irreducible varieties V and W , we say that V is separable if the function field $K(V)$ is separable over K . We say that a morphism $\phi : W \rightarrow V$ over K is separable if it is dominant and the function field $K(W)$ is separable over the function field $K(V)$.

Before summarizing our results, we make some comments for Differential Basis Theorem in positive characteristic. Y.Ershov [13] showed that the completions of the theory of separably closed fields are determined by specifying the degree of imperfection e ; that is, the degree of K over K^p , $[K : K^p] = p^e$. In my Thesis [20], all the work has been done with the restriction of what we call *differential degree of imperfection* being finite, $[C_K : K^p] < \infty$. Thus the following Basis theorem are used through the Thesis [24, §III]; namely, every radical differential ideal in $K\{\bar{x}\}$ is finitely generated if and only if $[C_K : K^p]$ is finite. However, in the paper [21], we extended the work including infinite differential degree of imperfection, $[C_K : K^p] = \infty$, then we have the following Basis Theorem [24, §III]; every separable prime differential ideal of $K\{\bar{x}\}$ is finitely generated as a radical differential ideal.

To sum up, separably differentially closed fields are characterised as the following [20, 21].

Theorem 2.1. *Let (K, δ) be a differential field of arbitrary characteristic. Then, the following are equivalent:*

- (1) (K, δ) is separably differentially closed.
- (2) for every differentially algebraic extension (L, δ) of (K, δ) , if L/K is separable (as fields) then (K, δ) is existentially closed in (L, δ) .
- (3) (K, δ) is constrainedly closed.
- (4) for each n and every separable prime differential ideal P of $K\{x_1, \dots, x_n\}$, if $g \in K\{x_1, \dots, x_n\} \setminus P$ then there is $\bar{a} \in K^n$ such that $f(\bar{a}) = 0$ for all $f \in P$ and $g(\bar{a}) \neq 0$.
- (5) For any nonzero $f, g \in K\{x\}$ with $s_f \neq 0$ and $\text{ord } g < \text{ord } f$, there exists $a \in K$ such that $f(a) = 0$ and $g(a) \neq 0$.
- (6) Let V and W be K -irreducible affine varieties over K with $W \subseteq \tau V$, W separable, and $\pi|_W : W \rightarrow V$ separable. If O_V and O_W are nonempty Zariski-open subsets over K of V and W , respectively, then there is a K -rational point $a \in O_V$ such that $(a, \delta a) \in O_W$.

Remark that the theorem above is reduced to the case of characteristic zero by the following points; namely, if K is characteristic zero, then

- (i) Separably differentially closed fields are differentially closed fields.
- (ii) Separable prime differential ideals are prime differential ideals
- (iii) Constrainedly closed differential fields and constrained elements (by definition in this section) are constrainedly closed differential fields and constrained elements (by the definition in previous section)

The completions of the theory of separably differentially closed fields are determined by specifying the differential degree of imperfection ϵ and characteristic p . Adding differential λ -functions as differential analogous to its algebraic one, one can show that the theory is stable.

Given (K, δ) a differential field of positive characteristic p with a differential degree of imperfection ϵ as analogous to the separable closure, its suitable closure should preserve same differential degree of imperfection. Then $\bar{K} \supseteq K$ is said to be a *separable differential closure* of K if \bar{K} is separably differentially closed with differential degree of imperfection ϵ and for any separably differentially closed extension L of K with the differential degree of imperfection ϵ , there exists a differential K -embedding $f : \bar{K} \hookrightarrow L$.

The existence of the separable differential closure is guaranteed by the constrained construction [21, chapter 5]. We showed that the existence and uniqueness of prime model extension over each differential fields of positive characteristic p with a differential degree of imperfection ϵ [20, 21]. Since by quantifier elimination in $\text{SDCF}_{p,\epsilon}^\ell$, any embeddings of the model is elementary. Thus, the existence of a prime model over K is deduced by the existence of a separable differential closure. The results that the theory is stable and countable implies the uniqueness of the prime model extensions [21, chapter 6].

One can naturally ask if a separable differential closure is minimal in general (for instance, the minimality of a (separable) differential closure of \mathbb{F}_p). Also we have a question for a version of Zilber's Dichotomy in SDCF . (Remark that if (K, δ)

is separably differentially closed fields, then K and C_K are separably closed fields with infinite differential degree of imperfection [21], but we only know that Zilber's Dichotomy holds for the case of finite degree of imperfection [18].)

3. DIFFERENTIAL LARGE FIELDS WITH STABLE THEORY

Infinite ω -stable fields and infinite superstable fields are algebraically closed by A. Macintyre [32] and G. Cherlin and S. Shelah [10] respectively. C. Wood [47] showed that separably closed fields are stable, and conversely the famous conjecture that "Infinite stable fields are separably closed" has remained open for a long time. (Remark that T. Scalon showed that infinite stable fields are Artin-Schreier closed [23]). Several special cases of the conjecture have been shown; for instance, W. Jonson, C. Tran, E. Walsberg and J. Ye [22] showed that a large stable field is separably closed where we recall that a field K is large if every irreducible variety over K with smooth K -rational point has Zariski dense set of K -rational points.

O. León Sánchez and M. Tressl [31] introduced the notion of differentially largeness in characteristic zero as an analogue to largeness of fields; a differential field (K, δ) of characteristic zero is differentially large if it is large and (K, δ) is existentially closed in every differential extension in which K is existentially closed as a field. The author and W. Mikolaj conjectured that every differential large stable fields is separably differentially closed. O. León Sánchez's (unpublished) work suggested that a differential field (K, δ) of arbitrary characteristic is differentially large if and only if it is large and (K, δ) is existentially closed in every differential extension in which K is existentially closed as a field. He also axiomatised them as follows; (K, δ) is differentially large if and only if for every pair of nonzero (f, g) in $K\{x\}$ with $S_f \neq 0$ and $\text{ord } f > \text{ord } g$, if $f = 0$ and $gS_f \neq 0$ has a algebraic solution in K , then $f = 0$ and $g \neq 0$ has a differential solution in K . Then one can easily show that every differentially large stable field is separably differentially closed; assuming that (K, δ) is differentially large and stable. Since an infinite stable fields is separably closed, K is separably closed, and (K, δ) is differentially large, we have (K, δ) being existentially closed in every differential field extension which is separable, that is, (K, δ) is separably differentially closed.

Corollary 3.1. *Every differentially large stable field is separably differentially closed.*

4. DIFFERENCE-DIFFERENTIAL FIELDS

R.Cohn [6], in 1970, studied difference-differential algebra, and then Hardouin and Singer [17] studied linear difference-differential Galois theory. E. Hrushovski showed the existence of model companion of the theory of difference-differential fields in characteristic zero and R. Bustamante [3, 4] studied their algebraic and model-theoretic properties. These studies are extended to the partial differential case by O. León Sánchez [29] (see also [16]). The theory for both ordinary and partial cases is supersimple and satisfy Zilber's Dichotomy which is shown by using proper jet spaces [39] (see also [34]).

Let (K, δ) be a differential field of characteristic zero. A difference-differential field (K, δ, σ) is a differential field (K, δ) with an automorphism σ which commutes with the derivation δ . A version of Basis Theorem for difference-differential fields is

the following; every perfect (δ, σ) -ideal is finitely generated where a perfect (δ, σ) -ideal I is a differential ideal and $f^m \sigma(f)^n \in I$ implies $f \in I$.

We now introduce two types of the form of model companion of the theory of difference-differential fields in the spirit of R. Bustamante and O. León Sánchez with restriction to the ordinal differential case.

Fact 4.1. *Let (K, δ, σ) be a difference-differential field of characteristic zero. Then, the following are equivalent:*

- (1) (K, δ, σ) is existentially closed.
- (2) Suppose that (K, δ) is differentially closed. Let V and W be K -irreducible δ -varieties over K with $W \subseteq V \times V^\sigma$ such that W projects generically both onto V and V^σ . If O_V and O_W are nonempty δ -open subsets over K of V and W , respectively, then there is a K -rational point $a \in O_V$ such that $(a, \delta a) \in O_W$.
- (3) Suppose that (K, δ) is differentially closed. Let U, V and W be K -irreducible affine varieties over K with $U \subseteq V \times V^\sigma$ and $W \subseteq \tau V$ such that U projects generically both onto V and V^σ and W projects generically onto U . If $\pi_1(W)^\sigma = \pi_2(W)$ and δ -generic point of W projects onto a δ -generic point of $\pi_1(W)^\sigma$ and onto δ -generic point of $\pi_2(W)$, then there is a K -rational point $a \in V$ such that $(a, \sigma a) \in U$ and $(a, \delta a, \sigma a, \sigma \delta a) \in W$ where we let $\pi_1 : \tau(V) \times \tau(V)^\sigma \rightarrow \tau(V)$ and $\pi_2 : \tau(V) \times \tau(V)^\sigma \rightarrow \tau(V)^\sigma$, with which identify $\tau(V \times V^\sigma)$.

Note that in the notations of (3) in the fact above, we say that a point a in an extension of K is a δ -generic of W over K if $(a, \delta a)$ is a generic of W over K , and $\text{tr.deg}(\delta^i a/K(a, \dots, \delta^{i-1} a)) = \text{tr.deg}(\delta a/K(a))$ for all $i > 1$. We denote by V^σ the δ -variety obtained by applying σ to the coefficients of the differential polynomials defining V . Remark that the statement of (2) above is described by differential variety, but that of (3) is by algebraic variety.

One can naturally ask if the theory of separably differentially closed fields with a generic differential automorphism has model companion. We note that Z. Chatzidakis [8] showed that in the case of separably closed fields, such model companion exists.

REFERENCES

- [1] L. Blum Generalized algebraic structures: A model theoretic approach. Ph.D. Thesis, Massachusetts Institute of Technology, 1968.
- [2] L. Blum. Differentially Closed Fields: A Model Theoretic Tour. Contributions to Algebra, Academic Press Inc., 1977.
- [3] R. Bustamante. Théorie des modèles des corps différentiellement clos avec un automorphisme générique. PhD Thesis. Université Paris 7, 2005.
- [4] R. Bustamante Medina. Differentially closed fields of characteristic zero with a generic automorphism. Revista de Matemática: Teoría y Aplicaciones, 14(1):81-100, 2007.
- [5] Guy Casale, James Freitag, and Joel Nagloo. Ax-Lindemann-Weierstrass with derivatives and the genus 0 Fuchsian groups. Annals of Mathematics, 192(3):721-765, 2020.
- [6] R. Cohn. A difference-differential basis theorem. Canadian J. Math. 22, No. 6, pp.1224-1237, 1970.
- [7] Z. Chatzidakis and Ehud Hrushovski. Model theory of difference fields. Trans. Amer. Math. Soc., 351(8):2997-3071, 1999.
- [8] Z. Chatzidakis, Generic automorphism of separably closed fields, Illinois Journal of Mathematics Volume 45, Number 3, Fall 2001, Pages 693-733 S 0019-2082.

- [9] Z. Chatzidakis, E. Hrushovski, Y. Peterzil, Model theory of difference fields, II: Periodic ideals and the trichotomy in all characteristics, *Proceedings of the London Math. Society* (3) 85 (2002), 257-311.
- [10] G. Cherlin, S. Shelah. Superstable fields and groups. *Annals of Mathematical Logic*. 18:227–270, 1980.
- [11] F. Delon. Ideaux et types sur les corps séparablement clos. Supplément au Bulletin de la société Mathématique de France. Mémoire No.33, Tome 116, 1988.
- [12] Matthew DeVilbiss and James Freitag. Generic differential equations are strongly minimal. *Compositio Mathematica*. 2023.
- [13] Y. Ershov. Fields with a solvable theory (English translation). *Sov. Math. Doklady*, 8:575–576, 1967.
- [14] James Freitag and Thomas Scanlon. Strong minimality and the j -function. *Journal of the European Mathematical Society*, 20(1):119-136, 2017.
- [15] J. Gogolok. Model theory of derivations of the Frobenius map revisited. *Journal of Symbolic Logic*, DOI:10.1017/jsl.2021.85.
- [16] N. Guzy and C. Riviere. On existentially closed partial differential fields with an automorphism. Preprint 2007.
- [17] C. Hardouin and M. Singer. Differential Galois theory of linear difference equations. *Mathematische Annalen* 342(2), pp.333-377, 2008.
- [18] E. Hrushovski, The Mordell-Lang conjecture for function fields, *J. Amer. Math. Soc.* 9 no. 3 (1996) 667-690.
- [19] E. Hrushovski. The Manin-Mumford conjecture and the model theory of difference fields. *Annals of Pure and Applied Logic*, 112(1), pp.43-115, 2001.
- [20] K. Ino, Model theory of separably differentially closed fields. PhD Thesis, The University of Manchester, 2023.
- [21] K. Ino, O. León Sánchez, Separably differentially closed fields, 2023. Preprint (<http://arxiv.org/abs/2302.11319>).
- [22] W. Johnson, C. Tran, E. Walsberg, and J. Ye, The Étale-open topology and the stable fields conjecture, 2020. (arXiv:2101.07782). To appear in *Journal of the European Mathematical Society*.
- [23] Itay Kaplan, Thomas Scanlon, and Frank O. Wagner, Artin-Schreier extensions in NIP and simple fields, *Israel Journal of Mathematics* 185 (2011), no. 1, 141-153.
- [24] E. R. Kolchin. *Differential algebra and algebraic groups*. Academic Press, 1973.
- [25] E. R. Kolchin. Constrained extensions of differential fields. *Advances in Mathematics*, 12:141–170, 1974.
- [26] P. Kowalski. Derivations of the Frobenius map. *Journal of Symbolic Logic*, 70(1):99-110, 2005.
- [27] O. León Sánchez, Geometric axioms for differentially closed field in several commuting derivations. *Journal of Algebra*, 362:107-116, 2012. (Corrigendum, 382:332-334, 2013.)
- [28] O. León Sánchez, Relative D-groups and differential Galois theory in several derivations. *Transactions AMS*, 367: 7613-7638, 2015.
- [29] O. León Sánchez, On the model companion of partial differential fields with an automorphism. *Israel Journal of Mathematics*, 2016.
- [30] O. León Sánchez, ALGEBRO-GEOMETRIC AXIOM FOR $DCF_{0,m}$. *Fundamenta Mathematicae*, 243, 1, p. 1-8, 2018.
- [31] O. León Sánchez, and M. Tressl. Differentially large fields. To appear in *Algebra and Number Theory*, 2023.
- [32] A. Macintyre, ω_1 -categorical fields, *Fundamenta Mathematicae* 70 (1971) no. 3, 253-270.
- [33] D. Marker, M. Messmer, A. Pillay. *Model theory of fields*. Lecture Notes in Logic 5. Springer, 1996.
- [34] R. Moosa and T. Scanlon. Jet and prolongation spaces. *Journal of the Inst. of Math. Jussieu* 9, pp.391-430, 2010.
- [35] M. Morley. Categoricity in power. *Transactions of the Amer. Math. Soc.*, 114:514–538, 1965.
- [36] J. Nagloo and A. Pillay, On Algebraic relations between solutions of a generic Painlevé equation, *J. Reine Angew. Math. (Crelles Journal)* 726 (2017) 1-27.
- [37] D. Pierce, A. Pillay. A note on the axioms for differentially closed fields of characteristic zero. *Journal of Algebra*, 204:108–115, 1998.
- [38] A. Pillay. Differential Galois theory I. *Illinois Journal of Mathematics*. Vol. 42, N. 4, 1998.

- [39] A. Pillay and M. Ziegler. Jet spaces of varieties over differential and difference fields. *Selecta Mathematica. New series* 9, pp.579-599, 2003.
- [40] A. Robinson, On the concept of differentially closed field, *Bull. Res. Council. Isr. Sect. F* 8 (1959), 113-118.
- [41] M. Rosenlicht, The nonminimality of the differential closure, *Pacific J. Math.* 52 (1974), 529-537.
- [42] A. Seidenberg. Some basic theorems in differential algebra (characteristic p , arbitrary). *Transactions of the Amer. Math. Soc.*, 73(1):174-190, 1952.
- [43] S. Shelah. Uniqueness and characterization of prime models over sets for totally transcendental first-order theories, *J. Symbolic Logic* 37 (1972), 107-113.
- [44] S. Shelah. Differentially closed fields. *Israel Journal of Mathematics*, 16:314-328, 1973.
- [45] K. Tent and M. Ziegler. *A course in model theory. Lecture Notes in Logic. Vol.40, ASL*, 2012.
- [46] C. Wood. The model theory of differential fields revisited. *Israel Journal of Mathematics*. 25:331-352, 1976.
- [47] C. Wood. Notes on the stability of separably closed fields. *Journal of Symbolic Logic*. 44(3):412-416, 1979.

KAI INO, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, OXFORD ROAD, MANCHESTER, UNITED KINGDOM M13 9PL

E-mail address: `kai.ino@manchester.ac.uk`