

On the Stability of Out-flowing Compressible Viscous Gas

Yucong Huang

School of Mathematics, University of Edinburgh

Joint work with Shinya Nishibata

1 Introduction

For dimensions $n \geq 2$, the isentropic compressible flow is governed by the following system of partial differential equations:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho U) = 0, \\ \partial_t(\rho U) + \operatorname{div}(\rho U \otimes U) + \nabla P = \operatorname{div} \mathbb{S}, \end{cases} \quad (\mathbf{CNS})$$

where $\rho \geq 0$ is mass density, $U = (U^1, \dots, U^n)$ is velocity, $P \geq 0$ is pressure given by,

$$P(\rho) = K\rho^\gamma \quad \text{where } K > 0 \text{ and } \gamma \geq 1.$$

$\mathbb{S} = \mathbb{S}(\nabla U)$ is the viscous stress tensor given by

$$\mathbb{S}(\nabla U) := \mu_1(\nabla U + \nabla U^\top) + \mu_2 \mathbb{I}_n \operatorname{div} U,$$

where \mathbb{I}_n is $n \times n$ identity matrix, $\mu_1 > 0$ is the shear viscosity coefficient and quantity $\frac{2}{n}\mu_1 + \mu_2 \geq 0$ is the bulk viscosity coefficient. Suppose fluid occupies the exterior domain of a unit ball, which is given by $\Omega := \{x \in \mathbb{R}^n \mid |x| > 1\}$. Moreover, we consider the spherically symmetric solution:

$$\rho(t, x) = \rho(t, r), \quad U(t, x) = u(t, r) \frac{x}{r}, \quad \text{where } r := |x|.$$

Then the problem (\mathbf{CNS}) can be reformulated in the variables $(t, r) \in (0, \infty) \times [1, \infty)$ as:

$$\begin{cases} \partial_t \rho + \frac{\partial_r(r^{n-1} \rho u)}{r^{n-1}} = 0, \\ \rho(\partial_t u + u \partial_r u) + \partial_r P(\rho) = \mu \partial_r \left(\frac{\partial_r(r^{n-1} u)}{r^{n-1}} \right), \end{cases} \quad (\mathbf{SCNS})$$

where $\mu := 2\mu_1 + \mu_2$. For the outflow problem, we consider the boundary and far-field conditions:

$$u(1, t) = u_b < 0, \quad \lim_{r \rightarrow \infty} (\rho, u)(r, t) = (\rho_+, u_+), \quad (\mathbf{B})$$

where $\rho_+ > 0$ and $u_+ \in \mathbb{R}$ are the constant reference density and velocity.

2 Stationary Solution

The stationary solution to (SCNS)–(B), denoted as $(\tilde{\rho}, \tilde{u})(r)$, solves the time-independent problem:

$$\begin{cases} \partial_r(r^{n-1}\tilde{\rho}\tilde{u}) = 0, \\ \tilde{\rho}\tilde{u}\partial_r\tilde{u} + \partial_r P(\tilde{\rho}) = \mu\partial_r\left(\frac{\partial_r(r^{n-1}\tilde{u})}{r^{n-1}}\right), \\ \tilde{u}(1) = u_b, \quad \lim_{r \rightarrow \infty} (\tilde{\rho}, \tilde{u})(r) = (\rho_+, u_+). \end{cases} \quad (\mathbf{ST})$$

Suppose the solution exists, then integrating the first equation in $s \in [1, r]$, we obtain that

$$\tilde{u}(r) = u_b \frac{\tilde{\rho}(1)}{\tilde{\rho}(r)} r^{1-n} \quad \text{for } r \geq 1.$$

Since $\tilde{\rho}(r) \rightarrow \rho_+ < \infty$ and $r^{1-n} \rightarrow 0$ as $r \rightarrow \infty$, it follows that if $(\tilde{\rho}, \tilde{u})(r)$ is a solution to (ST), then it necessarily satisfies the condition:

$$u_+ = \lim_{r \rightarrow \infty} \tilde{u}(r) = 0.$$

I. Hashimoto and A. Matsumura in [1] obtained the existence and uniqueness of a $\mathcal{C}^2[1, \infty)$ solution to (ST) for both inflow and outflow problems, provided $|u_b| \leq \delta$ for some $\delta > 0$. For small initial perturbation, I. Hashimoto, S. Nishibata, and S. Sugizaki in [2] showed the long time stability of $(\tilde{\rho}, \tilde{u})$ for both inflow and outflow cases. However, it turns out that the small requirement on initial perturbation can be relaxed for outflow case $u_b < 0$, by employing careful a-priori estimates in the Lagrangian coordinate. The main aim of the present article is to give a survey on the long time stability of outflow stationary solution $(\tilde{\rho}, \tilde{u})$ with large initial perturbation.

3 Main Result

Theorem 1 (S. Nishibata and Y.H., To Appear). *Assume $n \geq 2$ and $1 \leq \gamma \leq 2$. Let initial data (ρ_0, u_0) satisfy*

$$\begin{aligned} r^{\frac{n-1}{2}}(\rho_0 - \tilde{\rho}), \quad r^{\frac{n-1}{2}}\partial_r(\rho_0 - \tilde{\rho}), \quad r^{\frac{n-1}{2}}(u_0 - \tilde{u}), \quad r^{\frac{n-1}{2}}\partial_r(u_0 - \tilde{u}) &\in L^2(1, \infty), \\ \rho_0 &\in \mathcal{C}^1[1, \infty) \quad \text{and} \quad u_0 \in \mathcal{C}^2[1, \infty), \end{aligned}$$

and the initial compatibility condition:

$$u_0(1) = u_b \quad \text{and} \quad \left(\rho_0 u_0 \partial_r u_0 + \mu \partial_r \left(\frac{\partial_r(r^{n-1}u_0)}{r^{n-1}} \right) - \partial_r P(\rho_0) \right) \Big|_{r=1} = 0.$$

Then, there exists $\delta_0 = \delta_0(\rho_0, u_0, \rho_+, \mu, \gamma, K, n) > 0$ so that, if $|u_b| \leq \delta_0$, then initial boundary value problem (SCNS)–(B) has a unique classical solution (ρ, u) such that for any $T > 0$:

$$\begin{aligned} r^{\frac{n-1}{2}}(\rho - \tilde{\rho}), \quad r^{\frac{n-1}{2}}\partial_r(\rho - \tilde{\rho}), \quad r^{\frac{n-1}{2}}(u - \tilde{u}), \quad r^{\frac{n-1}{2}}\partial_r(u - \tilde{u}) &\in \mathcal{C}^0([0, T]; L^2(1, \infty)), \\ \rho &\in \mathcal{C}^{1+\sigma, 1+\sigma/2}, \quad u \in \mathcal{C}^{2+\sigma, 1+\sigma/2} \quad \text{for some } \sigma \in (0, 1). \end{aligned}$$

Furthermore, (ρ, u) satisfies the large time convergence:

$$\lim_{t \rightarrow \infty} \sup_{r \in [1, \infty)} |(\rho(r, t) - \tilde{\rho}(r), u(r, t) - \tilde{u}(r))| = 0.$$

It is worth noting that the weighted H^1 Sobolev norm of $(\rho - \tilde{\rho}, u - \tilde{u})$ are not required to be small, but merely finite (and possibly large). Before ending this section, we mention few previous results on the compressible Navier-Stokes equations:

Adhesion boundary problem $u_b = 0$. A. Matsumura and T. Nishida in 1983 [7] considered the heat-conducting flow, posed in a general 3D exterior domain Ω with adhesion boundary:

$$u|_{\partial\Omega} = 0, \quad \partial_n \theta|_{\partial\Omega} = 0 \quad \text{where } \partial_n \text{ denotes the normal derivative.}$$

And the flow is also under the influence of an external force f . They showed that any small perturbation in $H^3(\Omega)$ to the stationary solution $(\tilde{\rho}(x), 0, \tilde{\theta})$ is stable in large time:

$$\lim_{t \rightarrow \infty} \sup_{x \in \Omega} |(\rho, u, \theta) - (\tilde{\rho}, 0, \tilde{\theta})| = 0.$$

T. Nakamura, S. Nishibata, and S. Yanagi in 2004 [11] studied the spherically symmetric isentropic flow with adhesion boundary condition: $u_b = 0$, and potential force f . They showed the large time stability for any spherical large perturbation to the stationary solution $(\tilde{\rho}(r), 0)$, under the H^1 and Hölder norm. Then N. Nakamura and N. Nishibata in 2008 [9] extended this result to the spherically symmetric heat-conducting model.

Inflow and outflow problems. A. Matsumura in 2001 [6] classified the stationary states to the inflow and outflow problems in 1D half-space $x \in [0, \infty)$. Depending on (ρ_b, u_b) and (ρ_+, u_+) , stationary solutions can either be viscosity shock, rarefaction wave, boundary layer solution, or certain superposition of the three.

A. Matsumura and K. Nishihara in 2001 [8] studied the isentropic inflow problem posed in 1D half-space. Suppose the stationary solution $(\tilde{\rho}, \tilde{u})$ is a superposition of boundary layer solution and weak rarefaction wave. Then they showed that any small perturbation to $(\tilde{\rho}, \tilde{u})$ is stable in large time $t \rightarrow \infty$ under the H^1 norm.

S. Kawashima, S. Nishibata, and P. Zhu in 2003 [4] considered the isentropic outflow problem posed in 1D half-space. They obtained the necessary and sufficient condition for the well-posedness of stationary solution. Moreover, they showed its time asymptotic stability under small perturbations.

T. Nakamura and S. Nishibata in 2011 [10] considered the heat-conducting inflow problem in 1D half-space. They have obtained a necessary and sufficient condition for the well-posedness of stationary solutions, and shown that they are stable under small perturbation in the H^1 and Hölder norm.

4 Strategy

The strategy for proving Theorem 1 is to reformulate equations (SCNS) in the Lagrangian coordinates, then obtain a set of a-priori estimates, exclusive to this Lagrangian formulation. For outflow problem, the transformation between Eulerian coordinate (t, r) and Lagrangian coordinate (t, x) is given by:

$$\begin{aligned} t &= t \quad \text{and} \quad r = R(t, x), \quad \text{where } R(t, x) \text{ is the solution to:} \\ x &= B(t) + \int_1^{R(t, x)} \rho(t, r) r^{n-1} dr \quad \text{and} \quad B(t) := -u_b \int_0^t \rho(1, s) ds. \end{aligned}$$

Note that $B(t)$ physically represents the total amount of mass flown out from the boundary $\{r = 1\}$ during the time period $[0, t]$. By Implicit Function theorem, and equations (SCNS), one can verify that

$$\partial_t R(t, x) = u(t, R(t, x)), \quad \partial_x R(t, x) = \frac{R(t, x)^{1-n}}{\rho(t, R(t, x))} \quad \text{for } t \geq 0 \text{ and } x \in [B(t), \infty).$$

This is similar to the standard differential relations of Lagrangian coordinate for adhesion problem. However, in this case the Lagrangian domain is characterised by the free boundary $x = B(t)$. Set $v = 1/\rho$ to be the specific volume. Then the original Eulerian equations (SCNS) is reformulated as

$$\begin{cases} \partial_t v = \partial_x (R^{n-1} u) \\ \partial_t u + R^{n-1} \partial_x p(v) = \mu R^{n-1} \partial_x \left(\frac{\partial_x (R^{n-1} u)}{v} \right) \\ u(t, B(t)) = u_b \end{cases} \quad \begin{array}{l} \text{for } t \geq 0 \text{ and } x \in [B(t), \infty), \\ \\ \text{for } t \geq 0, \end{array} \quad (\mathbf{L})$$

where $p(v) = Kv^{-\gamma}$, and $R = R(t, x)$ is a function of (v, u) given by

$$R(t, x) = \left(1 + n \int_{B(t)}^x v(t, y) dy \right)^{\frac{1}{n}}.$$

Subtracting (L) with the stationary equations (ST), one can derive the relative energy estimate for difference function $(\phi, \psi) := (v - 1/\tilde{\rho}, u - \tilde{u})$ as follows:

Lemma 1 (Relative Energy Estimate). *Suppose $1 \leq \gamma \leq 2$. Then there exists a generic constant $C > 0$ and $\delta = \delta(\rho_+, \mu, \gamma, K) > 0$ such that if $|u_b| \leq \delta$ then for arbitrary $T > 0$*

$$\begin{aligned} & \sup_{t \in [0, T]} \int_{B(t)}^\infty \mathcal{E}(t, x) dx + |u_b| \int_0^T \left| \frac{|\phi|^2}{v} \right|_{x=B(t)} dt \\ & + \mu \int_0^T \int_{B(t)}^\infty \left(\frac{v\psi^2}{r^2} + \frac{r^{2(n-1)} |\partial_x \psi|^2}{v} \right) dx dt \\ & + C^{-1} \int_0^T \int_{B(t)}^\infty \left\{ |u_b|^3 \frac{|\phi|^2}{r^{3n-2}} + |u_b| \frac{|\psi|^2}{r^n} \right\} dx dt \leq \int_0^\infty \mathcal{E}(0, x) dx. \end{aligned}$$

The main advantage for the Lagrangian formulation is the point-wise representation formula for $v(t, x)$, which was originally derived by A. V. Kazhikhov and V. V. Shelukhin in 1977 [5] for one dimensional adhesion problem in bounded interval with heat-conducting flow. Combining this formula for $v(t, x)$ with the relative energy estimate Lemma 1, one can obtain the upper and lower bound on specific volume $C_0^{-1} \leq v(t, x) \leq C_0$ for all $x \in [B(t), \infty)$, where $C_0 > 0$ is a constant that depends only on the initial data. Once the upper and lower bound on $v(t, x)$ is obtained, it can be used to obtain the H^1 -estimates for (ϕ, ψ) . For $\phi(t, x)$, one uses the following coercivity structure, which is exclusive to the Lagrangian formulation:

$$\partial_t \left(\mu \frac{\partial_x \phi}{v} - \frac{\psi}{r^{n-1}} \right) + \frac{\gamma K}{\mu} v^{-\gamma} \left(\mu \frac{\partial_x \phi}{v} - \frac{\psi}{r^{n-1}} \right) = (\text{lower order terms in } \phi \text{ and } \psi).$$

Note that the above is a modification of the argument used by Ya I Kanel' in 1968 [3], who originally derived the coercivity equation for one dimensional Cauchy problem for isentropic flow. The H^1 estimate for ψ can be attained using the dissipation structure of the momentum equation. Finally, applying the Parabolic Schauder estimates, one completes the desired a-priori estimate.

References

- [1] I. Hashimoto and A. Matsumura. Existence of radially symmetric stationary solutions for the compressible Navier-Stokes equation. *Methods Appl. Anal.*, 28(3):299–311, 2021.
- [2] I. Hashimoto, S. Nishibata, and S. Sugizaki. Asymptotic behavior of spherically symmetric solutions to the compressible Navier-Stokes equations towards stationary waves. *To appear*.
- [3] Ya I Kanel'. A model system of equations for the one-dimensional motion of a gas. *Differentsial'nye Uravneniya*, 4(4):721–734, 1968.
- [4] Shuichi Kawashima, Shinya Nishibata, and Peicheng Zhu. Asymptotic stability of the stationary solution to the compressible Navier-Stokes equations in the half space. *Comm. Math. Phys.*, 240(3):483–500, 2003.
- [5] Aleksandr Vasil'evich Kazhikhov and Vladimir V Shelukhin. Unique global solution with respect to time of initial-boundary value problems for one-dimensional equations of a viscous gas. *Prikladnaia Matematika i Mekhanika*, 41:282–291, 1977.
- [6] Akitaka Matsumura. Inflow and outflow problems in the half space for a one-dimensional isentropic model system of compressible viscous gas. *Methods Appl. Anal.*, 8(4):645–666, 2001.
- [7] Akitaka Matsumura and Takaaki Nishida. Initial-boundary value problems for the equations of motion of compressible viscous and heat-conductive fluids. *Comm. Math. Phys.*, 89(4):445–464, 1983.
- [8] Akitaka Matsumura and Kenji Nishihara. Large-time behaviors of solutions to an inflow problem in the half space for a one-dimensional system of compressible viscous gas. *Comm. Math. Phys.*, 222(3):449–474, 2001.
- [9] Tohru Nakamura and Shinya Nishibata. Large-time behavior of spherically symmetric flow of heat-conductive gas in a field of potential forces. *Indiana Univ. Math. J.*, 57(2):1019–1054, 2008.
- [10] Tohru Nakamura and Shinya Nishibata. Stationary wave associated with an inflow problem in the half line for viscous heat-conductive gas. *J. Hyperbolic Differ. Equ.*, 8(4):651–670, 2011.
- [11] Tohru Nakamura, Shinya Nishibata, and Shigenori Yanagi. Large-time behavior of spherically symmetric solutions to an isentropic model of compressible viscous fluid in a field of potential forces. *Math. Models Methods Appl. Sci.*, 14(12):1849–1879, 2004.
- [12] Tohru Nakamura, Shinya Nishibata, and Takeshi Yuge. Convergence rate of solutions toward stationary solutions to the compressible Navier-Stokes equation in a half line. *J. Differential Equations*, 241(1):94–111, 2007.
- [13] Atusi Tani. On the first initial-boundary value problem of compressible viscous fluid motion. *Publications of the Research Institute for Mathematical Sciences*, 13(1):193–253, 1977.

School of Mathematics
The University of Edinburgh
James Clerk Maxwell Building
Peter Guthrie Tait Rd
Edinburgh EH9 3FD, UK
E-mail address: `yucong.huang@ed.ac.uk`