

THE ALGEBRAIC AND ANABELIAN GEOMETRY OF CONFIGURATION SPACES

SHINICHI MOCHIZUKI AND AKIO TAMAGAWA

July 2007

ABSTRACT. In this paper, we study the *pro- Σ fundamental groups of configuration spaces*, where Σ is either the set of all prime numbers or a set consisting of a single prime number. In particular, we show, via two somewhat distinct approaches, that, in many cases, the “*fiber subgroups*” of such fundamental groups arising from the various natural projections of a configuration space to lower-dimensional configuration spaces may be *characterized group-theoretically*.

Contents:

- §0. Notations and Conventions
- §1. Surface Groups
- §2. Configuration Space Groups
- §3. Direct Products of Profinite Groups
- §4. Product-theoretic Quotients
- §5. Divisors and Units on Coverings of Configuration Spaces
- §6. Nearly Abelian Groups
- §7. A Discrete Analogue

Introduction

Let $n \geq 1$ be an integer; X a *hyperbolic curve* of type (g, r) [where $2g - 2 + r > 0$] over an algebraically closed field k of characteristic 0. Denote by

$$X_n \subseteq P_n$$

the *n-th configuration space* associated to X , i.e., the open subscheme of the direct product P_n of n copies of X obtained by removing the various *diagonals* from P_n

2000 Mathematical Subject Classification. Primary 14H30; Secondary 14H10.

[cf. Definition 2.1, (i)]. By omitting the factors corresponding to various subsets of the set of n copies of X , we obtain various *natural projection morphisms*

$$X_n \rightarrow X_m$$

for nonnegative integers $m \leq n$ [cf. Definition 2.1, (ii)]. Next, let $\Sigma_{\mathcal{C}}$ be either the set of all prime numbers or a set consisting of a single prime number. Write \mathcal{C} for the class of all finite groups of order a product of primes $\in \Sigma_{\mathcal{C}}$. Then by considering the *maximal pro- \mathcal{C} quotient of the étale fundamental group*, which we denote by “ $\pi_1^{\mathcal{C}}(-)$ ”, we obtain various *natural surjections*

$$\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(X_m)$$

arising from the natural projection morphisms considered above. We shall refer to the kernel of such a surjection $\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(X_m)$ as a *fiber subgroup* of $\pi_1^{\mathcal{C}}(X_n)$ of *length* $n - m$ and *co-length* m [cf. Definition 2.3, (iii)]. Also, we shall refer to a closed subgroup of $\pi_1^{\mathcal{C}}(X_n)$ that arises as the inverse image of a closed subgroup of $\pi_1^{\mathcal{C}}(P_n)$ via the natural surjection $\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(P_n)$ [induced by the inclusion $X_n \hookrightarrow P_n$] as *product-theoretic* [cf. Definition 2.3, (ii)].

The present paper is concerned with the issue of the *group-theoretic characterization* of these *fiber subgroups*. Our *main results* [cf. Corollaries 4.8, 6.3] may be summarized as follows:

- (i) Suppose that $g \geq 2$. Let $H \subseteq \pi_1^{\mathcal{C}}(X_n)$ be a *product-theoretic open subgroup*. Then the subgroups $H \cap F$ of H — where F ranges over the various *fiber subgroups* of $\pi_1^{\mathcal{C}}(X_n)$ — may be characterized *group-theoretically* [cf. Corollary 4.8].
- (ii) Suppose that (g, r) is not equal to $(0, 3)$ or $(1, 1)$. Then the *fiber subgroups* of $\pi_1^{\mathcal{C}}(X_n)$ may be characterized *group-theoretically* [cf. Corollary 6.3].

The proof of (i) is obtained as a consequence of the following result [cf. Theorem 4.7]:

- (iii) In the notation of (i), every normal closed subgroup $J \subseteq H$ such that the quotient group H/J is *abelian* and *torsion-free* is, in fact, *product-theoretic*.

The proof of (iii) is based on a slightly complicated computation involving *Chern classes* [cf. §4], together with the well-known fact that the action of the Galois group of a finite Galois covering of a curve of genus ≥ 2 on the Tate module of the Jacobian of the covering curve contains the *regular representation* [cf. Proposition 1.3]. On the other hand, the proof of (ii), due to the second author, makes essential use to the notion of a “*nearly abelian group*”, i.e., a profinite group G which admits a normal closed subgroup $N \subseteq G$ which is *topologically normally generated by a single element* $\in G$ such that G/N contains an open abelian subgroup [cf. Definition 6.1]. It is worth noting that at the time of writing, we are unable to prove *either* an analogue of (i) for $g < 2$ *or* an analogue of (ii) when (g, r) is equal to $(0, 3)$ or $(1, 1)$.

The original proof of (i) [due to the first author] given in §4 may be regarded as a consequence of various *explicit group-theoretic manifestations* of certain *algebraic-geometric* properties. This proof of (i) motivated the second author to develop a more direct approach to understanding these essentially *purely algebraic-geometric* properties. This approach, which is exposed in §5, allows one to prove a *stronger version* [cf. Theorem 5.6], in the case of *proper hyperbolic curves*, of Theorem 4.7 and, moreover, implies certain interesting consequences concerning the *non-existence of units on finite étale coverings of a sufficiently generic hyperbolic curve* [cf. Corollary 5.7].

The contents of the present paper may be summarized as follows: Basic well-known facts concerning the profinite fundamental groups of *hyperbolic curves* and *configuration spaces*, including a certain mild generalization of a theorem of *Lubotzky-Melnikov-van den Dries*, are reviewed in §1, §2, respectively. In §3, we discuss the *group-theoreticity* of direct product decompositions of profinite groups. In §4, §6, we present the proofs, via somewhat different techniques, of the main results (i), (ii) discussed above. In §5, we discuss the algebraic geometry of *divisors* and *units* on configuration spaces, a theory which yields an alternate approach to the theory of §4, in the case of *proper hyperbolic curves*. Finally, in §7, we observe that these results (i), (ii) imply a certain *discrete analogue* [cf. Corollary 7.4] of (i), (ii).

Section 0: Notations and Conventions

Numbers:

The notation \mathbb{Q} will be used to denote the field of *rational numbers*. The notation $\mathbb{Z} \subseteq \mathbb{Q}$ will be used to denote the set, group, or ring of *rational integers*. The notation $\mathbb{N} \subseteq \mathbb{Z}$ will be used to denote the set or [additive] monoid of *nonnegative integers*. If l is a prime number, then the notation \mathbb{Q}_l (respectively, \mathbb{Z}_l) will be used to denote the *l -adic completion* of \mathbb{Q} (respectively, \mathbb{Z}). The [topological] field of complex numbers will be denoted \mathbb{C} .

Topological Groups:

Let G be a *Hausdorff topological group*, and $H \subseteq G$ a *closed subgroup*. Let us write

$$Z_G(H) \stackrel{\text{def}}{=} \{g \in G \mid g \cdot h = h \cdot g, \forall h \in H\}$$

for the *centralizer* of H in G . Also, we shall write $Z(G) \stackrel{\text{def}}{=} Z_G(G)$ for the *center* of G .

We shall say that a profinite group G is *slim* if for every open subgroup $H \subseteq G$, the centralizer $Z_G(H)$ is trivial. Note that every *finite normal closed subgroup* $N \subseteq G$ of a slim profinite group G is *trivial*. [Indeed, this follows by observing that for any normal open subgroup $H \subseteq G$ such that $N \cap H = \{1\}$, consideration of the

inclusion $N \hookrightarrow G/H$ reveals that the conjugation action of H on N is *trivial*, i.e., that $N \subseteq Z_G(H) = \{1\}$.]

We shall write G^{ab} for the *abelianization* of G , i.e., the quotient of G by the closure of the commutator subgroup of G . We shall denote the group of automorphisms of G by $\text{Aut}(G)$. Conjugation by elements of G determines a homomorphism $G \rightarrow \text{Aut}(G)$ whose image consists of the *inner automorphisms* of G . We shall denote by $\text{Out}(G)$ the quotient of $\text{Aut}(G)$ by the [normal] subgroup consisting of the inner automorphisms. In particular, if G is *center-free*, then we have an *exact sequence* $1 \rightarrow G \rightarrow \text{Aut}(G) \rightarrow \text{Out}(G) \rightarrow 1$.

Curves:

Suppose that $g \geq 0$ is an *integer*. Then if S is a scheme, a *family of curves of genus g*

$$X \rightarrow S$$

is defined to be a smooth, proper, geometrically connected morphism of schemes $X \rightarrow S$ whose geometric fibers are curves of genus g .

Suppose that $g, r \geq 0$ are *integers* such that $2g - 2 + r > 0$. We shall denote the *moduli stack of r -pointed stable curves of genus g* over \mathbb{Z} (where we assume the points to be *ordered*) by $\overline{\mathcal{M}}_{g,r}$ [cf. [DM], [Knud] for an exposition of the theory of such curves]. The open substack $\mathcal{M}_{g,r} \subseteq \overline{\mathcal{M}}_{g,r}$ of smooth curves will be referred to as the *moduli stack of smooth r -pointed stable curves of genus g* or, alternatively, as the *moduli stack of hyperbolic curves of type (g, r)* . The *divisor at infinity* $\overline{\mathcal{M}}_{g,r} \setminus \mathcal{M}_{g,r}$ of $\overline{\mathcal{M}}_{g,r}$ is a *divisor with normal crossings* on the \mathbb{Z} -smooth algebraic stack $\overline{\mathcal{M}}_{g,r}$, hence determines a *log structure* on $\overline{\mathcal{M}}_{g,r}$; denote the resulting log stack by $\overline{\mathcal{M}}_{g,r}^{\text{log}}$. For any integer $r' > r$, the operation of “forgetting the last $r' - r$ points” determines a [1-]morphism of log algebraic stacks

$$\overline{\mathcal{M}}_{g,r'}^{\text{log}} \rightarrow \overline{\mathcal{M}}_{g,r}^{\text{log}}$$

which factors as a composite of structure morphisms of various tautological log stable curves [cf. [Knud]], hence is *log smooth*.

A *family of hyperbolic curves of type (g, r)*

$$X \rightarrow S$$

is defined to be a morphism which factors $X \hookrightarrow Y \rightarrow S$ as the composite of an open immersion $X \hookrightarrow Y$ onto the complement $Y \setminus D$ of a relative divisor $D \subseteq Y$ which is finite étale over S of relative degree r , and a family $Y \rightarrow S$ of curves of genus g . One checks easily that, if S is *normal*, then the pair (Y, D) is *unique up to canonical isomorphism*. We shall refer to Y (respectively, D) as the *compactification* (respectively, *divisor of cusps*) of X . A *family of hyperbolic curves $X \rightarrow S$* is defined to be a morphism $X \rightarrow S$ such that the restriction of this morphism to each connected component of S is a *family of hyperbolic curves of type (g, r)* for

some integers g, r as above. A family of hyperbolic curve of type $(0, 3)$ will be referred to as a *tripod*.

Section 1: Surface Groups

In the present §1, we discuss various well-known preliminary facts concerning the sorts of profinite groups that arise from étale fundamental groups of *hyperbolic curves*.

Definition 1.1. Let \mathcal{C} be a family of finite groups containing the trivial group; Σ a set of prime numbers.

(i) We shall refer to a finite group as a Σ -*group* if every prime dividing its order belongs to Σ . We shall refer to a finite group belonging to \mathcal{C} as a \mathcal{C} -*group* and to a profinite group every finite quotient of which is a \mathcal{C} -group as a *pro- \mathcal{C} group*. We shall refer to \mathcal{C} as a *full formation* [cf. [FJ], p. 343] if it is closed under taking quotients, subgroups, and extensions.

(ii) Suppose that \mathcal{C} is a full formation; write $\Sigma_{\mathcal{C}}$ for the set of primes p such that $\mathbb{Z}/p\mathbb{Z}$ is a \mathcal{C} -group and $\widehat{\mathbb{Z}} \twoheadrightarrow \widehat{\mathbb{Z}}_{\mathcal{C}}$ for the maximal pro- \mathcal{C} quotient of $\widehat{\mathbb{Z}}$. Then we shall say that the formation \mathcal{C} is *nontrivial* if there exists a nontrivial \mathcal{C} -group [or, equivalently, if $\Sigma_{\mathcal{C}}$ is nonempty]. We shall say that the formation \mathcal{C} is *primary* if $\Sigma_{\mathcal{C}}$ is of cardinality one. We shall say that the formation \mathcal{C} is *solvable* if every \mathcal{C} -group is solvable. We shall say that the formation \mathcal{C} is *total* if every finite group is a \mathcal{C} -group. We shall say that \mathcal{C} is a *PT-formation* if it is *either* primary *or* total. We shall say that \mathcal{C} is *invertible on a scheme S* if every prime of $\Sigma_{\mathcal{C}}$ is invertible on S .

(iii) Suppose that \mathcal{C} is a full formation; let G be a profinite group. If G admits an open subgroup H which is abelian, then we shall say that G is *almost abelian*. If G admits an open subgroup H which is pro- \mathcal{C} , then we shall say that G is *almost pro- \mathcal{C}* . We shall refer to a quotient $G \twoheadrightarrow Q$ as *almost pro- \mathcal{C} -maximal* if for some open subgroup $H \subseteq G$ with maximal pro- \mathcal{C} quotient [cf. [FJ], p. 344] $H \twoheadrightarrow P$, we have $\text{Ker}(G \twoheadrightarrow Q) = \text{Ker}(H \twoheadrightarrow P)$. [Thus, any almost pro- \mathcal{C} -maximal quotient of G is almost pro- \mathcal{C} .] If G is topologically finitely generated, and, moreover, the abelianization J^{ab} of every open subgroup $J \subseteq G$ is torsion-free, then we shall say that G is *strongly torsion-free*.

Remark 1.1.1. The notion of a full formation is a special case of the notion of a *Melnikov formation* [cf. [FJ], p. 343]. In the present paper, [partly for the sake of simplicity] we restrict ourselves to full formations.

Remark 1.1.2. Let \mathcal{C} be a full formation. Then [it follows immediately from the definitions that] a *solvable* finite group is a $\Sigma_{\mathcal{C}}$ -group [cf. Definition 1.1, (ii)]

if and only if it is a \mathcal{C} -group. In particular, if \mathcal{C} is *solvable*, then it is *completely determined* by the set of primes $\Sigma_{\mathcal{C}}$.

Remark 1.1.3. Recall that every finite group whose order is a prime power is *nilpotent*, hence, in particular, *solvable*. Thus, [cf. Remark 1.1.2] a *primary* full formation \mathcal{C} is *completely determined* by the unique prime number $\in \Sigma_{\mathcal{C}}$.

Remark 1.1.4. One verifies immediately that in the various definitions in Definition 1.1, (iii), of terms of the form “*almost* \mathbb{P} ”, where “ \mathbb{P} ” is some property, an *equivalent definition* is obtained if one requires the open subgroup “ H ” to be *normal*.

Remark 1.1.5. If G is a *strongly torsion-free* profinite group, then one verifies immediately [by considering abelianizations of open subgroups of G] that G is *torsion-free* in the usual sense, i.e., that G has no nontrivial elements of finite order.

Definition 1.2. Let \mathcal{C} be a full formation. We shall say that a profinite group is a [*pro- \mathcal{C}*] *surface group* (respectively, an *almost pro- \mathcal{C} -surface group*) if it is isomorphic to the maximal pro- \mathcal{C} quotient (respectively, to some almost pro- \mathcal{C} -maximal quotient) of the étale fundamental group of a hyperbolic curve [cf. §0] over an algebraically closed field of characteristic zero [or, equivalently, the profinite completion of the topological fundamental group of a hyperbolic Riemann surface of finite type]. We shall refer to an almost pro- \mathcal{C} -surface group as *open* (respectively, *closed*) if it admits (respectively, does not admit) a pro- \mathcal{C} free [cf. [FJ], p. 345] open subgroup.

Remark 1.2.1. Thus, in the notation of Definition 1.2, every pro- \mathcal{C} surface group is an almost pro- \mathcal{C} -surface group. On the other hand, if \mathcal{C} is *not total*, then one verifies immediately that there exist almost pro- \mathcal{C} -surface groups which are not pro- \mathcal{C} surface groups. Nevertheless, every almost pro- \mathcal{C} -surface group admits a *normal open subgroup* which is a *pro- \mathcal{C} surface group*.

Remark 1.2.2. We recall that if Π is a *pro- \mathcal{C} surface group* arising from a hyperbolic curve [cf. Definition 1.2] of type (g, r) , then Π is *topologically generated* by $2g + r$ generators *subject to a single [well-known!] relation*, and Π^{ab} [cf. §0] is a *free abelian pro- \mathcal{C} group* of rank $2g - 1 + r$ (if $r > 0$), $2g$ (if $r = 0$). In particular, [since every open subgroup of Π is again a pro- \mathcal{C} surface group, it follows that] Π is *strongly torsion-free*. Moreover, for any $l \in \Sigma_{\mathcal{C}}$, the *l -cohomological dimension* of Π is equal to 1 (if $r > 0$), 2 (if $r = 0$); $\dim_{\mathbb{Q}_l}(H^2(\Pi, \mathbb{Q}_l)) = \dim_{\mathbb{F}_l}(H^2(\Pi, \mathbb{F}_l))$ is equal to 0 (if $r > 0$), 1 (if $r = 0$). In particular, the quantity

$$\chi(\Pi) = \sum_{i=0}^2 (-1)^i \cdot \dim_{\mathbb{Q}_l}(H^i(\Pi, \mathbb{Q}_l)) = \sum_{i=0}^2 (-1)^i \cdot \dim_{\mathbb{F}_l}(H^i(\Pi, \mathbb{F}_l)) = 2 - 2g - r$$

is a *group-theoretic invariant* of Π which [as is well-known] satisfies the property that

$$\chi(\Pi_1) = [\Pi : \Pi_1] \cdot \chi(\Pi)$$

for any open subgroup $\Pi_1 \subseteq \Pi$. Finally, we recall that this formula admits a *representation-theoretic generalization*, which will play a crucial role in §4 below, in the form of the following elementary consequence:

Proposition 1.3. (Inclusion of the Regular Representation) *Let $Y \rightarrow X$ be a finite [possibly ramified] Galois covering of smooth proper hyperbolic curves over an algebraically closed field k of characteristic prime to the order of $G \stackrel{\text{def}}{=} \text{Gal}(Y/X)$; l a prime number that is invertible in k . Write V for the G -module determined by the first étale cohomology module $H_{\text{ét}}^1(Y, \mathbb{Q}_l)$. Then the G -module V contains the **regular representation** of G as a direct summand.*

Proof. Indeed, this follows immediately from the computation of the Galois module V in [Milne], p. 187, Corollary 2.8 [cf. also [Milne], p. 187, Remark 2.9], in light of our assumption that X is proper hyperbolic, hence of *genus* ≥ 2 . \circ

Proposition 1.4. (Slimness) *Let \mathcal{C} be a nontrivial full formation. Then every almost pro- \mathcal{C} -surface group Π is slim.*

Proof. Indeed, this follows immediately by considering the *conjugation action* of Π/N on $N^{\text{ab}} \otimes \mathbb{Z}_l$, where $l \in \Sigma_{\mathcal{C}}$, for sufficiently small normal open subgroups $N \subseteq \Pi$ [cf. Remark 1.2.1]. That is to say, in light of the interpretation of a certain quotient of $N^{\text{ab}} \otimes \mathbb{Z}_l$ as the Tate module arising from the l -power torsion points of the Jacobian of the compactification of the covering determined by N of any hyperbolic curve that gives rise to Π [cf. the proof of [Mzk3], Lemma 1.3.1], it follows that this conjugation action is *faithful*. Another [earlier] approach to the *slimness* of surface groups may be found in [Naka], Corollary 1.3.4. \circ

Remark 1.4.1. The property involving the *regular representation* discussed in Proposition 1.3 may be regarded as a *stronger version* [in the case of coverings of curves of genus ≥ 2] of the *faithfulness* of the action of Π/N on [a certain quotient of] $N^{\text{ab}} \otimes \mathbb{Z}_l$ that was applied in the proof of Proposition 1.4, hence, in particular, as a stronger version of the *slimness* of surface groups.

Next, we give a mild generalization to arbitrary surface groups of a well-known result for free pro- \mathcal{C} groups due to *Lubotzky-Melnikov-van den Dries*. In particular, the argument given below in the proof of Theorem 1.5 may be regarded as a *short elementary proof* of [a certain portion of] the *theorem of Lubotzky-Melnikov-van den Dries*, as exposed in [FJ], Proposition 24.10.3; [FJ], Proposition 24.10.4, (a).

Theorem 1.5. (Normal Closed Subgroups of Surface Groups) *Let \mathcal{C} be a full formation; Π an almost pro- \mathcal{C} -surface group; $N \subseteq \Pi$ a topologically*

finitely generated normal closed subgroup. Then N is either **trivial** or of **finite index**.

Proof. First, we observe that we may assume without loss of generality that \mathcal{C} is *nontrivial*. Since Π is *slim*, hence does not contain any *nontrivial finite normal closed subgroups* [cf. §0], it follows that we may always replace Π by an open subgroup of Π . In particular, [cf. Remark 1.2.1] we may assume, without loss of generality, that Π is a *pro- \mathcal{C} surface group*. Now suppose that N is *nontrivial* and of *infinite index*. Then there exists an $l \in \Sigma_{\mathcal{C}}$ such that N contains a nontrivial subgroup $A \subseteq N$ which is a quotient of \mathbb{Z}_l . In particular, there exists a normal open subgroup $\Pi_1 \subseteq \Pi$ such that the image of A in Π/Π_1 is nontrivial. Now set $\Pi_A \stackrel{\text{def}}{=} \Pi_1 \cdot A \subseteq \Pi$, $N_A \stackrel{\text{def}}{=} N \cap \Pi_A$ [so Π_A, N_A are open subgroups of Π , N , respectively]. Then N_A is a *topologically finitely generated normal closed subgroup of infinite index* of Π_A such that $A \subseteq N_A$ *surjects* onto the [nontrivial, abelian!] image of Π_A in Π/Π_1 . In particular, by replacing $N \subseteq \Pi$ by $N_A \subseteq \Pi_A$, we may assume without loss of generality that the image of N in Π^{ab} is *nontrivial*.

Since Π is *topologically finitely generated*, there exists a descending sequence of normal open subgroups

$$\dots \subseteq H_n \subseteq \dots \subseteq \Pi$$

[where n ranges over the positive integers] of Π which is, moreover, *exhaustive*, i.e., $\bigcap_n H_n = \{1\}$. Thus, if we set $N_n \stackrel{\text{def}}{=} H_n \cdot N$ [for $n \geq 1$], then [one verifies immediately that] we obtain a descending sequence of normal open subgroups

$$\dots \subseteq N_n \subseteq \dots \subseteq \Pi$$

[where n ranges over the positive integers] of Π such that $\bigcap_n N_n = N$ [cf. the fact that N is *closed*!]. Since N is of *infinite index* in Π , it follows that $[\Pi : N_n] \rightarrow \infty$ as $n \rightarrow \infty$, hence [cf. Remark 1.2.2] that $|\chi(N_n)| \rightarrow \infty$ as $n \rightarrow \infty$. In particular, there exists an n such that the rank [as a free abelian pro- \mathcal{C} group] of N_n^{ab} is $\geq s + 2$, where we write s for any positive integer such that there exist s elements of N that topologically generate N . Since, moreover, the image of N in Π^{ab} , hence *a fortiori* in N_n^{ab} is *nontrivial*, it follows that there exists, for some $l \in \Sigma_{\mathcal{C}}$, a nontrivial homomorphism $\mathbb{Z}_l \rightarrow N_n^{\text{ab}}$ that factors through N . Now write

$$N_n \twoheadrightarrow \Pi^*$$

for the *maximal pro- l quotient* of N_n [so Π^* is a *pro- l surface group*], $N^* \subseteq \Pi^*$ for the image of N in Π^* . Thus, $N^* \subseteq \Pi^*$ is a *topologically finitely generated normal closed subgroup* whose image in [the free \mathbb{Z}_l -module of finite rank] $(\Pi^*)^{\text{ab}}$ is a *nontrivial* \mathbb{Z}_l -submodule $M \subseteq (\Pi^*)^{\text{ab}}$ whose rank is $\leq s$, hence \leq the rank of $(\Pi^*)^{\text{ab}}$ minus 2. In particular, there exists an element $x \in \Pi^*$ such that if we denote by $F^* \subseteq \Pi^*$ the [necessarily *topologically finitely generated*!] closed subgroup topologically generated by N^* and x , then we obtain inclusions of closed subgroups

$$N^* \subseteq F^* \subseteq \Pi^*$$

such that N^* is of infinite index in F^* , and F^* is of infinite index in Π^* [as may be seen by considering the ranks of the images of these subgroups in Π^{ab}].

Now observe that for any two open subgroups $J_2 \subseteq J_1 \subseteq \Pi^*$, the induced morphism $H^2(J_1, \mathbb{Z}_l) \rightarrow H^2(J_2, \mathbb{Z}_l)$ maps a generator of $H^2(J_1, \mathbb{Z}_l)$ to $[J_1 : J_2]$ times a generator $H^2(J_2, \mathbb{Z}_l)$ [where we recall that $H^2(J_1, \mathbb{Z}_l), H^2(J_2, \mathbb{Z}_l)$ are either both zero or both isomorphic to \mathbb{Z}_l , depending on whether Π is an open or a closed surface group]. [Indeed, this follows immediately by thinking about degrees of coverings of proper hyperbolic curves! We refer to Remark 4.1.1; Lemma 4.2, (i) [and its proof], below, for more details on this well-known circle of ideas.] In particular, since F^* is a subgroup of infinite index in Π^* , it follows immediately [by considering open subgroups $J \subseteq \Pi^*$ containing F^*] that F^* is a *pro- l group* whose *[l -]cohomological dimension is ≤ 1* . Thus, by [RZ], Theorem 7.7.4, F^* is a *topologically finitely generated free pro- l group* [i.e., in particular, a pro- l open surface group], and $N^* \subseteq F^*$ is a *nontrivial topologically finitely generated closed normal subgroup of infinite index*. Put another way, it suffices to prove Theorem 1.5 in the case where Π is a *pro- l open surface group*.

Thus, we return to the original notation of the statement of Theorem 1.5, under the further assumption that Π is a *pro- l open surface group* [and N is *nontrivial of infinite index*]. Since N is of infinite index in Π , by replacing Π by an open subgroup containing N [cf. the argument above involving the “ H_n ” and “ N_n ”], we may assume that *the rank of Π is $>$ the rank of N* . Thus, [cf., e.g., [RZ], the proof of Theorem 7.7.4] $\dim_{\mathbb{F}_l}(\Pi^{\text{ab}} \otimes \mathbb{F}_l) > \dim_{\mathbb{F}_l}(N^{\text{ab}} \otimes \mathbb{F}_l)$. Next, observe that there exists a normal open subgroup $H \subseteq \Pi$ such that the natural surjection $N \twoheadrightarrow N^{\text{ab}} \otimes \mathbb{F}_l$ factors through $N/(N \cap H) \xrightarrow{\sim} (N \cdot H)/H$. But this implies that the inclusion $N \hookrightarrow N \cdot H$ induces a homomorphism $N^{\text{ab}} \otimes \mathbb{F}_l \rightarrow (N \cdot H)^{\text{ab}} \otimes \mathbb{F}_l$ that admits a *splitting* $(N \cdot H)^{\text{ab}} \otimes \mathbb{F}_l \rightarrow ((N \cdot H)/H)^{\text{ab}} \otimes \mathbb{F}_l \xrightarrow{\sim} (N/(N \cap H))^{\text{ab}} \otimes \mathbb{F}_l \xrightarrow{\sim} N^{\text{ab}} \otimes \mathbb{F}_l$. In particular, by replacing Π by the open subgroup $N \cdot H$ [where we note that $\dim_{\mathbb{F}_l}((N \cdot H)^{\text{ab}} \otimes \mathbb{F}_l) \geq \dim_{\mathbb{F}_l}(\Pi^{\text{ab}} \otimes \mathbb{F}_l)$], we may assume, without loss of generality, that the natural homomorphism $N^{\text{ab}} \otimes \mathbb{F}_l \rightarrow \Pi^{\text{ab}} \otimes \mathbb{F}_l$ is *injective*, but *not surjective*. Thus, it follows [cf., e.g., [RZ], the proof of Theorem 7.7.4] that there exists a collection of free generators $\{\gamma_i\}_{i \in I}$ [where I is a finite set] of Π such that for some nonempty *proper* subset $I' \subseteq I$, the $\{\gamma_i\}_{i \in I'}$ form a collection of free generators of N . But, as is well-known [and easily verified, by applying the universal property of free pro- l groups, together with the *existence of non-normal cyclic subgroups of finite l -groups!*], this contradicts the *normality* of N . \circ

Section 2: Configuration Space Groups

In the present §2, we discuss various well-known preliminary facts concerning the sorts of profinite groups that arise from étale fundamental groups of *configuration spaces* associated to hyperbolic curves.

First, let us suppose that we have been given a log scheme

$$Z^{\log}$$

which is *log regular* [cf., [Kato2], Definition 2.1]; write $U_Z \subseteq Z$ for the *interior* of Z^{\log} [i.e., the open subscheme on which the log structure of Z^{\log} is *trivial*]. By abuse of notation, we shall often use the notation for a scheme to denote the log scheme with trivial log structure determined by the scheme. If \mathcal{C} is a *full formation* that is *invertible* on Z , and Z is *connected*, then we shall write

$$\pi_1^{\mathcal{C}}(Z^{\log})$$

for the *maximal pro- \mathcal{C} quotient* of the *étale fundamental group* [obtained by considering *Kummer log étale coverings*, for some choice of basepoint — cf. [Ill] for more details] of Z^{\log} . Thus, by the *log purity theorem* of Fujiwara-Kato [cf. [Ill]; [Mzk1], Theorem B], the natural morphism $U_Z \rightarrow Z^{\log}$ induces a [continuous outer] *isomorphism* $\pi_1^{\mathcal{C}}(U_Z) \xrightarrow{\sim} \pi_1^{\mathcal{C}}(Z^{\log})$.

Next, suppose that S is a *regular scheme*, and that

$$X \rightarrow S$$

is a *family of hyperbolic curves of type (g, r)* over S , with *compactification* $X \hookrightarrow Y \rightarrow S$ and *divisor of cusps* $D \subseteq Y$ [cf. §0]. For simplicity, we assume that the finite étale covering $D \rightarrow S$ is *split*. Let $n \in \mathbb{N}$.

Definition 2.1.

(i) For positive integers $i, j \leq n$ such that $i < j$, write

$$\pi_{i,j} : P_n \stackrel{\text{def}}{=} X \times_S \dots \times_S X \rightarrow X \times_S X$$

for the projection of the product P_n of n copies of $X \rightarrow S$ to the i -th and j -th factors. Write E for the set [of cardinality n] of factors of P_n . Then we shall refer to as the *n -th configuration space associated to $X \rightarrow S$* the S -scheme

$$X_n \rightarrow S$$

which is the open subscheme determined by the complement in P_n of the union of the various inverse images via the $\pi_{i,j}$ [as (i, j) ranges over the pairs of positive integers $\leq n$ such that $i < j$] of the image of the diagonal embedding $X \hookrightarrow X \times_S X$. We shall refer to as the *n -th log configuration space associated to $X \rightarrow S$* the [log smooth] log scheme over S

$$Z_n^{\log} \rightarrow S$$

obtained by pulling back the [log smooth] [1-]morphism $\overline{\mathcal{M}}_{g,r+n}^{\log} \rightarrow \overline{\mathcal{M}}_{g,r}^{\log}$ given by “forgetting the last n points” [cf. §0] via the classifying [1-]morphism $S \rightarrow \overline{\mathcal{M}}_{g,r}^{\log}$ determined [up to a permutation of the r remaining points] by $X \rightarrow S$. We shall refer to E as the *index set* of the configuration space X_n , or, alternatively, of the log configuration space Z_n^{\log} .

(ii) In the notation of (i), let $E' \subseteq E$ be a subset of cardinality n' ; $E'' \stackrel{\text{def}}{=} E \setminus E'$; $n'' \stackrel{\text{def}}{=} n - n'$. Then by “*forgetting*” the factors of E that belong to E' , we obtain a *natural projection morphism*

$$p_{E'} = p^{E''} : X_n \rightarrow X_{n''}$$

[and similarly in the logarithmic case], which we shall refer to as the *projection morphism of profile E'* , or, alternatively, the *projection morphism of co-profile E''* . Also, in this situation, we shall refer to n' (respectively, n'') as the *length* (respectively, *co-length*) of this projection morphism.

Remark 2.1.1. One verifies immediately that in the notation of Definition 2.1, (i), X_n may be naturally identified with the *interior* of Z_n^{log} .

Remark 2.1.2. One verifies immediately that in the notation of Definition 2.1, (ii), each projection morphism $p_{E'} = p^{E''} : X_n \rightarrow X_{n''}$ is itself the n' -th *configuration space* associated to a *family of hyperbolic curves of type $(g, r + n'')$* over $X_{n''}$ that embeds as a dense open subscheme of the pull-back via $X_{n''} \rightarrow S$ of the original family of hyperbolic curves $X \rightarrow S$.

Proposition 2.2. (Fundamental Groups of Configuration Spaces) *In the notation of the above discussion, suppose further that the following conditions hold:*

- (a) S is **connected**;
- (b) \mathcal{C} is a **PT-formation** which is **invertible** on S ;
- (c) for each $l \in \Sigma_{\mathcal{C}}$, the **images** of the **cyclotomic character** $\pi_1(S) \rightarrow \mathbb{F}_l^\times$ and the **natural Galois action**

$$\pi_1(S) \rightarrow \text{Aut}(\pi_1(Y_{\bar{s}})^{\text{ab}} \otimes \mathbb{F}_l)$$

*arising from the family of curves $Y \rightarrow S$ are **\mathcal{C} -groups** [a condition which is vacuous if \mathcal{C} is **total**].*

Let $n \geq 1$ be an integer, \bar{s} a **geometric point** of S , and \bar{x} a **geometric point** of X_{n-1} [where we write $X_0 \stackrel{\text{def}}{=} S$]; we shall denote the fibers over geometric points by means of subscripts. Then:

(i) Any projection morphism $X_n \rightarrow X_{n-1}$ of length one determines a **natural exact sequence**

$$1 \rightarrow \pi_1^{\mathcal{C}}((X_n)_{\bar{x}}) \rightarrow \pi_1^{\mathcal{C}}(X_n) \rightarrow \pi_1^{\mathcal{C}}(X_{n-1}) \rightarrow 1$$

of **pro- \mathcal{C} groups**.

(ii) The profinite group $\pi_1^{\mathcal{C}}((X_n)_{\bar{s}})$ is **slim and topologically finitely generated**.

(iii) The natural sequence

$$1 \rightarrow \pi_1^{\mathcal{C}}((X_n)_{\bar{s}}) \rightarrow \pi_1^{\mathcal{C}}(X_n) \rightarrow \pi_1^{\mathcal{C}}(S) \rightarrow 1$$

is **exact**.

(iv) Suppose that k, k' are **separably closed fields**; $k \subseteq k'$; $S = \text{Spec}(k)$; $S' \stackrel{\text{def}}{=} \text{Spec}(k')$; \bar{s} (respectively, \bar{s}') is the geometric point of S (respectively, S') determined by the identity morphism of S (respectively, S'). Then the natural morphism $\pi_1^{\mathcal{C}}((X_n \times_S S')_{\bar{s}'}) \rightarrow \pi_1^{\mathcal{C}}((X_n)_{\bar{s}})$ is an **isomorphism**.

(v) Suppose that $S = \text{Spec}(R)$, where R is a **complete discrete valuation ring**; that \bar{s} arises from an algebraic closure of the residue field of R ; and that $\bar{\eta}$ is a geometric point of S that arises from an algebraic closure of the quotient field K of R . Then the operation of specialization of the normalization of X_n in a covering of $X_n \times_R K$ determines an **isomorphism** $\pi_1^{\mathcal{C}}((X_n)_{\bar{\eta}}) \xrightarrow{\sim} \pi_1^{\mathcal{C}}((X_n)_{\bar{s}})$.

Proof. First, let us observe that since the kernel of the natural surjection $\pi_1^{\mathcal{C}}(X_{\bar{s}}) \rightarrow \pi_1^{\mathcal{C}}(Y_{\bar{s}})$ is topologically normally generated by the inertia groups of the cusps [which are isomorphic to $\widehat{\mathbb{Z}}_{\mathcal{C}}(1)$, where the “(1)” denotes a “Tate twist”, and “ $\widehat{\mathbb{Z}}_{\mathcal{C}}$ ” is as in Definition 1.1, (ii)], condition (c) [together with our assumption that the divisor of cusps of $X \rightarrow S$ is *split*] implies that for each $l \in \Sigma_{\mathcal{C}}$, the *image* of the natural Galois action

$$\pi_1(S) \rightarrow \text{Aut}(\pi_1(X_{\bar{s}})^{\text{ab}} \otimes \mathbb{F}_l)$$

arising from the family of hyperbolic curves $X \rightarrow S$ is a \mathcal{C} -group.

Now we *claim* that to complete the proof of Proposition 2.2, it suffices to verify assertion (v). Indeed, let us assume that assertion (v) *holds* and reason by induction on $n \geq 1$. [That is to say, if $n \geq 2$, then we assume that assertions (i), (ii), and (iii) have already been verified for “ $n - 1$ ”.] Then let us first observe that assertion (iv) follows from assertion (v) by a standard argument in elementary algebraic geometry [cf., e.g., [Mzk4], Proposition 2.3, (ii), in the case where k is of characteristic zero, $n = 1$; since \mathcal{C} is *invertible* on U_Z , and we are free to apply assertion (v), the case of positive characteristic k and arbitrary n is entirely similar]. Next, observe that [in light of Remark 2.1.2; the easily verified fact that the family $X_n \rightarrow X_{n-1}$ also satisfies conditions (a), (b), (c)] assertion (i) is a special case of assertion (iii) for “ $n = 1$ ”; thus, [by applying the induction hypothesis] we may assume that assertion (i) holds if $n \geq 2$. Since, moreover, the property of being a slim topologically finitely generated profinite group holds for a profinite group which is an extension of a profinite group G_1 by a profinite group G_2 whenever it holds for G_1 and G_2 , assertion (ii) [for “ n ”] follows immediately, by applying the induction hypothesis, from assertion (i) (when $n \geq 2$) and Proposition 1.4. As for assertion (iii), let us first observe that by assertions (iv), (v) [and various standard arguments in elementary algebraic geometry], we may assume without

loss of generality that \bar{s} arises from an algebraic closure of the function field K of S . Thus, by considering the natural action of $G_K \stackrel{\text{def}}{=} \pi_1(\text{Spec}(K), \bar{s})$ on \bar{s} , we obtain a *natural outer action*

$$G_K \rightarrow \text{Out}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}))$$

which is *compatible* with the natural outer action of G_K on $\pi_1^{\mathcal{C}}((P_n)_{\bar{s}})$ [which may be identified with the product of n copies of $\pi_1^{\mathcal{C}}(X_{\bar{s}})$], relative to the natural inclusion $X_n \hookrightarrow P_n$ [cf. Definition 2.1, (i)]. In particular, since [by *Zariski-Nagata purity* — i.e., the classical non-logarithmic version of the “log purity theorem” quoted above] the kernel of the natural surjection $\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}) \twoheadrightarrow \pi_1^{\mathcal{C}}((P_n)_{\bar{s}})$ is topologically normally generated by the *inertia groups* of the divisors of $(P_n)_{\bar{s}}$ lying in the complement $(P_n \setminus X_n)_{\bar{s}}$ [which are isomorphic to quotients of $\widehat{\mathbb{Z}}_{\mathcal{C}}(1)$], condition (c) [together with the observation at the beginning of the present proof] implies that for each $l \in \Sigma_{\mathcal{C}}$, the *image* of the natural Galois action

$$G_K \rightarrow \text{Aut}(\pi_1((X_n)_{\bar{s}})^{\text{ab}} \otimes \mathbb{F}_l)$$

is a \mathcal{C} -group, hence [cf. Remark 1.1.3 when \mathcal{C} is *primary*] that the homomorphism $G_K \rightarrow \text{Out}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}))$ *factors* through the *maximal pro- \mathcal{C} quotient* $G_K^{\mathcal{C}}$ of G_K . Note, moreover, that [again] by *Zariski-Nagata purity* [i.e., the classical non-logarithmic version of the “log purity theorem” quoted above], the kernel of the natural surjection $G_K^{\mathcal{C}} \twoheadrightarrow \pi_1^{\mathcal{C}}(S)$ is topologically normally generated by the various *inertia groups* determined by the prime divisors of S . On the other hand, by assertion (v), the images of these inertia groups in $\text{Out}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}))$ are *trivial*. Thus, we obtain a homomorphism $\pi_1^{\mathcal{C}}(S) \rightarrow \text{Out}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}))$, hence — by pulling back the natural exact sequence

$$1 \rightarrow \pi_1^{\mathcal{C}}((X_n)_{\bar{s}}) \rightarrow \text{Aut}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}})) \rightarrow \text{Out}(\pi_1^{\mathcal{C}}((X_n)_{\bar{s}})) \rightarrow 1$$

[cf. assertion (ii); §0] via this homomorphism — an exact sequence as in assertion (iii). This completes the proof of the *claim*.

Finally, we consider assertion (v). First, we remark that assertion (v) is a special case of the more general result of [Vid], Théorème 2.2; since, however, [Vid] has yet to be published at the time of writing, we give a self-contained [modulo published results] proof of assertion (v), as follows. We begin by observing that by the *log purity theorem*, we have natural isomorphisms

$$\pi_1^{\mathcal{C}}((X_n)_{\bar{s}}) \xrightarrow{\sim} \pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{s}}); \quad \pi_1^{\mathcal{C}}((X_n)_{\bar{\eta}}) \xrightarrow{\sim} \pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{\eta}})$$

[cf. Definition 2.1, (i); Remark 2.1.1]. Now suppose that $W_0^{\log} \rightarrow (Z_n^{\log})_{\bar{s}}$ is a connected *Kummer log étale covering*. Since $(Z_n^{\log})_{\bar{s}}$ is *log regular*, it thus follows that W_0^{\log} is also log regular, hence, in particular, *normal*. By the definition of “log étale”, one may deform this covering to a *formal Kummer log étale covering* over the \mathfrak{m}_R -completion [where \mathfrak{m}_R is the maximal ideal of R] of Z_n^{\log} . Moreover, the underlying scheme of this formal covering may be *algebrized* [cf. [EGA III], Théorème 5.4.5; the easily verified fact that Z_n is *projective*], hence determines a

finite morphism $W \rightarrow Z_n$. Now it follows from the well-known local structure of Kummer log étale coverings that the formal covering that gave rise to W is S -flat, hence that W itself is S -flat, with normal special fiber $W_{\bar{s}} \cong W_0$. Since S is, of course, normal, we thus conclude [cf. [EGA IV], Corollaire 6.5.4, (ii)] that W is normal and connected, hence irreducible. By considering the formal covering that gave rise to W at completions of closed points of Z_n lying in the interior $X_n \subseteq Z_n$, it follows, moreover, that $W \rightarrow Z_n$ is generically étale. Thus, it makes sense to speak of the ramification divisor in Z_n of $W \rightarrow Z_n$. On the other hand, again by considering the formal covering that gave rise to W , it follows immediately that this ramification divisor is contained in the complement of X_n in Z_n , hence [by the log purity theorem!] that $W \rightarrow Z_n$ determines a Kummer log étale covering $W^{\log} \rightarrow Z_n^{\log}$ whose special fiber $W_{\bar{s}}^{\log} \rightarrow (Z_n^{\log})_{\bar{s}}$ may be naturally identified with the given covering $W_0^{\log} \rightarrow (Z_n^{\log})_{\bar{s}}$. Thus, by algebrizing morphisms between formal Kummer log étale coverings [cf. [EGA III], Théorème 5.4.1], we conclude that the deformation and algebrization procedure just described determines an equivalence of categories between the categories of Kummer log étale coverings of $(Z_n^{\log})_{\bar{s}}$, Z_n^{\log} . In particular, we obtain a natural isomorphism $\pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{s}}) \xrightarrow{\sim} \pi_1^{\mathcal{C}}(Z_n^{\log})$.

On the other hand, again by the log purity theorem, it follows immediately that we obtain an isomorphism

$$\pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{\eta}}) \xrightarrow{\sim} \varinjlim_{S'} \pi_1^{\mathcal{C}}(Z_n^{\log} \times_S S')$$

[where S' ranges over the normalizations of S in the various finite extensions of K in the function field of $\bar{\eta}$], hence, by applying the isomorphisms

$$\pi_1^{\mathcal{C}}(Z_n^{\log} \times_S S') \xrightarrow{\sim} \pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{s}})$$

[where we regard \bar{s} as a geometric point of the various S'] obtained above, we obtain an isomorphism $\pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{\eta}}) \xrightarrow{\sim} \pi_1^{\mathcal{C}}((Z_n^{\log})_{\bar{s}})$, as desired. \circ

Remark 2.2.1. Another proof of Proposition 2.2, (iii), in the case $n = 1$ may be found in [Stix], Proposition 2.3.

Definition 2.3. Let \mathcal{C} be a PT-formation.

(i) We shall say that a profinite group is a $[\text{pro-}\mathcal{C}]$ configuration space group if it is isomorphic to the maximal pro- \mathcal{C} quotient of the étale fundamental group

$$\pi_1^{\mathcal{C}}(X_n)$$

of the n -th configuration space X_n for some $n \geq 1$ [cf. Definition 2.1, (i)] of a hyperbolic curve X over an algebraically closed field of characteristic $\notin \Sigma_{\mathcal{C}}$ [where we note that in this situation, if we take S to be the spectrum of this algebraically closed field, then the conditions (a), (b), (c) of Proposition 2.2 are satisfied].

(ii) Let X be a hyperbolic curve over an algebraically closed field of characteristic $\notin \Sigma_{\mathcal{C}}$; X_n the n -th configuration space [for some $n \geq 1$] associated to X .

Then we shall refer to a closed subgroup $H \subseteq \pi_1^{\mathcal{C}}(X_n)$ as being *product-theoretic* if H arises as the inverse image via the natural surjection

$$\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(P_n)$$

[cf. Definition 2.1, (i)] of a closed subgroup of $\pi_1^{\mathcal{C}}(P_n)$.

(iii) Let X, X_n be as in (ii); write E for the *index set* of X_n . Let $E' \subseteq E$ be a subset of cardinality n' ; $E'' \stackrel{\text{def}}{=} E \setminus E'$; $n'' \stackrel{\text{def}}{=} n - n'$; $p_{E'} = p^{E''} : X_n \rightarrow X_{n''}$ the projection morphism of profile E' . Then we shall refer to the *kernel*

$$F \subseteq \pi_1^{\mathcal{C}}(X_n)$$

of the induced *surjection* $\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(X_{n''})$ [cf. Remark 2.1.2; Proposition 2.2, (iii)] as the *fiber subgroup* of $\pi_1^{\mathcal{C}}(X_n)$ of *profile* E' , or, alternatively, as the *fiber subgroup* of $\pi_1^{\mathcal{C}}(X_n)$ of *co-profile* E'' . Also, we shall refer to n' (respectively, n'') as the *length* (respectively, *co-length*) of F .

Proposition 2.4. (Fiber Subgroups of Configuration Spaces) *Let \mathcal{C} be a PT-formation; X a hyperbolic curve over an algebraically closed field of characteristic $\notin \Sigma_{\mathcal{C}}$; X_n the n -th configuration space [for some $n \geq 1$] associated to X ; E the index set of X_n ; $\Pi \stackrel{\text{def}}{=} \pi_1^{\mathcal{C}}(X_n)$; $E'_1, E'_2 \subseteq E$ subsets whose respective complements we denote by $E''_1, E''_2 \subseteq E$; $F_1, F_2 \subseteq \Pi$ the fiber subgroups with respective profiles $E'_1, E'_2 \subseteq E$. Then:*

(i) *The description of Remark 2.1.2 determines on F_2 (respectively, Π/F_2) a structure of configuration space group with index set E'_2 (respectively, E''_2).*

(ii) *$F_1 \subseteq F_2$ if and only if $E'_1 \subseteq E'_2$. Moreover, in this situation, $F_1 \subseteq F_2$ is the fiber subgroup of F_2 with profile $E'_1 \subseteq E'_2$ [i.e., relative to the structure of F_2 as the “ $\pi_1^{\mathcal{C}}(-)$ ” of a configuration space that arises from the description given in Remark 2.1.2].*

(iii) *The image of F_1 in Π/F_2 is the fiber subgroup of Π/F_2 with profile $E'_1 \cap E''_2 \subseteq E''_2$ [i.e., relative to the structure of Π/F_2 as the “ $\pi_1^{\mathcal{C}}(-)$ ” of a configuration space that arises from the description given in Remark 2.1.2].*

(iv) *The subgroup of Π topologically generated by F_1, F_2 is the fiber subgroup F_3 with profile $E'_3 \stackrel{\text{def}}{=} E'_1 \cup E'_2$. In particular, if E''_1, E''_2 are disjoint and of cardinality one, then F_1, F_2 topologically generate Π .*

(v) *In the situation of (iv), suppose that the length of F_1, F_2 is equal to 1. Then there exists a normal closed subgroup $K \subseteq \Pi$ satisfying the following properties: (a) $K \subseteq F_3$; (b) K is topologically normally generated in F_3 by a single element; (c) the images of F_1, F_2 in F_3/K commute.*

(vi) *F_2 is topologically generated by the fiber subgroups [of Π] of length 1 whose profiles are contained in E'_2 . In particular, Π is topologically generated by its fiber subgroups of length 1.*

Proof. Assertions (i), (ii) are immediate from the definitions [and Remark 2.1.2]. Next, let us consider assertion (vi). In light of assertions (i), (ii), it suffices to verify assertion (vi) in the case where $F_2 = \Pi$; also, we may assume without loss of generality that F_1 is of *length* 1. Then, by induction on n [cf. also assertion (i)], Π/F_1 is topologically generated by its fiber subgroups of length 1. Since the inverse image in Π of any fiber subgroup of length 1 of Π/F_1 is clearly a fiber subgroup of length 2, it follows [cf. assertions (i), (ii)] that we may assume without loss of generality that $n = 2$. But then it suffices to observe that if $F_\alpha, F_\beta \subseteq \Pi$ are fiber subgroups whose profiles $E'_\alpha, E'_\beta \subseteq E$ are disjoint subsets of length 1, then the natural morphism $F_\alpha \subseteq \Pi \rightarrow \Pi/F_\beta$ [which is simply the morphism induced on “ $\pi_1^{\mathcal{C}}(-)$ ’s” by an open immersion of hyperbolic curves] is a *surjection*. This completes the proof of assertion (vi). Now assertion (iv) follows formally from assertion (vi); also, in light of assertion (vi), assertion (iii) follows immediately from the definitions.

Finally, we consider assertion (v). First, let us observe that when $n = 2$, assertion (v) follows by observing that the kernel of the natural surjection $\pi_1^{\mathcal{C}}(X_2) \rightarrow \pi_1^{\mathcal{C}}(P_2)$ [cf. Definition 2.3, (ii)] is topologically normally generated by the inertia group of the *diagonal divisor* of X_2 , which is isomorphic to $\widehat{\mathbb{Z}}_{\mathcal{C}}(1)$ [hence topologically generated by a *single element*]. Now assertion (v) follows immediately for arbitrary n , by applying assertions (i), (ii), (iv). \circ

Remark 2.4.1. Note that it follows immediately from Proposition 2.2, (ii); Proposition 2.4, (i) [or, alternatively, (vi)], that the *fiber subgroups* of $\pi_1^{\mathcal{C}}(X_n)$ are *topologically finitely generated normal closed subgroups*.

Section 3: Direct Products of Profinite Groups

In the present §3, we study *quotients of products of profinite groups*. In particular, we show that, in certain cases, the product decomposition of a direct product of profinite groups is “*group-theoretic*”.

Definition 3.1. Let G be a profinite group. Then we shall say that G is *indecomposable* if, for any isomorphism of profinite groups $G \xrightarrow{\sim} H \times J$, where H, J are profinite groups, it follows that either H or J is the trivial group. We shall say that G is *strongly indecomposable* if every open subgroup of G is indecomposable.

Proposition 3.2. **(The Indecomposability of Surface Groups)** *Let \mathcal{C} be a nontrivial full formation. Then every almost pro- \mathcal{C} -surface group Π is strongly indecomposable.*

Proof. Since every open subgroup of an almost pro- \mathcal{C} -surface group is again an almost pro- \mathcal{C} -surface group, it suffices to show that Π is *indecomposable*. Suppose

that we have an isomorphism of profinite groups $\Pi \cong H \times J$, where H, J are *nonabelian* [since Π is *slim* — cf. Proposition 1.4!] *infinite* [again since Π is *slim*, hence does not contain any *nontrivial finite normal closed subgroups* — cf. §0] *profinite groups*. Note that since H, J , are *infinite*, it follows that for any open subgroup Π_1 , we may always replace Π by an open subgroup of Π_1 . In particular, [cf. Remark 1.2.1] we may assume, without loss of generality, that Π is a *pro- \mathcal{C} surface group* arising from a curve of *genus* ≥ 2 . Now we *claim* that for every prime number $l \in \Sigma_{\mathcal{C}}$, there exist finite quotients $H \twoheadrightarrow Q_H, J \twoheadrightarrow Q_J$ such that l divides the order of Q_H, Q_J . Indeed, suppose that l does *not* divide the order of any finite quotient of H . Then there exists a *proper* normal open subgroup $N_H \subseteq H$ such that if we set $N \stackrel{\text{def}}{=} N_H \times J \subseteq \Pi$, then the conjugation action of $\Pi/N \cong H/N_H$ on $N^{\text{ab}} \otimes \mathbb{Z}_l \cong (N_H^{\text{ab}} \otimes \mathbb{Z}_l) \times (J^{\text{ab}} \otimes \mathbb{Z}_l) \cong J^{\text{ab}} \otimes \mathbb{Z}_l$ is *trivial*, which, as was seen in the proof of Proposition 1.4, leads to a contradiction. This completes the proof of the *claim*.

Thus, by replacing Π by the *maximal pro- l quotient* of a suitable open subgroup of Π for some $l \in \Sigma_{\mathcal{C}}$ [and replacing \mathcal{C} by the primary formation determined by l], we may assume without loss of generality that Π, H, J are *pro- l groups*. Note, moreover, that since H, J are *nonabelian* pro- l groups, it follows that $\dim_{\mathbb{F}_l}(H^{\text{ab}} \otimes \mathbb{F}_l) \geq 2, \dim_{\mathbb{F}_l}(J^{\text{ab}} \otimes \mathbb{F}_l) \geq 2$ [cf., e.g., [RZ], Proposition 7.7.2]. On the other hand, observe that the cup product morphism

$$H^1(H, \mathbb{F}_l) \otimes H^1(J, \mathbb{F}_l) \rightarrow H^2(\Pi, \mathbb{F}_l)$$

is an *injection*. [Indeed, this follows immediately by considering the *spectral sequences* associated to the surjections $\Pi \cong H \times J \twoheadrightarrow J, H \twoheadrightarrow \{1\}$, where we note that the latter surjection may be regarded as a *quotient* of the former surjection.] But this implies that $\dim_{\mathbb{F}_l}(H^2(\Pi, \mathbb{F}_l)) \geq 2$, which [cf. Remark 1.2.2] is absurd. This completes the proof of Proposition 3.2. \circ

Remark 3.2.1. Note that the *strong indecomposability* of Proposition 3.2 may also be derived as an immediate consequence of Theorem 1.5, in light of the *slimness* of Proposition 1.4.

Proposition 3.3. (Quotients of Direct Products) *Let G_1, \dots, G_n be profinite groups, where $n \geq 1$ is an integer;*

$$\phi : \Pi \stackrel{\text{def}}{=} \prod_{i=1}^n G_i \twoheadrightarrow Q$$

*a surjection of profinite groups. Then there exist normal closed subgroups $H_i \subseteq G_i$ [for $i = 1, \dots, n$], $N \subseteq Q$ such that $N \subseteq Z(Q)$ [cf. §0], and the composite $\Pi \twoheadrightarrow Q/N$ of ϕ with the surjection $Q \twoheadrightarrow Q/N$ induces an **isomorphism***

$$\bar{\Pi} \stackrel{\text{def}}{=} \prod_{i=1}^n \bar{G}_i \xrightarrow{\sim} Q/N$$

— where we write $\overline{G}_i \stackrel{\text{def}}{=} G_i/H_i$. In particular, if Q is **center-free**, then we obtain an isomorphism $\overline{\Pi} \xrightarrow{\sim} Q$; if Q is **center-free** and **indecomposable**, then we obtain an isomorphism $\overline{G}_i \xrightarrow{\sim} Q$ for some $i \in \{1, \dots, n\}$.

Proof. Indeed, write $I \stackrel{\text{def}}{=} \text{Ker}(\phi) \subseteq \Pi$; $I_i \subseteq G_i$ for the inverse image of I via the natural injection $\iota_i : G_i \hookrightarrow \Pi$ into the i -th factor; $H_i \subseteq G_i$ for the image of I under the natural projection $\pi_i : \Pi \rightarrow G_i$ to the i -th factor [where $i \in \{1, \dots, n\}$]. Thus, we have inclusions

$$\Pi_I \stackrel{\text{def}}{=} \prod_{i=1}^n I_i \subseteq I \subseteq \Pi_H \stackrel{\text{def}}{=} \prod_{i=1}^n H_i \subseteq \Pi$$

inside Π . Now observe that the *commutator* of any element

$$(1, \dots, 1, g_i, 1, \dots, 1) \in \Pi$$

[i.e., all of whose components, except possibly the i -th component $g_i \in G_i$, are equal to 1] with an element $h \in I$ yields an element of I [since I is *normal* in Π] which lies in the image of ι_i , hence determines an element of $I_i \subseteq G_i$, which is in fact equal to the commutator $[g_i, \pi_i(h)] \in G_i$ [where we observe that $\pi_i(h) \in H_i$] computed in G_i . In particular, since $g_i \in G_i$ is *arbitrary*, and any element of H_i arises as such a “ $\pi_i(h)$ ”, it follows that the commutator subgroup $[G_i, H_i]$ is contained in I_i . But this implies that the commutator subgroup $[\Pi, \Pi_H]$ is normally generated in Π by elements of $\Pi_I \subseteq I$, hence [since I is *normal* in Π] is contained in I . Put another way, if we set $N \subseteq Q$ equal to the image in $\Pi/I \xrightarrow{\sim} Q$ of Π_H , then it follows that $N \subseteq Z(Q)$. On the other hand, it is immediate from the definitions that ϕ determines an isomorphism $\prod_{i=1}^n (G_i/H_i) \xrightarrow{\sim} Q/N$, as desired. \circ

Remark 3.3.1. Proposition 3.3 may be regarded as being motivated by the following elementary fact concerning *products of rings*: If R_1, \dots, R_n [where $n \geq 1$ is an integer] are [not necessarily commutative] rings with unity and

$$\phi : R \stackrel{\text{def}}{=} \prod_{i=1}^n R_i \rightarrow Q$$

is a *surjection* of rings with unity, then there exist two-sided ideals $I_i \subseteq R_i$ [for $i = 1, \dots, n$] such that ϕ induces an *isomorphism*

$$\overline{R} \stackrel{\text{def}}{=} \prod_{i=1}^n \overline{R}_i \xrightarrow{\sim} Q$$

— where we write $\overline{R}_i \stackrel{\text{def}}{=} R_i/I_i$. [Indeed, this follows immediately by observing that if, for $i = 1, \dots, n$, we write $e_i \in R$ for the element whose i -th component is 1 and whose other components are 0, then any element $f \in \text{Ker}(\phi)$ may be written in the

form $f = f \cdot e_1 + \dots + f \cdot e_n$, where each $f \cdot e_i \in \text{Ker}(\phi)$ [since $\text{Ker}(\phi)$ is a *two-sided ideal!*.]

Remark 3.3.2. Proposition 3.3 is due to the *second author*. We observe in passing that when, in the notation of Proposition 3.3, Q is an *almost pro- \mathcal{C} -surface group* for some nontrivial full formation \mathcal{C} [hence *slim* and *strongly indecomposable* — cf. Propositions 1.4, 3.2], and the G_i are *topologically finitely generated*, one may give a different proof of Proposition 3.3 by applying Theorem 1.5 to the images J_i of the various composites of ϕ with the natural inclusions $\iota_i : G_i \hookrightarrow \Pi$ — which allows one to conclude [in light of the *slimness* of Q !] that *only one* of the J_i [as i ranges over the integers $1, \dots, n$] can be *nontrivial*. In fact, this argument was the approach originally taken by the *first author* to proving Proposition 3.3 and, moreover, underlies the proof of the main result of this paper via the approach of the *second author* given in §6 below. On the other hand, this argument [unlike the *very elementary* proof of Proposition 3.3 given above!] has the drawback that it depends on the generalization of the result of *Lubotzky-Melnikov-van den Dries* given in Theorem 1.5. This drawback was pointed out by the second author to the first author when the first author first informed the second author of this restricted version of Proposition 3.3 and, indeed, served to motivate the second author to obtain the more elementary proof of Proposition 3.3 given above.

Corollary 3.4. (**Group-theoreticity of Product Decompositions**) *Let $n, m \geq 1$ be integers;*

$$G_1, \dots, G_n; \quad H_1, \dots, H_m$$

*nontrivial profinite groups which are **slim** and **strongly indecomposable** [e.g., **almost pro- \mathcal{C} -surface groups** for some **nontrivial full formation \mathcal{C}** — cf. Propositions 1.4, 3.2];*

$$G \subseteq \Pi_G \stackrel{\text{def}}{=} \prod_{i=1}^n G_i; \quad H \subseteq \Pi_H \stackrel{\text{def}}{=} \prod_{j=1}^m H_j$$

open subgroups;

$$\alpha : G \xrightarrow{\sim} H$$

*an **isomorphism** of profinite groups. For $i = 1, \dots, n; j = 1, \dots, m$, write $G_i^{\rightarrow} \subseteq G_i, H_j^{\rightarrow} \subseteq H_j$ for the respective images of G, H via the natural projections $\Pi_G \twoheadrightarrow G_i, \Pi_H \twoheadrightarrow H_j$. Then $n = m$; there exist a **unique permutation** σ of the set $\{1, \dots, n\}$ and **unique isomorphisms** of profinite groups $\alpha_i : G_i^{\rightarrow} \xrightarrow{\sim} H_{\sigma(i)}^{\rightarrow}$ [for $i = 1, \dots, n$] such that the restriction of [the composite with the inclusion into Π_H of] the isomorphism*

$$(\alpha_1, \dots, \alpha_n) : \left(\Pi_G \supseteq \right) \prod_{i=1}^n (G_i^{\rightarrow}) \xrightarrow{\sim} \prod_{i=1}^n (H_{\sigma(i)}^{\rightarrow}) \left(\subseteq \Pi_H \right)$$

to G coincides with [the composite with the inclusion into Π_H of] α .

Proof. First, we observe that the *uniqueness* assertions follow immediately from the *nontriviality* and *slimness* of the profinite groups $G_1, \dots, G_n, H_1, \dots, H_m$. Thus, it suffices to verify the *existence* of σ and the α_i . For $i = 1, \dots, n$; $j = 1, \dots, m$, write

$$G_i^{\bar{}} \subseteq \Pi_G; \quad H_j^{\bar{}} \subseteq \Pi_H$$

for the respective intersections of G, H with the images of the natural injections $G_i \hookrightarrow \Pi_G, H_j \hookrightarrow \Pi_H$;

$$G_i^{\neq} \subseteq \Pi_G; \quad H_j^{\neq} \subseteq \Pi_H$$

for the respective intersections of G, H with the kernels of the natural projections $\Pi_G \twoheadrightarrow G_i, \Pi_H \twoheadrightarrow H_j$. Now we *claim* that for each $j = 1, \dots, m$, the kernel of the composite

$$\psi_j : G \rightarrow H_j$$

of α with the natural projection $(H \subseteq) \Pi_H \twoheadrightarrow H_j$ contains G_i^{\neq} , for a *unique* $i \in \{1, \dots, n\}$. Indeed, since the image of ψ_j is *open*, hence *slim*, it follows [cf. §0] that this image has *no nontrivial finite normal closed subgroups*; since the G_i^{\neq} are *normal closed subgroups* of G , it thus suffices to prove that the kernel of the *restriction* of ψ_j to the open subgroup of $G \subseteq \Pi_G$ determined by the *direct product* of the $G_{i'}^{\bar{}}$ [for $i' = 1, \dots, n$] contains the intersection of this open subgroup with G_i^{\neq} , for a *unique* i . But [since H_j is *slim* and *strongly indecomposable*] this follows formally from Proposition 3.3. This completes the proof of the *claim*.

Note, moreover, that [in the notation of the *claim*] the assignment $j \mapsto i$ determines a map $\{1, \dots, m\} \rightarrow \{1, \dots, n\}$, which, in light of the *injectivity* of α , is easily verified to be *surjective*. But this implies that $m \geq n$; thus, by applying this argument to α^{-1} , we obtain that $m = n$. In particular, the map $\{1, \dots, m\} \rightarrow \{1, \dots, n\}$ considered above is a *bijection*, whose inverse we denote by σ . By rearranging the indices, we may assume without loss of generality that σ is the *identity*.

Now it follows from the definition of [the map that gave rise to] σ that we obtain a *surjection*

$$\alpha_i : G_i^{\twoheadrightarrow} \twoheadrightarrow H_i^{\twoheadrightarrow}$$

for each $i = 1, \dots, n$, such that the restriction of [the composite with the inclusion into Π_H of] the *surjection*

$$(\alpha_1, \dots, \alpha_n) : \left(\Pi_G \supseteq \right) \prod_{i=1}^n (G_i^{\twoheadrightarrow}) \twoheadrightarrow \prod_{i=1}^n (H_i^{\twoheadrightarrow}) \left(\subseteq \Pi_H \right)$$

to G coincides with [the composite with the inclusion into Π_H of] α . In particular, since α is injective, it follows that the kernel of each α_i is a *finite closed normal subgroup* of an open subgroup of G_i . Thus, by the *slimness* of G_i , we conclude [cf. §0] that the α_i are *injective*, as desired. \circ

Section 4: Product-theoretic Quotients

In the present §4, we show that in the case of genus ≥ 2 , the [closure of the] *commutator subgroup* of a *product-theoretic* open subgroup of a configuration space group is, up to torsion, again *product-theoretic* [cf. Theorem 4.7]. This result, combined with the theory of §3, implies a rather strong result, in the case of genus ≥ 2 , concerning the *group-theoreticity* of the various *fiber subgroups* associated to a configuration space group [cf. Corollary 4.8].

Let Y be a *connected smooth variety* over an *algebraically closed field* k which [for simplicity] we assume to be *of characteristic zero*.

Definition 4.1. Let $j \geq 1$ be an integer. Then we shall refer to Y as *j -good* if for every positive integer $j' \leq j$ and every class

$$\eta \in H_{\text{ét}}^{j'}(Y, \mathbb{Z}/N\mathbb{Z})$$

[where “ $H_{\text{ét}}^{j'}(-)$ ” denotes étale cohomology, and $N \geq 1$ is an integer], there exists a finite étale covering $Y' \rightarrow Y$ such that $\eta|_{Y'} = 0$.

Remark 4.1.1. As is well-known, it follows immediately from the Hochschild-Serre spectral sequence in étale cohomology [cf., e.g., [Milne], p. 105, Theorem 2.20] that one has a *natural isomorphism*

$$H^{j'}(\pi_1(Y), \widehat{\mathbb{Z}}) \xrightarrow{\sim} H_{\text{ét}}^{j'}(Y, \widehat{\mathbb{Z}})$$

for all nonnegative integers $j' \leq j$ whenever Y is *j -good*. Also, we observe that it is immediate from the definitions that the condition “1-good” is *vacuous*.

Let

$$f : Z \rightarrow Y$$

be a *family of hyperbolic curves* over Y ; $y \in Y(k)$. We shall denote fibers over y by means of a subscript “ y ”. Suppose that we have also been given a *section*

$$s : Y \rightarrow Z$$

of f , whose image we denote by $D_s \subseteq Z$. Write $U_Z \subseteq Z$ for the open subscheme given by the *complement* of D_s ; $\mathcal{L} \stackrel{\text{def}}{=} \mathcal{O}_Z(D_s)$; $\mathbb{L}^\times \rightarrow Z$ for the complement of the zero section of the geometric line bundle determined by \mathcal{L} ;

$$U_Z \rightarrow \mathbb{L}^\times$$

for the morphism determined by the natural inclusion $\mathcal{O}_Z \hookrightarrow \mathcal{O}_Z(D_s) = \mathcal{L}$. Thus, $U_Z \rightarrow Y$ is also a *family of hyperbolic curves*. Now if we denote by “ $\pi_1(-)$ ” the

étale fundamental group [for an appropriate choice of basepoint], then we have a *natural commutative diagram*

$$\begin{array}{ccccccccc}
1 & \longrightarrow & \pi_1((U_Z)_y) & \longrightarrow & \pi_1(U_Z) & \longrightarrow & \pi_1(Y) & \longrightarrow & 1 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
1 & \longrightarrow & \pi_1(\mathbb{L}_y^\times) & \longrightarrow & \pi_1(\mathbb{L}^\times) & \longrightarrow & \pi_1(Y) & \longrightarrow & 1 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
1 & \longrightarrow & \pi_1(Z_y) & \longrightarrow & \pi_1(Z) & \longrightarrow & \pi_1(Y) & \longrightarrow & 1
\end{array}$$

in which the first and third horizontal sequences are *exact* [cf. Proposition 2.2, (iii)]. Write $I_s \subseteq \pi_1(U_Z)$ for the *inertia group* [well-defined up to conjugation in $\pi_1(U_Z)$] associated to the divisor D_s . Thus, $I_s \cong \widehat{\mathbb{Z}}(1)$ [where the “(1)” denotes a “Tate twist”].

Lemma 4.2. (The Line Bundle Associated to a Cusp) *In the notation of the above discussion, suppose further that Y is j -good, for some integer $j \geq 2$. Then:*

(i) Z is j -good.

(ii) $\pi_1(\mathbb{L}^\times)$ fits into a short exact sequence:

$$1 \rightarrow \widehat{\mathbb{Z}}(1) \rightarrow \pi_1(\mathbb{L}^\times) \rightarrow \pi_1(Z) \rightarrow 1$$

Moreover, the resulting extension class $\in H^2(\pi_1(Z), \widehat{\mathbb{Z}}(1)) \cong H_{\text{ét}}^2(Z, \widehat{\mathbb{Z}}(1))$ [cf. (i); Remark 4.1.1] is the **first Chern class** of the line bundle \mathcal{L} .

(iii) The sequence $1 \rightarrow \pi_1(\mathbb{L}_y^\times) \rightarrow \pi_1(\mathbb{L}^\times) \rightarrow \pi_1(Y) \rightarrow 1$ of the above commutative diagram is **exact**.

(iv) The morphism of fundamental groups $\pi_1(U_Z) \rightarrow \pi_1(\mathbb{L}^\times)$ induces an **isomorphism** $I_s \xrightarrow{\sim} \text{Ker}(\pi_1(\mathbb{L}^\times) \rightarrow \pi_1(Z))$. In particular, the vertical arrows of the commutative diagram of the above discussion are **surjections**.

(v) Write $\pi_1(U_Z/Z) \stackrel{\text{def}}{=} \text{Ker}(\pi_1(U_Z) \rightarrow \pi_1(Z)) \subseteq \pi_1((U_Z)_y)$. Then the quotient of $\pi_1(U_Z/Z)$ by

$$\pi_1(U_Z/\mathbb{L}^\times) \stackrel{\text{def}}{=} \text{Ker}(\pi_1(U_Z) \rightarrow \pi_1(\mathbb{L}^\times)) \subseteq \pi_1(U_Z/Z) \quad (\subseteq \pi_1(U_Z))$$

is the **maximal quotient** of $\pi_1(U_Z/Z)$ on which the conjugation action by $\pi_1((U_Z)_y)$ is **trivial**.

Proof. First, we consider assertion (i). In light of the exact sequence $1 \rightarrow \pi_1(Z_y) \rightarrow \pi_1(Z) \rightarrow \pi_1(Y) \rightarrow 1$ [together with the Leray-Serre spectral sequence for $Z \rightarrow Y$], it follows immediately that to show that Z is j -good, it suffices to show that Z_y is

j -good. But this follows immediately from the fact that the cohomological dimension of Z_y is equal to 1 when Z_y is *affine* [cf., e.g., [Milne], p. 253, Theorem 7.2] and from the well-known isomorphism $H_{\text{ét}}^2(Z_y, \mathbb{Z}/N\mathbb{Z}) \cong (\mathbb{Z}/N\mathbb{Z})(-1)$ determined by considering fundamental classes of points [together with the fact that the cohomological dimension of Z_y is equal to 2 — cf., e.g., [Milne], p. 276, Theorem 11.1], when Z_y is *proper*. This completes the proof of assertion (i).

In light of assertion (i), assertion (ii) follows from [Mzk2], Lemmas 4.4, 4.5. Assertion (iii) follows immediately by considering the natural commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \widehat{\mathbb{Z}}(1) & \longrightarrow & \pi_1(\mathbb{L}_y^\times) & \longrightarrow & \pi_1(Z_y) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \widehat{\mathbb{Z}}(1) & \longrightarrow & \pi_1(\mathbb{L}^\times) & \longrightarrow & \pi_1(Z) \longrightarrow 1 \end{array}$$

[in which the rows are *exact*, by assertion (ii); the vertical arrow on the left is an *isomorphism*], together with the exact sequence $1 \rightarrow \pi_1(Z_y) \rightarrow \pi_1(Z) \rightarrow \pi_1(Y) \rightarrow 1$. Assertion (iv) (respectively, (v)) follows immediately from the argument of the proof of [Mzk5], Lemma 4.2, (ii) (respectively, [Mzk5], Lemma 4.2, (iii)). \circ

Now let l be a *prime number*; suppose that Y is *2-good*. Also, let us suppose that, for $i = 1, \dots, m$ [where $m \geq 1$ is an integer], we have been given a *section*

$$s_i : Y \rightarrow Z$$

of f , whose image we denote by $D_{s_i} \subseteq Z$. Write $U_i \subseteq Z$ for the open subscheme given by the *complement* of D_{s_i} ; $W_Z \stackrel{\text{def}}{=} \bigcap_{i=1}^m U_i \subseteq Z$; $\mathcal{L}_i \stackrel{\text{def}}{=} \mathcal{O}_Z(D_{s_i})$; $\mathbb{L}_i^\times \rightarrow Z$ for the complement of the zero section of the geometric line bundle determined by \mathcal{L}_i ;

$$W_Z \rightarrow \mathbb{L}_i^\times$$

for the morphism determined by the natural inclusion $\mathcal{O}_Z \hookrightarrow \mathcal{O}_Z(D_{s_i}) = \mathcal{L}_i$. Also, let us suppose that $W_Z \rightarrow Y$ is a *family of hyperbolic curves* [i.e., that the images of the s_i do not intersect]. By forming the quotient of the exact sequence of Lemma 4.2, (ii), by the pro-prime-to- l portion of $\widehat{\mathbb{Z}}(1)$, we obtain *extensions*

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{Z}_l(1) & \longrightarrow & \mathbb{E}_{i,y} & \longrightarrow & \pi_1(Z_y) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathbb{Z}_l(1) & \longrightarrow & \mathbb{E}_i & \longrightarrow & \pi_1(Z) \longrightarrow 1 \end{array}$$

for $i = 1, \dots, m$. Also, let us write

$$\kappa_i \in H_{\text{ét}}^2(Z, \mathbb{Z}_l(1))$$

for the *fundamental class* associated to D_{s_i} [i.e., the *first Chern class* of the line bundle \mathcal{L}_i — cf. Lemma 4.2, (ii)].

Lemma 4.3. (Multi-section Splittings) *In the notation of the above discussion:*

(i) *The natural homomorphism*

$$\pi_1(W_Z) \rightarrow \prod_{i=1}^m \mathbb{E}_i$$

[where the product is a fiber product over $\pi_1(Z)$] **is surjective.**

(ii) *The natural quotient $\pi_1(W_Z) \twoheadrightarrow \pi_1(W_Z)^{\text{ab}} \otimes \mathbb{Z}_l$ factors through the quotient determined by the surjection of (i).*

(iii) *For $i = 1, \dots, m$, let $\lambda_i \in \mathbb{Z}_l$. Then there **exists** a surjection $\pi_1(W_Z) \twoheadrightarrow \mathbb{Z}_l(1)$ — which, by (ii), necessarily **factors** through the surjection of (i), hence determines a **surjection***

$$\prod_{i=1}^m \mathbb{E}_i \twoheadrightarrow \mathbb{Z}_l(1)$$

— that restricts to multiplication by λ_i on the copy of $\mathbb{Z}_l(1)$ in \mathbb{E}_i if and only if the class

$$\sum_{i=1}^m \lambda_i \cdot \kappa_i \in H_{\text{ét}}^2(Z, \mathbb{Z}_l(1))$$

vanishes.

Proof. First, we consider assertion (i). In light of the exact sequences of Proposition 2.2, (iii), and Lemma 4.2, (iii), it suffices to show the surjectivity of $\pi_1((W_Z)_y) \rightarrow \prod_{i=1}^m \mathbb{E}_{i,y}$. But this follows immediately, in light of Lemma 4.2, (iv), by considering the various *inertia groups* $\subseteq \pi_1((W_Z)_y)$ of the cusps of $(W_Z)_y$. This completes the proof of assertion (i). Assertion (ii) follows immediately, in light of Lemma 4.2, (iv) [and induction on n], from the fact that the kernel of the natural surjection $\pi_1(W_Z) \twoheadrightarrow \pi_1(Z)$ is *topologically normally generated* by the *inertia groups of cusps*. Finally, we observe that assertion (iii) follows immediately from the definitions. \circ

Lemma 4.4. (The Section Arising from the Graph of a Morphism) *In the notation of the above discussion, suppose further that $Z \rightarrow Y$ is given by the projection to the second factor $C \times_k C \rightarrow C$, where we write $C \stackrel{\text{def}}{=} Z_y$, that C is **proper**, and that $s : Y \rightarrow Z$ is given by the **graph** of a k -morphism $\sigma : C \rightarrow C$. Then the component of the first Chern class of \mathcal{L} in the middle direct summand of*

$$H_{\text{ét}}^2(Z, \mathbb{Z}_l(1)) \cong H_{\text{ét}}^2(C, \mathbb{Z}_l(1)) \oplus (H_{\text{ét}}^1(C, \mathbb{Z}_l) \otimes H_{\text{ét}}^1(C, \mathbb{Z}_l(1))) \oplus H_{\text{ét}}^2(C, \mathbb{Z}_l(1))$$

[cf. the Künneth isomorphism in étale cohomology, discussed, e.g., in [Milne], p. 258, Theorem 8.5] is given by applying the endomorphism $\sigma^* \otimes \text{id}$ of the module $H_{\text{ét}}^1(C, \mathbb{Z}_l) \otimes H_{\text{ét}}^1(C, \mathbb{Z}_l(1))$ to the element of this module determined by the morphism $\text{Hom}_{\mathbb{Z}_l}(H_{\text{ét}}^1(C, \mathbb{Z}_l(1)), \mathbb{Z}_l) \rightarrow H_{\text{ét}}^1(C, \mathbb{Z}_l)$ given by the inverse of the morphism

$H_{\text{ét}}^1(C, \mathbb{Z}_l) \rightarrow \text{Hom}_{\mathbb{Z}_l}(H_{\text{ét}}^1(C, \mathbb{Z}_l(1)), \mathbb{Z}_l)$ arising from the cup product $H_{\text{ét}}^1(C, \mathbb{Z}_l) \otimes H_{\text{ét}}^1(C, \mathbb{Z}_l(1)) \rightarrow H_{\text{ét}}^2(C, \mathbb{Z}_l(1)) \cong \mathbb{Z}_l$ in étale cohomology.

Proof. Indeed, this follows immediately from [Milne], p. 287, Lemma 12.2. \circ

Lemma 4.5. (Linear Independence for Vector Spaces) *Let G be a finite group, whose order we denote by $|G|$; K a field; V a finite-dimensional K -vector space equipped with a linear action by G such that the G -module V contains the regular representation of G as a direct summand; $N \geq 1$ an integer. Write*

$$W \stackrel{\text{def}}{=} V \oplus \dots \oplus V$$

for the direct sum of N copies of V ; $\iota_i \in \text{Hom}_K(V, W)$ [where $i = 1, \dots, N$] for the inclusion $V \hookrightarrow W$ into the i -th factor. Then the $N \cdot |G|$ elements

$$\iota_i \circ g$$

[where $i = 1, \dots, N$; $g \in G$] of $\text{Hom}_K(V, W)$ are **linearly independent**.

Proof. Indeed, any nontrivial linear relation between these elements implies — by applying the various linear morphisms $\text{Hom}_K(V, W) \rightarrow \text{Hom}_K(V, V)$ obtained by *projecting* onto the various factors of V in W — a nontrivial linear relation between the endomorphisms $\in \text{Hom}_K(V, V)$ determined by the elements of G , in contradiction to the assumption that the G -module V contains the *regular representation* of G as a direct summand. \circ

Lemma 4.6. (Linear Independence for Configuration Spaces) *In the notation of the above discussion, suppose further that:*

(a) *there exists a commutative diagram*

$$\begin{array}{ccc} Z & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X \times_k X_n & \longrightarrow & X_n \end{array}$$

where the upper horizontal arrow is the given morphism $Z \rightarrow Y$; the lower horizontal arrow is the projection to the second factor; $n \geq 1$ is an integer; X_n is the **n -th configuration space** associated to some **hyperbolic curve** X over k ; the vertical arrows are **finite étale Galois coverings** arising from the coverings of $X \times_k X_n, X_n$ determined by taking the **direct product** of copies of a finite étale Galois covering $Z_0 \rightarrow X$ [so Z_y may be **identified** with Z_0];

(b) *the genus of the compactification B of X is ≥ 2 ;*

- (c) if we write $C \rightarrow B$ for the **normalization** of B in Z_0 , then we have $m = n \cdot \deg(C/B)$, and the $s_i : Y \rightarrow Z$ are the various liftings of the n **tautological sections** $X_n \rightarrow X \times_k X_n$ arising from the definition of the configuration space X_n .

[Thus, the fact that $W_Z \rightarrow Y$ is a **family of hyperbolic curves** follows immediately from Remark 2.1.2; the fact that X_n , hence also Y , is **2-good** follows, by induction on n , from Lemma 4.2, (i). Moreover, W_Z forms a finite étale covering of X_{n+1} that arises from a **product-theoretic** open subgroup of $\pi_1(X_{n+1})$.] Then:

(i) The morphisms of the commutative diagram of (a) determine an **isomorphism** of k -schemes $Z \xrightarrow{\sim} Z_0 \times_k Y$.

(ii) The isomorphism of (i) determines an isomorphism of **first cohomology groups** $H_{\text{ét}}^1(Z, \mathbb{Z}_l) \xrightarrow{\sim} H_{\text{ét}}^1(Z_0, \mathbb{Z}_l) \times H_{\text{ét}}^1(Y, \mathbb{Z}_l)$.

(iii) The images of the κ_i in $H_{\text{ét}}^2(Z, \mathbb{Q}_l(1))$ are **linearly independent** [over \mathbb{Q}_l].

Proof. Note that the projection to the first factor $X \times_k X_n \rightarrow X$ determines a morphism $Z \rightarrow Z_0 (\subseteq C)$. Thus, we obtain an isomorphism as in assertion (i); assertion (ii) follows immediately from assertion (i). Now it remains to verify assertion (iii). Suppose that the section $s : Y \rightarrow Z$ arises from a point $\in Z_0(k)$. Write $\kappa \in H_{\text{ét}}^2(Z, \mathbb{Z}_l(1))$ for the *fundamental class* associated to D_s . For $i = 1, \dots, m$, set $\kappa'_i \stackrel{\text{def}}{=} \kappa_i - \kappa$. Next, let us *observe* that κ and the κ_i all map [cf. the Leray-Serre spectral sequence for $Z \rightarrow Y$] to the *same element* of $H_{\text{ét}}^2(Z_y, \mathbb{Q}_l(1))$ — a \mathbb{Q}_l -vector space of dimension 0 [cf., e.g., [Milne], p. 253, Theorem 7.2] if Z_y is *affine* and dimension 1 [cf., e.g., [Milne], p. 276, Theorem 11.1, (a)] if Z_y is *proper*. In particular, it follows that κ'_i maps to 0 in $H_{\text{ét}}^2(Z_y, \mathbb{Q}_l(1))$. Note, moreover, that [since $\kappa = 0$ whenever Z_y is *affine*] this *observation* also implies that to verify the linear independence of the images of the κ_i in $H_{\text{ét}}^2(Z, \mathbb{Q}_l(1))$, it suffices to verify the linear independence of the images of the κ'_i in $H_{\text{ét}}^2(Z, \mathbb{Q}_l(1))$. Thus, we conclude that the κ'_i determine classes

$$\eta_i \in H_{\text{ét}}^1(Y, H_{\text{ét}}^1(Z_y, \mathbb{Q}_l(1))) \cong H_{\text{ét}}^1(Y, \mathbb{Q}_l) \otimes H_{\text{ét}}^1(Z_0, \mathbb{Q}_l(1))$$

[cf. the Leray-Serre spectral sequence for $Z \rightarrow Y$], and that to verify the linear independence of the images of the κ_i in $H_{\text{ét}}^2(Z, \mathbb{Q}_l(1))$, it suffices to verify that the η_i are *linearly independent*.

On the other hand, it follows immediately from the definitions that the κ_i arise as pull-backs via the various projections $Y \rightarrow Z_0 \hookrightarrow C$, $Z \rightarrow Z_0 \hookrightarrow C$ of the classes [cf. Lemma 4.4] determined by the *graphs* $\subseteq C \times_k C$ of the various $\sigma : C \rightarrow C$, for $\sigma \in \text{Gal}(C/B)$. In particular, the η_i arise as pull-backs via these various projections of the classes in

$$H_{\text{ét}}^1(C, \mathbb{Q}_l) \otimes H_{\text{ét}}^1(C, \mathbb{Q}_l(1)) (\hookrightarrow H_{\text{ét}}^1(Y, \mathbb{Q}_l) \otimes H_{\text{ét}}^1(Z_0, \mathbb{Q}_l(1)))$$

determined [cf. Lemma 4.4] by the *graphs* of the various $\sigma \in G \stackrel{\text{def}}{=} \text{Gal}(C/B)$. On the other hand, by Proposition 1.3 [cf. our assumption that the genus of B is $\geq 2!$], it follows that the G -module $V \stackrel{\text{def}}{=} H_{\text{ét}}^1(C, \mathbb{Q}_l)$ contains the *regular representation* of G as a direct summand. Note, moreover, that the n inclusions $V = H_{\text{ét}}^1(C, \mathbb{Q}_l) \hookrightarrow H_{\text{ét}}^1(Y, \mathbb{Q}_l)$ [determined up to composition with the action of G on V] arising from the n projections $X_n \rightarrow X$ determine a map of \mathbb{Q}_l -vector spaces

$$\left(\bigoplus H_{\text{ét}}^1(C, \mathbb{Q}_l) \right) \rightarrow H_{\text{ét}}^1(Y, \mathbb{Q}_l)$$

[where the direct sum is over n copies of $H_{\text{ét}}^1(C, \mathbb{Q}_l)$] which is *injective*. Thus, we are, in effect, in the situation of Lemma 4.5, so the linear independence of the η_i follows from the linear independence asserted in Lemma 4.5. \circ

Theorem 4.7. (Strongly Torsion-free Pro-solvable Product-theoreticity)
 Let X be a **hyperbolic curve of genus ≥ 2 over an algebraically closed field k of characteristic zero**; $n \geq 1$ an integer; X_n the **n -th configuration space associated to X** ; $H \subseteq \pi_1(X_n)$ a **product-theoretic open subgroup**; G a **strongly torsion-free pro-solvable profinite group**. Then the kernel of any continuous homomorphism

$$H \rightarrow G$$

is **product-theoretic**.

Proof. First, we *claim* that it suffices to verify Theorem 4.7 in the case where $G = \mathbb{Z}_l$ [for some prime number l]. Indeed, since G is topologically finitely generated [cf. Definition 1.1, (iii)], and arbitrary intersections of product-theoretic closed subgroups of $\pi_1(X_n)$ are clearly product-theoretic, Theorem 4.7 for arbitrary [torsion-free] *abelian* G follows immediately from the case “ $G = \mathbb{Z}_l$ ”. Thus, by replacing H, G *successively* by appropriate open subgroups of H, G , Theorem 4.7 for arbitrary [strongly torsion-free] *pro-solvable* G follows immediately from the [torsion-free] abelian case. This completes the proof of the *claim*. Thus, in the following, we assume that $G = \mathbb{Z}_l$.

Now observe that Theorem 4.7 is vacuous for $n = 1$. Thus, by induction on n , it suffices to verify Theorem 4.7 for “ $n + 1$ ” under the assumption that it holds for “ n ”. Next, let us observe that it follows immediately from the definition of “product-theoretic” that any covering of X_{n+1} that arises from a product-theoretic open subgroup $J \subseteq \pi_1(X_{n+1})$ is *dominated* by a covering of the form “ $W_Z \rightarrow X_{n+1}$ ” for W_Z as in Lemma 4.6. Thus, to complete the proof of Theorem 4.7, it suffices to show that the kernel of any quotient $J \twoheadrightarrow J^{\text{ab}} \otimes \mathbb{Z}_l \twoheadrightarrow \mathbb{Z}_l$ is *product-theoretic*, for J corresponding to a covering “ $W_Z \rightarrow X_{n+1}$ ” as in Lemma 4.6. In particular, by applying Lemma 4.3, (iii), in light of the *linear independence* asserted in Lemma 4.6, (iii), and the *induction hypothesis* [which may be applied to “ Y ”, via Lemma 4.6, (ii)], we conclude that the kernel of such a quotient $J \twoheadrightarrow \mathbb{Z}_l$ is *product-theoretic*, as desired. \circ

Remark 4.7.1. Note that Theorem 4.7 is *false* if the genus of X is < 2 and $n \geq 2$. Indeed, to construct a counter-example for arbitrary $n \geq 2$, it suffices to construct a counter-example for $n = 2$. If, moreover, U is the hyperbolic curve determined by an open subscheme of X , then consideration of the natural morphism $U_2 \hookrightarrow X_2$ shows that the existence of a counter-example for X_2 implies the existence of a counter-example for U_2 . Thus, we may assume, without loss of generality, that $n = 2$, and X is either of type $(0, 3)$ or of type $(1, 1)$. But, in either of these cases, it is well-known that there exists a dominant map $X_2 \rightarrow X$ that extends to a map $X \times_k X \rightarrow B$ [where B is a compactification of X] that maps the open subscheme $X_2 \subseteq X \times_k X$ into $X \subseteq B$. Thus, by pulling back an appropriate infinite cyclic covering of some finite étale covering of X , one obtain an infinite cyclic covering of some finite étale covering of $X \times_k X$ that is [infinitely] ramified over the diagonal of $X \times_k X$.

Corollary 4.8. (Group-theoreticity of Projections of Configuration Spaces I) *Let \mathcal{C} be a PT-formation. For $\xi = \alpha, \beta$, let X^ξ be a hyperbolic curve of genus ≥ 2 over an algebraically closed field k_ξ of characteristic zero; $n_\xi \geq 1$ an integer; $X_{n_\xi}^\xi$ the n_ξ -th configuration space associated to X^ξ ; E_ξ the index set of $X_{n_\xi}^\xi$; $H_\xi \subseteq \Pi^\xi \stackrel{\text{def}}{=} \pi_1^{\mathcal{C}}(X_{n_\xi}^\xi)$ a product-theoretic open subgroup. Let*

$$\gamma : H_\alpha \xrightarrow{\sim} H_\beta$$

be an isomorphism of profinite groups. Then γ induces a bijection $\sigma : E_\alpha \xrightarrow{\sim} E_\beta$ [so $n_\alpha = n_\beta$] such that

$$\gamma(F_\alpha \cap H_\alpha) = F_\beta \cap H_\beta$$

for all fiber subgroups $F_\alpha \subseteq \Pi^\alpha$, $F_\beta \subseteq \Pi^\beta$, whose respective profiles $E'_\alpha \subseteq E_\alpha$, $E'_\beta \subseteq E_\beta$ correspond via σ .

Proof. First, let us observe that to complete the proof of Corollary 4.8, it suffices to construct a bijection $\sigma : E_\alpha \xrightarrow{\sim} E_\beta$ [so $n_\alpha = n_\beta$] such that

$$\gamma(F_\alpha \cap H_\alpha) = F_\beta \cap H_\beta$$

for all fiber subgroups $F_\alpha \subseteq \Pi^\alpha$, $F_\beta \subseteq \Pi^\beta$ of co-length one whose respective profiles correspond via σ . Indeed, this follows immediately by applying *induction* on $n \stackrel{\text{def}}{=} n_\alpha = n_\beta$ [cf. also Proposition 2.4, (i), (ii)].

Next, for $j = 1, \dots, n_\xi$, let us write

$$K_j^\xi \subseteq H_\xi$$

for the intersection with H_ξ of the fiber subgroup $\subseteq \Pi^\xi$ of co-length one with co-profile given by the element of E_ξ labeled by j . Thus, [cf. Proposition 2.4, (iv);

the fact that fiber subgroups of co-length one are *normal closed subgroups of infinite index*] for *distinct* $j, j' \in \{1, \dots, n_\xi\}$, $K_j^\xi, K_{j'}^\xi$ topologically generate an *open subgroup* of Π^ξ ; in particular, K_j^ξ is *not* contained in $K_{j'}^\xi$.

Now we *claim* that to complete the proof of Corollary 4.8, it suffices to prove that the following *statement* holds [in general]:

For each $i \in E_\alpha$, there exists a $j \in E_\beta$ such that $K_j^\beta \subseteq \gamma(K_i^\alpha)$.

Indeed, by applying this statement to γ, γ^{-1} , we conclude that for each $i \in E_\alpha$, there exist $j \in E_\beta, i' \in E_\alpha$ such that $\gamma(K_{i'}^\alpha) \subseteq K_j^\beta \subseteq \gamma(K_i^\alpha)$, hence that $K_{i'}^\alpha \subseteq K_i^\alpha$. But, as observed above, this implies that $i' = i$, hence that $K_j^\beta = \gamma(K_i^\alpha)$. Moreover, this relation “ $K_j^\beta = \gamma(K_i^\alpha)$ ” determines an assignment $i \mapsto j$, hence a mapping $\sigma : E_\alpha \rightarrow E_\beta$, which is a *bijection*, relative to which intersections with H_α, H_β of fiber subgroups of co-length one with corresponding profiles correspond via γ . This completes the proof of the *claim*.

To verify the “*statement*” of the above *claim*, we reason as follows: Let $l \in \Sigma_{\mathcal{C}}$. Write $H_\alpha/K_i^\alpha \twoheadrightarrow G$ for the *maximal pro- l quotient* of H_α/K_i^α ;

$$\phi : H_\beta \xrightarrow{\sim} H_\alpha \twoheadrightarrow H_\alpha/K_i^\alpha \twoheadrightarrow G$$

for the surjection determined by γ^{-1} . Then observe that since H_α/K_i^α is a *pro- \mathcal{C} surface group*, it follows that G is a *pro- l surface group*, hence *strongly torsion-free* [cf. Remark 1.2.2] and *pro-solvable* [cf. Remark 1.1.3]. Thus, it follows from Theorem 4.7 that ϕ factors through the quotient $H_\beta \twoheadrightarrow Q_\beta$ determined by the quotient “ $\pi_1^{\mathcal{C}}(X_{n_\beta}^\beta) \twoheadrightarrow \pi_1^{\mathcal{C}}(P_{n_\beta}^\beta)$ ” [i.e., the image of $H_\beta \subseteq \pi_1^{\mathcal{C}}(X_{n_\beta}^\beta)$ in $\pi_1^{\mathcal{C}}(P_{n_\beta}^\beta)$] corresponding to the quotient that was denoted “ $\pi_1^{\mathcal{C}}(X_n) \twoheadrightarrow \pi_1^{\mathcal{C}}(P_n)$ ” in Definition 2.3, (ii). In particular, since Q_β admits an open subgroup with a direct product decomposition induced by the natural direct product decomposition of $\pi_1^{\mathcal{C}}(P_{n_\beta}^\beta)$, it thus follows [since G is *slim* and *strongly indecomposable* — cf. Propositions 1.4, 3.2] from Proposition 3.3 that there exists a $j \in E_\beta$ such that the image of K_j^β in G is a *finite normal closed subgroup*, hence *trivial* [since G is *slim* — cf. Proposition 1.4, §0]. But this implies that the image of K_j^β in the *pro- \mathcal{C} -surface group* H_α/K_i^α is a *topologically finitely generated* [cf. Proposition 2.2, (ii)] *normal closed subgroup*, which [since G is *infinite*] is *of infinite index*, hence — by Theorem 1.5 — *trivial*. This completes the proof of Corollary 4.8. \circ

Remark 4.8.1. It is interesting to note that in [NT], Theorem 3.1 [cf., especially, [NT], Lemmas 3.3 – 3.6], a certain analogue of Theorem 4.8 is shown for *graded Lie algebras*. More generally, the idea of studying configuration spaces from an *abelian* point of view dates back at least to [Naka].

Remark 4.8.2. The original motivation, for the first author, for developing the theory applied to prove Corollary 4.8 was the idea that by combining Corollary

4.8 with the techniques of [Mzk6], [Mzk7], one could obtain results in the *absolute anabelian geometry of configuration spaces over p -adic local fields*. It is the intention of the first author to carry out this application of Corollary 4.8 in a subsequent paper.

Section 5: Divisors and Units on Coverings of Configuration Spaces

In the present §5, we discuss a certain *generalization* [cf. Theorem 5.6; Remark 5.6.1], in the case of *proper hyperbolic curves*, of Theorem 4.7 [due to the second author]. Unlike the proof of Theorem 4.7 given in §4, the proof of this generalization does not rely on the notion of “goodness” or properties involving the “regular representation”. In this sense, the approach given in the present §5 is *more efficient* and relies on direct *algebraic-geometric properties* — such as the *disjointness of divisors* — of which the properties involving the “regular representation” applied in §4 may be thought of as a sort of “*étale-topological translation*”. On the other hand, the approach of §4 [which was discovered first, by the first author], though less efficient, is applicable to both *affine* and *proper hyperbolic curves*, and, moreover, has the virtue of relying on *explicit group-theoretic manifestations* of these algebraic-geometric properties; it was this explicitness that served to render the approach of §4 more readily accessible to the intuition of the first author. Finally, we discuss certain consequences [cf. Corollary 5.7] of the theory of the present §5 concerning the “*non-existence of units*” on finite étale coverings of a sufficiently generic hyperbolic curve.

We begin by reviewing some essentially well-known *generalities concerning log schemes*.

Definition 5.1. Let X^{\log} be a *fine log scheme* [cf. [Kato1]].

(i) Denote by M_X the étale sheaf of monoids on X that defines the log structure on X^{\log} . Thus, we have a natural injection $\mathcal{O}_X^\times \hookrightarrow M_X$, which we shall use to regard \mathcal{O}_X^\times as a subsheaf of M_X . We shall refer to the quotient sheaf of monoids

$$M_X^{\text{char}} \stackrel{\text{def}}{=} M_X / \mathcal{O}_X^\times$$

as the *characteristic* of X^{\log} and to the associated sheaf of groupifications

$$M_X^{\text{char-gp}}$$

as the *group-characteristic* of X^{\log} . Thus, [since X^{\log} is *fine*] the fibers of M_X^{char} (respectively, $M_X^{\text{char-gp}}$) are finitely generated torsion-free abelian monoids (respectively, abelian groups). For $n \in \mathbb{N}$, we shall denote by

$$U_X^{[n]} \subseteq X$$

and refer to as the n -interior of X^{\log} the subset [cf. Proposition 5.2, (i), (ii) below] of points [of the scheme X lying under geometric points of the scheme X] at which the fiber of $M_X^{\text{char-gp}}$ is of $\text{rank} \leq n$. Thus, $U_X^{[0]}$ is the interior $U_X \subseteq X$ of X^{\log} [i.e., the open subscheme of points at which the log structure of X^{\log} is *trivial*].

(ii) Let M be a finitely generated [abstract] abelian monoid; $N \geq 1$ an integer. We shall say that M is \mathbb{Q} -regular [with exponent N] if for some $n \in \mathbb{N}$, the map

$$\mathbb{N}^n \rightarrow \mathbb{N}^n$$

[where \mathbb{N}^n is the monoid determined by the product of n copies of \mathbb{N}] given by multiplication by N factors as a composite of injections of monoids $\mathbb{N}^n \hookrightarrow M \hookrightarrow \mathbb{N}^n$. We shall say that X^{\log} is *weakly \mathbb{Q} -regular* (respectively, *strongly \mathbb{Q} -regular*) if, for every geometric point \bar{x} of X , the fiber of $M_{\bar{x}}^{\text{char}}$ at \bar{x} is a \mathbb{Q} -regular monoid (respectively, \mathbb{Q} -regular monoid with exponent invertible in the residue field of \bar{x}).

Proposition 5.2. (Generalities on Log Schemes) *Let X^{\log} be a fine log scheme; $n \in \mathbb{N}$; l a prime number invertible on X . Then:*

(i) *The n -interior $U_X^{[n]} \subseteq X$ is open.*

(ii) *Suppose that X^{\log} is log regular. Then the complement of the n -interior $U_X^{[n]}$ is a closed subset of X of codimension $> n$; the complement $D_X \stackrel{\text{def}}{=} X \setminus U_X$ [equipped with the reduced induced scheme structure] is a divisor on X .*

(iii) *Suppose that X^{\log} is log regular and weakly \mathbb{Q} -regular. Then X is locally \mathbb{Q} -factorial [i.e., every Weil divisor on X admits a positive multiple which is Cartier]. Moreover, if X is connected, and G is any torsion-free profinite group, then any homomorphism of profinite groups*

$$\phi : \pi_1(U_X^{[1]}) \rightarrow G$$

factors through the natural surjection of étale fundamental groups $\pi_1(U_X^{[1]}) \rightarrow \pi_1(X)$.

(iv) *Suppose that X^{\log} is log smooth over a field k [equipped with the trivial log structure] and strongly \mathbb{Q} -regular; let $F \subseteq X$ be a closed subset of codimension $\geq n$. Then the natural map on étale cohomology*

$$H_{\text{ét}}^j(X, \mathbb{Q}_l) \rightarrow H_{\text{ét}}^j(X \setminus F, \mathbb{Q}_l)$$

is an isomorphism for $j \leq 2n - 2$ and an injection for $j = 2n - 1$.

(v) *Under the assumptions of (iv), suppose further that X is connected; write*

$$D_X = \bigcup_{i \in I} D_{X,i}$$

— where I is a finite set; the $D_{X,i} \subseteq X$ are **irreducible** divisors. [Thus, since X^{\log} is log regular, hence **normal** [cf. [Kato2], Theorem 4.1], we have a natural surjection of étale fundamental groups

$$\pi_1(U_X) \twoheadrightarrow \pi_1(X)$$

whose kernel contains the **inertia groups** of the $D_{X,i}$; the maximal pro- l quotient of each of these inertia groups is naturally isomorphic to some quotient of $\mathbb{Z}_l(1)$.] Then we have a **natural exact sequence**

$$\begin{aligned} 0 \rightarrow \mathrm{Hom}(\pi_1(X), \mathbb{Q}_l(1)) &\rightarrow \mathrm{Hom}(\pi_1(U_X), \mathbb{Q}_l(1)) \\ &\rightarrow \bigoplus_{i \in I} \mathbb{Q}_l \rightarrow H_{\text{ét}}^2(X, \mathbb{Q}_l(1)) \rightarrow H_{\text{ét}}^2(U_X, \mathbb{Q}_l(1)) \end{aligned}$$

— where the “Hom’s” denote the modules of continuous homomorphisms of topological groups; the second arrow is the arrow determined by the natural surjection $\pi_1(U_X) \twoheadrightarrow \pi_1(X)$; the third arrow is the arrow determined by the [copies of $\mathbb{Z}_l(1)$ that naturally surject onto the] **inertia groups** of the $D_{X,i}$ [and the natural identification of \mathbb{Q}_l with $\mathrm{Hom}(\mathbb{Z}_l(1), \mathbb{Q}_l(1))$]; the fourth arrow is the arrow that sends the $1 \in \mathbb{Q}_l$ in the direct summand labeled “ i ” to the **fundamental class** $c(D_{X,i})$ of the Weil divisor $D_{X,i}$ [which is well-defined, by (iii)].

Proof. Indeed, assertion (i) follows immediately from the definition of a *fine* log scheme [cf. [Kato1], §2.1-3]. In light of assertion (i), the portion of assertion (ii) concerning $U_X^{[n]}$ follows immediately from the inequality $\dim(\mathcal{O}_{X,\bar{x}}) \geq \mathrm{rank}_{\mathbb{Z}}((M_X^{\mathrm{char-gp}})_{\bar{x}})$ [where \bar{x} is a geometric point of X ; $\mathcal{O}_{X,\bar{x}}$ is the corresponding strict henselization of a local ring of X] — cf. the definition of “log regular” in [Kato2], Definition 2.1. Since X is *normal* [cf. [Kato2], Theorem 4.1], the portion of assertion (ii) concerning D_X follows immediately from the description given in [Kato2], Theorem 11.6, of the monoid M_X in terms of *rational functions* on X .

To verify assertion (iii) (respectively, (iv)), let us first observe that it follows immediately from our assumptions that X^{\log} is *log regular* (respectively, *log smooth* over k) and *weakly \mathbb{Q} -regular* (respectively, *strongly \mathbb{Q} -regular*) that every point of X admits an étale neighborhood $V \rightarrow X$ such that there exists a finite (respectively, [finite] Kummer log étale) dominant morphism $W^{\log} \rightarrow V^{\log}$ [where we equip V with the log structure pulled back from X] such that the scheme W is *regular* (respectively, *smooth* over k) and connected, and the log structure of W^{\log} arises from a *divisor with normal crossings* on W . Now the *local \mathbb{Q} -factoriality* portion of assertion (iii) follows immediately by pulling back a given Weil divisor on X to the *regular* scheme W [which yields a *Cartier* divisor on W] and then pushing forward via $W \rightarrow V$ [which multiplies the original divisor on V by the degree of the morphism $W \rightarrow V$]. To verify the portion of assertion (iii) concerning *étale fundamental groups*, we may assume without loss of generality that X , V , and W are *strictly henselian*, and that the morphism $V \rightarrow X$ is an *isomorphism*. Now by *Zariski-Nagata purity* [i.e., the classical non-logarithmic version of the “log purity theorem” quoted in §2], it follows that $\pi_1(U_W^{[1]}) \xrightarrow{\sim} \pi_1(W) = \{1\}$. On the other

hand, since the morphism $W \rightarrow X$ is *finite*, it follows that the natural morphism $\{1\} = \pi_1(U_W^{[1]}) \rightarrow \pi_1(U_X^{[1]})$ has *open image*, hence that $\pi_1(U_X^{[1]})$ is *finite*. Thus, our assumption that G is *torsion-free* implies that ϕ is *trivial*. This completes the proof of assertion (iii).

To verify assertion (iv), let us first observe that assertion (iv) holds when X is *smooth* over k . Indeed, in this case, by applying *noetherian induction* to F and possibly base-changing to a finite inseparable extension of k , we may assume without loss of generality that F is *smooth* over k ; but then the content of assertion (iv) is well-known [cf., e.g., [Milne], p. 244, Remark 5.4, (b)]. In the case of *arbitrary* X^{\log} , we argue as follows: Write $\iota : X_F \stackrel{\text{def}}{=} X \setminus F \hookrightarrow X$ for the natural inclusion. Then [by applying a well-known exact sequence in étale cohomology] it suffices to verify that $\mathbb{R}^j \iota_{\text{ét},*}(\mathbb{Q}_l) = 0$ for $0 < j \leq 2n - 1$. Since we have already verified assertion (iv) for k -smooth X , we may assume that $F \cap U_X = \emptyset$. In particular, it suffices [cf. [Milne], p. 88, Theorem 1.15] to verify, for an arbitrary *strictly henselization* \underline{V} of V at a closed point of V , that $H^j(\underline{V}_F, \mathbb{Q}_l) = 0$ for $0 < j \leq 2n - 1$ [where we write $\underline{V}_F \stackrel{\text{def}}{=} X_F \times_X \underline{V}$]. On the other hand, let us observe that since $\zeta^{\log} : \underline{W}_F^{\log} \stackrel{\text{def}}{=} \underline{V}_F \times_V W^{\log} \rightarrow \underline{V}_F^{\log} \stackrel{\text{def}}{=} \underline{V}_F \times_V V^{\log}$ is *Kummer log étale*, it follows that one may define a “*trace morphism*”

$$\zeta_{\text{ét},*}((\mathbb{Q}_l)_{\underline{W}_F}) \rightarrow (\mathbb{Q}_l)_{\underline{V}_F}$$

[where we use the subscripts “ \underline{W}_F ”, “ \underline{V}_F ” to denote the constant sheaf on \underline{W}_F , \underline{V}_F] that restricts, relative to $\zeta_{\text{ét}}^*$, to multiplication by the degree $\deg(\zeta)$ of ζ on $(\mathbb{Q}_l)_{\underline{V}_F}$. [Indeed, this is immediate for the restriction $\zeta_U : U_{\underline{W}_F} \rightarrow U_{\underline{V}_F}$ to the respective interiors, since this restriction is *finite étale*. On the other hand, since \underline{V}_F is *normal*, we have a *natural isomorphism* $(\mathbb{Q}_l)_{\underline{V}_F} \xrightarrow{\sim} \theta_* \theta^*((\mathbb{Q}_l)_{\underline{V}_F})$, where we write $\theta : U_{\underline{V}_F} \hookrightarrow \underline{V}_F$ for the natural inclusion of the interior. Thus, we obtain a trace morphism as desired by restricting to the interiors, applying the trace morphism on the interiors, and then applying this natural isomorphism.] Thus, by taking étale cohomology, one obtains a *trace morphism* $\tau : H^j(\underline{W}_F, \mathbb{Q}_l) \rightarrow H^j(\underline{V}_F, \mathbb{Q}_l)$ such that the composite $\tau \circ \rho$ with the restriction morphism $\rho : H^j(\underline{V}_F, \mathbb{Q}_l) \rightarrow H^j(\underline{W}_F, \mathbb{Q}_l)$ is equal to multiplication by $\deg(\zeta)$ on $H^j(\underline{V}_F, \mathbb{Q}_l)$. Since, moreover, we have already verified assertion (iv) for k -smooth X , it follows that $H^j(\underline{W}_F, \mathbb{Q}_l) = 0$ for $0 < j \leq 2n - 1$, hence that $H^j(\underline{V}_F, \mathbb{Q}_l) = 0$ for $0 < j \leq 2n - 1$, as desired.

Finally, we consider assertion (v). When $X = U_X^{[1]}$ [so X, D_X are *smooth* over k], assertion (v) follows immediately by applying the well-known *Gysin sequence* in étale cohomology [cf., e.g., [Milne], p. 244, Remark 5.4, (b)]

$$0 \rightarrow H_{\text{ét}}^1(X, \mathbb{Q}_l(1)) \rightarrow H_{\text{ét}}^1(U_X, \mathbb{Q}_l(1)) \rightarrow \bigoplus_{i \in I} \mathbb{Q}_l \rightarrow H_{\text{ét}}^2(X, \mathbb{Q}_l(1)) \rightarrow H_{\text{ét}}^2(U_X, \mathbb{Q}_l(1))$$

and the natural isomorphisms $H_{\text{ét}}^1((-), \mathbb{Q}_l(1)) \cong \text{Hom}(\pi_1((-), \mathbb{Q}_l(1)), \mathbb{Q}_l(1))$, for “ $(-)$ ” equal to X, U_X . For *arbitrary* X^{\log} , we reduce immediately to the case where $X = U_X^{[1]}$ by applying assertions (ii), (iv). \circ

Remark 5.2.1. We recall in passing that the *local \mathbb{Q} -factoriality* portion of Proposition 5.2, (iii), is *false* for arbitrary [not necessarily weakly \mathbb{Q} -regular] log regular X^{\log} . Indeed, such an example appears in the Remark following [Mzk1], Corollary 1.8.

Now we return to our discussion of *configuration spaces*. Let X be a *proper hyperbolic curve of genus g_X* over an *algebraically closed field k of characteristic zero*, $n \geq 1$ an integer, l a prime number; write $X_n \subseteq P_n$ for the associated n -th *configuration space*, Z_n^{\log} for the associated n -th *log configuration space*, and E for the *index set* of X_n , Z_n^{\log} [cf. Definition 2.1, (i)]. Thus, X_n may be identified with the *interior U_{Z_n}* of Z_n^{\log} .

Proposition 5.3. (The Logarithmic Geometry of the Log Configuration Space) *In the notation of the above discussion: Write*

$$V \stackrel{\text{def}}{=} U_{Z_n}^{[1]}; \quad V^{\log} \stackrel{\text{def}}{=} Z_n^{\log} \times_{Z_n} V$$

for the **1-interior** of Z_n^{\log} and the log scheme obtained by restricting the log structure of Z_n^{\log} . For $j \geq 1$ an integer, let us denote by $\wedge^j E$ the set of **subsets of E of cardinality j** [so E may be identified with $\wedge^1 E$] and by

$$\wedge^* E \stackrel{\text{def}}{=} \bigcup_{j=1}^n \wedge^j E$$

the [disjoint] union of the subsets of cardinality $j \geq 1$. Then:

(i) We shall refer to a divisor on Z_n obtained as the pull-back via a projection morphism $Z_n \rightarrow X$ of co-length 1 [and co-profile $e \in \wedge^1 E = E$] of a point $\in X(k)$ as a **fiber divisor [of co-profile e] [on Z_n]**. Then all fiber divisors of co-profile $e \in \wedge^1 E = E$ on Z_n determine the **same fundamental class**

$$\eta_e \in H_{\text{ét}}^2(Z_n, \mathbb{Q}_l(1))$$

— which we shall refer to as the **fiber class of co-profile e** [on Z_n].

(ii) The **irreducible divisors** on Z_n contained in the divisor D_{Z_n} defining the log structure of Z_n^{\log} are in **natural bijective correspondence** with the elements of $(\wedge^* E) \setminus E$. That is to say, a point of V belongs to the irreducible divisor $D_\varepsilon \subseteq V$ corresponding to an element $\varepsilon \in (\wedge^* E) \setminus E$ if and only if it corresponds to a stable curve with precisely two irreducible components, one isomorphic to X , the other of genus zero, such that the marked points that lie on X are precisely the marked points determined by the factors $e \in \varepsilon' \stackrel{\text{def}}{=} E \setminus \varepsilon$. In particular, we obtain a **natural isomorphism of schemes**

$$D_\varepsilon \cong X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$$

— where $|\varepsilon|$, $|\varepsilon'|$ are the cardinalities of ε , ε' , respectively; the projection $D_\varepsilon \rightarrow X_{|\varepsilon'|+1}$ is induced by any projection $X_n \rightarrow X_{|\varepsilon'|+1}$ of co-profile ε^+ , for $\varepsilon^+ \in \wedge^{|\varepsilon'|+1} E$ an element such that $\varepsilon' \subseteq \varepsilon^+$; $Q_{|\varepsilon|-2}$ is the $(|\varepsilon| - 2)$ -th configuration space [i.e., $\text{Spec}(k)$, when $|\varepsilon| = 2$] of “the” **tripod** [cf. §0] over k . In particular, the index set of the configuration space $X_{|\varepsilon'|+1}$ appearing in this isomorphism may be naturally identified with the set “ E/ε ” obtained from E by **identifying** the elements of ε to a single element $[\varepsilon] \in E/\varepsilon$. We shall refer to the irreducible divisor on Z_n contained in D_{Z_n} that corresponds to $\varepsilon \in (\wedge^* E) \setminus E$ as the **log-prime divisor of co-profile** ε [on Z_n]; we shall refer to the fundamental class

$$\eta_\varepsilon \in H_{\text{ét}}^2(Z_n, \mathbb{Q}_l(1))$$

of the log-prime divisor of co-profile ε as the **log-prime class of co-profile** ε [on Z_n].

(iii) Let

$$W^{\log} \rightarrow V^{\log}$$

be a connected [finite] **Kummer log étale covering**. Then W^{\log} is **log smooth** over k and **strongly \mathbb{Q} -regular**, and W , D_W are **smooth** over k . We shall also refer to irreducible divisors on W that lie over fiber divisors on Z_n as **fiber divisors** on W , and to fundamental classes of fiber divisors on W as **fiber classes** on W ; in a similar vein, we shall refer to irreducible divisors on W that lie over log-prime divisors on Z_n as **log-prime divisors** on W , and to fundamental classes of log-prime divisors on W as **log-prime classes** on W . Then a fiber class on W is completely determined by its **co-profile**. Also, we shall refer to a class on W or Z_n as a **log-characteristic class** if it is either a fiber class or a log-prime class.

Proof. Assertion (i) follows, for instance, from [Milne], p. 276, Theorem 11.1, (a). Assertion (ii) follows immediately from the definition of Z_n^{\log} involving the [log] moduli stack of stable curves. Assertion (iii) follows immediately from the definitions and assertion (i). \circ

Lemma 5.4. (Line Bundles on Log-prime Divisors) *In the notation of Proposition 5.3, (ii):*

(i) *The isomorphism $D_\varepsilon \cong X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$ of Proposition 5.3, (ii), determines an isomorphism of Picard groups $\text{Pic}(D_\varepsilon) \xrightarrow{\sim} \text{Pic}(X_{|\varepsilon'|+1})$.*

(ii) *The **co-normal bundle** of D_ε is isomorphic [cf. (i)] to the line bundle obtained by pulling back the **canonical bundle** ω_X of X via the [unique!] projection $X_{|\varepsilon'|+1} \rightarrow X$ that arises from a projection morphism $X_n \rightarrow X$ of co-length 1 whose co-profile is **not** contained in ε' .*

Proof. First, we consider assertion (i). Since $Q_{|\varepsilon|-2}$ is an open subscheme of the affine space [of dimension $|\varepsilon| - 2$] over k , it follows that $D_\varepsilon \cong X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$ is isomorphic to an open subscheme of the affine space [of dimension $|\varepsilon| - 2$] over

$X_{|\varepsilon'|+1}$. Thus, the assertion concerning Picard groups follows immediately from elementary algebraic geometry [cf., e.g., [Fulton], Theorem 3.3, (a)].

As for assertion (ii), we observe that the description of the stable curves parametrized by D_ε given in Proposition 5.3, (ii), implies [in light of the well-known *local structure* of a node] that the co-normal bundle in question is naturally isomorphic to the tensor product of the pull-back of the canonical bundle described in the statement of assertion (ii) with some [necessarily *trivial* — by assertion (i)] line bundle on D_ε pulled back from the natural projection to $Q_{|\varepsilon|-2}$. This completes the proof of assertion (ii). \circ

Lemma 5.5. (Linear Independence of Log-characteristic Classes) *In the notation of Proposition 5.3, (iii): Set*

$$I_W \stackrel{\text{def}}{=} I_W^{\text{fiber}} \cup I_W^{\text{log-prime}}$$

— where we write $I_W^{\text{fiber}} \stackrel{\text{def}}{=} E$ and $I_W^{\text{log-prime}}$ for the set of **log-prime divisors** on W . Thus, if we think of the elements of I_W^{fiber} as the co-profiles of fiber classes of $H_{\text{ét}}^2(W, \mathbb{Q}_l(1))$, then we obtain a [fiber or log-prime] class

$$\eta_i \in H_{\text{ét}}^2(W, \mathbb{Q}_l(1))$$

for each element $i \in I_W$. Then:

(i) The **log-characteristic classes** $\{\eta_i\}_{i \in I_W}$ are **linearly independent** over \mathbb{Q}_l .

(ii) The **restricted fiber classes** $\{\eta_i|_{U_W}\}_{i \in I_W^{\text{fiber}}}$ are **linearly independent** [in $H_{\text{ét}}^2(U_W, \mathbb{Q}_l(1))$] over \mathbb{Q}_l .

Proof. First, we observe that the description of the kernel of the restriction map $H_{\text{ét}}^2(W, \mathbb{Q}_l(1)) \rightarrow H_{\text{ét}}^2(U_W, \mathbb{Q}_l(1))$ given in the exact sequence of Proposition 5.2, (v), implies that assertion (ii) follows immediately from assertion (i). For $i \in I_W^{\text{log-prime}}$, write

$$D_i$$

for the divisor [tautologically!] determined by i . Thus, [by the definition of V^{log} , W^{log}]

$$D_i \cap D_j = \emptyset$$

for all $i, j \in I_W^{\text{log-prime}}$ such that $i \neq j$. Also, we observe that it follows immediately from the definitions that, for $i \in I_W^{\text{log-prime}}$, the covering $W^{\text{log}} \rightarrow V^{\text{log}}$ determines a *finite étale morphism*

$$D_i \rightarrow D_\varepsilon \cong X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$$

[cf. Proposition 5.3, (ii)]. Finally, we observe that we may assume without loss of generality that the Kummer log étale covering $W^{\text{log}} \rightarrow V^{\text{log}}$ is *Galois*, with Galois group $\Gamma \stackrel{\text{def}}{=} \text{Gal}(W^{\text{log}}/V^{\text{log}})$.

Now let us verify assertion (i) by *induction* on n . Let

$$\sum_{i \in I_W} c_i \cdot \eta_i = 0$$

[where the $c_i \in \mathbb{Q}_l$] be a linear relation among the η_i . The case $n = 1$ is immediate from the definitions [cf. also [Milne], p. 276, Theorem 11.1, (a)]. Next, we consider the case $n = 2$. Let $j \in I_W^{\log\text{-prime}}$. Since $n = 2$, it follows immediately from the definitions that D_j is a *proper hyperbolic curve* such that the covering $W^{\log} \rightarrow V^{\log}$ induces a finite étale morphism of D_j onto the diagonal of $V = Z_n = X \times X$. In particular, it follows that we may restrict the above linear relation to D_j to obtain a linear relation

$$-c_j \cdot \eta_j|_{D_j} = \sum_{i \in I_W^{\text{fiber}}} c_i \cdot \eta_i|_{D_j}$$

among classes of $H_{\text{ét}}^2(D_j, \mathbb{Q}_l(1)) \cong \mathbb{Q}_l$ [cf. [Milne], p. 276, Theorem 11.1, (a)]. Moreover, since the class $\eta_j|_{D_j}$ is the pull-back to D_j of some \mathbb{Q}_l^\times -multiple of the fundamental class of the diagonal of $X \times X$ [cf. Lemma 5.4, (ii)], we thus conclude that $\eta_j|_{D_j} \neq 0$. Next, let us observe that, for $i \in I_W^{\text{fiber}}$, the classes η_i are *fixed* by the natural action of Γ on $H_{\text{ét}}^2(W, \mathbb{Q}_l(1))$. Thus, if we identify $H_{\text{ét}}^2(D_j, \mathbb{Q}_l(1))$ with \mathbb{Q}_l via the natural isomorphism $H_{\text{ét}}^2(D_j, \mathbb{Q}_l(1)) \cong \mathbb{Q}_l$, then we conclude that the elements $\eta_i|_{D_j}, \eta_j|_{D_j} \in \mathbb{Q}_l$ [where $i \in I_W^{\text{fiber}}$] are *independent* of the choice of j among all Γ -conjugates of j , hence that the element $c_j \in \mathbb{Q}_l$ is *independent* of the choice of j among all Γ -conjugates of j . But this implies that the linear relation $\sum_{i \in I_W} c_i \cdot \eta_i = 0$ arises as the pull-back to W of a similar linear relation on V . That is to say, we may assume without loss of generality that $W^{\log} = V^{\log}$. But then the \mathbb{Q}_l -linear independence of the unique log-prime class η_Δ and the two fiber classes η_1, η_2 follows, for instance, from the [easily verified] *non-singularity of the matrix of intersection numbers* among the classes $\eta_\Delta, \eta_1, \eta_2$ [where we recall that $\eta_\Delta \cdot \eta_1 = \eta_\Delta \cdot \eta_2 = \eta_1 \cdot \eta_2 = 1, \eta_1 \cdot \eta_1 = \eta_2 \cdot \eta_2 = 0, \eta_\Delta \cdot \eta_\Delta = 2 - 2g_X$]. This completes the proof of the case $n = 2$.

Now we assume that $n \geq 3$. Let D_j [where $j \in I_W^{\log\text{-prime}}$] be a *log-prime divisor of co-length 2* — i.e., which projects to a log-prime divisor of V whose *co-profile* ε is of cardinality 2. Then the covering $W^{\log} \rightarrow V^{\log}$ determines a finite étale morphism $D_j \rightarrow X_{n-1}$; that is to say, D_j appears as a “ U_W ” that arises in the case “ $n - 1$ ”. Moreover, it follows immediately from Lemma 5.4, (ii), that $\eta_j|_{D_j}$ is a \mathbb{Q}_l -multiple of the restriction to D_j of the *fiber class* of Z_{n-1}^{\log} of co-profile $[\varepsilon] \in E/\varepsilon$ [cf. Proposition 5.3, (ii)]. On the other hand, for $i \in I_W^{\log\text{-prime}}$ such that $i \neq j$, $\eta_i|_{D_j} = 0$; for $i \in I_W^{\text{fiber}}$ of co-profile $e \in E$, $\eta_i|_{D_j}$ is a \mathbb{Q}_l^\times -multiple of the restriction to D_j of the *fiber class* of Z_{n-1}^{\log} whose co-profile is the image of e in E/ε . In particular, by restricting the linear relation $\sum_{i \in I_W} c_i \cdot \eta_i = 0$ to D_j , it follows by applying assertion (ii) for “ $n - 1$ ” [i.e., here we apply the *induction hypothesis* on n] that $c_i = 0$ for all $i \in \varepsilon' \subseteq E = I_W^{\text{fiber}}$. Since $n \geq 3$, it follows that $\varepsilon' \neq \emptyset$. Thus, by *varying* j [i.e., varying ε], we conclude that $c_i = 0$ for all $i \in I_W^{\text{fiber}}$.

Now to complete the proof of Lemma 5.5, it suffices to show that the coefficients of any linear relation $\sum_{i \in I_W^{\log\text{-prime}}} c_i \cdot \eta_i = 0$ *vanish*. On the other hand,

by restricting to D_j , for $j \in I_W^{\log\text{-prime}}$, we obtain relations $c_j \cdot \eta_j|_{D_j} = 0$ for each $j \in I_W^{\log\text{-prime}}$. Thus, to complete the proof of Lemma 5.5, it suffices to show that

$$\eta_j|_{D_j} \neq 0$$

for $j \in I_W^{\log\text{-prime}}$. On the other hand, by Lemma 5.4, (ii), it follows that $\eta_j|_{D_j}$ is a \mathbb{Q}_l^\times -multiple of the pull-back to D_j of a class on $X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$ [for some $\varepsilon \in (\wedge^* E) \setminus E$, $\varepsilon' \stackrel{\text{def}}{=} E \setminus \varepsilon$] that arises as the pull-back to $X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2}$ via the projection morphism $X_{|\varepsilon'|+1} \times Q_{|\varepsilon|-2} \rightarrow X_{|\varepsilon'|+1}$ of the restriction to $X_{|\varepsilon'|+1}$ of a fiber class of $Z_{|\varepsilon'|+1}$. Since $|\varepsilon'| + 1 \leq n - 1$, it thus follows from assertion (ii) for “ $|\varepsilon'| + 1$ ” [i.e., here we apply the *induction hypothesis* on n] that the class $\eta_j|_{D_j} \neq 0$, as desired. This completes the proof of Lemma 5.5. \circ

Theorem 5.6. (Extendability of Coverings) *Let X be a proper hyperbolic curve over an algebraically closed field k of characteristic zero; $n \geq 1$ an integer; $X_n \subseteq P_n$ the n -th configuration space associated to X ; $(X_n^*)^{\log} \stackrel{\text{def}}{=} Z_n^{\log}$ the n -th log configuration space associated to X ;*

$$Y \rightarrow X_n$$

a finite étale morphism, where Y is connected; $Y^ \rightarrow X_n^*$ the normalization of X_n^* in Y ; G a strongly torsion-free profinite group;*

$$\phi : \pi_1(Y) \rightarrow G$$

a continuous homomorphism. Then:

(i) *The homomorphism ϕ factors uniquely through the natural surjection $\pi_1(Y) \twoheadrightarrow \pi_1(Y^*)$ induced by the open immersion $Y \hookrightarrow Y^*$.*

(ii) *Suppose further that the covering $Y \rightarrow X_n$ arises from a product-theoretic open subgroup of $\pi_1(X_n)$. Then the kernel of ϕ is product-theoretic.*

Proof. First, we consider assertion (i). In the notation of the discussion preceding Theorem 5.6: By the *log purity theorem* [cf. the discussion of §2], we have a natural isomorphism $\pi_1(U_V) \xrightarrow{\sim} \pi_1(V^{\log})$, where we observe that [it follows immediately from the definitions that] the interior U_V of V^{\log} may be identified with X_n . Thus, the finite étale covering $Y \rightarrow X_n$ may be identified with the interior U_W of a Kummer log étale covering $W^{\log} \rightarrow V^{\log}$. In particular, by the *log purity theorem*, we conclude that the finite morphism $Y^* \rightarrow X_n^*$ determines a Kummer log étale covering $(Y^*)^{\log} \rightarrow (X_n^*)^{\log}$. Thus, in summary, we obtain a commutative diagram

$$\begin{array}{ccccc} U_W = Y & \hookrightarrow & W^{\log} & \hookrightarrow & (Y^*)^{\log} \\ \downarrow & & \downarrow & & \downarrow \\ U_V = X_n & \hookrightarrow & V^{\log} & \hookrightarrow & (X_n^*)^{\log} \end{array}$$

in which the “hooked horizontal arrows” are open immersions; the complements of the open immersions $W \hookrightarrow Y^*$, $V \hookrightarrow X_n^*$ are of *codimension* ≥ 2 ; the vertical arrows are Kummer log étale coverings; by abuse of notation, we use notation for schemes to denote the corresponding log schemes with trivial log structure. In particular, it follows immediately from the fact that the log structure of $(X_n^*)^{\log}$ arises from a divisor with normal crossings that $(Y^*)^{\log}$ is *log smooth* over k and *strongly \mathbb{Q} -regular*. Thus, by Proposition 5.2, (iii) [which is applicable since $(Y^*)^{\log}$ is *log regular* and *weakly \mathbb{Q} -regular*], it follows that, to complete the proof of assertion (i), it suffices to show that ϕ factors through the natural surjection $\pi_1(Y) \twoheadrightarrow \pi_1(W)$.

Next, we *claim* that we may assume without loss of generality that G is *abelian*. Indeed, to show that ϕ factors through the natural surjection $\pi_1(Y) \twoheadrightarrow \pi_1(W)$, it suffices [by *Zariski-Nagata purity*, since W , D_W are *smooth* over k] to show that $\phi(\xi) = 1$ for *each element* $\xi \in \pi_1(Y)$ of the inertia group of an irreducible divisor of $D_W = W \setminus Y$. Let $\Xi \subseteq \pi_1(Y)$ be the closed subgroup *topologically generated* by ξ ; suppose that $\phi(\Xi) \neq \{1\}$. Then [cf. the argument applied at the beginning of the proof of Theorem 1.5] there exists an open subgroup $G_\xi \subseteq G$ containing $\phi(\Xi)$ such that $\phi(\Xi)$ [i.e., Ξ] has *nontrivial image* in G_ξ^{ab} . On the other hand, by applying assertion (i) to the situation where we *replace* Y by the finite étale covering $Y_\xi \rightarrow Y$ determined [via ϕ] by the open subgroup $G_\xi \subseteq G$, *replace* G by G_ξ^{ab} , and *replace* W by the *normalization* W_ξ of W in Y_ξ [where we note that $\xi \in \pi_1(Y_\xi)$ still satisfies the property of being an element of an inertia group of an irreducible divisor of $W_\xi \setminus Y_\xi$], we conclude that Ξ has *trivial image* in G_ξ^{ab} , a *contradiction*. This completes the proof of the *claim*.

Thus, [cf. the proof of Theorem 4.7] it suffices to verify assertion (i) in the case where $G = \mathbb{Z}_l$, for some prime number l . But in this case, the fact that the given homomorphism $\pi_1(Y) = \pi_1(U_W) \rightarrow G$ factors through $\pi_1(U_W) \twoheadrightarrow \pi_1(W)$ follows immediately from the *exact sequence* of Proposition 5.2, (v), in light of the \mathbb{Q}_l -*linear independence* asserted in Lemma 5.5, (i). This completes the proof of assertion (i).

Next, we consider assertion (ii). Write

$$Y^\dagger \rightarrow P_n$$

for the normalization of P_n ($\supseteq X_n$) in Y . Since Y^* is *normal* and maps to X_n^* , hence to P_n , we thus obtain a *birational morphism* $Y^* \rightarrow Y^\dagger$. Since the morphism $X_n^* \rightarrow P_n$ is *proper*, it thus follows that the morphism $Y^* \rightarrow Y^\dagger$ is *proper*. On the other hand, since $Y \rightarrow X_n$ arises from a *product-theoretic* open subgroup of $\pi_1(X_n)$, one verifies immediately that Y^\dagger is *smooth* over k , and that the kernel of the natural surjection $\pi_1(Y) \rightarrow \pi_1(Y^\dagger)$ is *product-theoretic*. In particular, by *Zariski-Nagata purity* [i.e., the classical non-logarithmic version of the “log purity theorem” quoted above], we conclude that the natural surjection $\pi_1(Y^*) \twoheadrightarrow \pi_1(Y^\dagger)$ is an *isomorphism*. But this implies that the given homomorphism $\pi_1(Y) \rightarrow G$ *factors* through the natural surjection $\pi_1(Y) \twoheadrightarrow \pi_1(Y^\dagger)$, hence, in particular, is *product-theoretic*. This completes the proof of assertion (ii). \circ

Remark 5.6.1. Theorem 5.6, (ii), *generalizes* Theorem 4.7, in the case of *proper hyperbolic curves*, to the case where “ G ” is *not necessarily pro-solvable*. This is

possible precisely because, unlike the theory of §4, the theory of the present §5 is applicable to coverings of configuration spaces that are *not necessarily product-theoretic* [cf. the reduction in the proof of Theorem 5.6, (i), to the case where G is *abelian*].

Remark 5.6.2. In an earlier version of the present manuscript, Theorem 5.6 was stated in a form that applied to *affine hyperbolic curves of genus ≥ 2* as well. In fact, however, the proof given for affine curves was found to contain a *gap* which the authors were unable to repair. Nevertheless, although such a gap exists for the analogue of Theorem 5.6, (i), for affine hyperbolic curves, this gap does not appear when the covering $Y \rightarrow X_n$ of Theorem 5.6 is assumed to be *product-theoretic* [i.e., the situation of Theorem 5.6, (ii)], under the further assumption that G be *pro-solvable* [cf. Remark 5.6.1]. In particular, it is a routine task to modify the argument given in the present §5 so as to yield a proof of Theorem 5.6, (ii) [under the further assumption that G be *pro-solvable*] — i.e., a *new proof* of Theorem 4.7 — for *arbitrary hyperbolic curves of genus ≥ 2* .

Remark 5.6.3. It is not clear to the authors at the time of writing whether or not one should expect the analogue of Theorem 5.6 to hold for affine hyperbolic curves. Nevertheless, one way to think about the “*extendability of coverings*” property discussed in Theorem 5.6 is the following: This property holds [by Theorem 5.6] for the “*tautological affine curve*” with a given fixed hyperbolic compactification — i.e., the tautological affine curve over a *configuration space* associated to a proper hyperbolic curve. That is to say, it may be the case that this sort of special property *only* holds for such affine curves with “*universal/tautological cusps*” and *not* for affine curves whose cusps are “*non-tautological*”.

Corollary 5.7. (Non-existence of Generic Units) *In the notation of Theorem 5.6, let us fix a projection morphism*

$$\phi_X : X_n \rightarrow B \stackrel{\text{def}}{=} X_{n-1}$$

*of length 1, which allows us to regard X_n as a family of hyperbolic curves over B . Denote the [scheme determined by the] generic point of B by η ; write $k(\eta)$ for the residue field of η , $\mathcal{X} \stackrel{\text{def}}{=} X_n \times_B \eta$. Let $k(\eta')$ be an arbitrary field extension of $k(\eta)$, $\eta' \rightarrow \eta$ the resulting morphism of schemes, \mathcal{Y} a **hyperbolic curve** over η' , and*

$$\mathcal{Y} \rightarrow \mathcal{X}_{\eta'} \stackrel{\text{def}}{=} \mathcal{X} \times_{\eta} \eta' = X_n \times_B \eta'$$

a finite étale covering over η' . Then every unit $u \in \Gamma(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}^{\times})$ on \mathcal{Y} is constant, i.e., is contained in the image of $k(\eta')$ in $\mathcal{O}_{\mathcal{Y}}$.

Proof. One reduces immediately by well-known elementary algebraic geometry arguments [i.e., replacing $k(\eta')$ by a finitely generated field extension of $k(\eta)$, extending $\mathcal{Y} \rightarrow \mathcal{X}_{\eta'}$ over some variety that admits $k(\eta')$ as its function field, and

restricting to a closed point of this variety] to the case where the morphism $\eta' \rightarrow \eta$ is *finite étale*. Now let us observe that by the *exact sequence* of Proposition 2.2, (i), it follows that, after possibly *replacing* \mathcal{Y} by an appropriate connected finite étale covering of \mathcal{Y} , we may assume without loss of generality that there exists a *commutative diagram*

$$\begin{array}{ccc} Y & \longrightarrow & X_n \\ \downarrow & & \downarrow \\ C & \longrightarrow & B \end{array}$$

in which the horizontal arrows are *connected finite étale coverings*; the vertical arrows are *families of hyperbolic curves*; the *divisor of cusps* $D_Y \subseteq \overline{Y}$ in the [relative, i.e., over C] *compactification* $\overline{Y} \rightarrow C$ of $Y \rightarrow C$ forms a *split* finite étale covering of C ; the covering $\mathcal{Y} \rightarrow \mathcal{X}$ *factors* through $Y_{\eta_C} \stackrel{\text{def}}{=} Y \times_C \eta_C$ [where $\eta_C \stackrel{\text{def}}{=} C \times_B \eta$] in such a way that the induced covering

$$\mathcal{Y} \rightarrow Y_{\eta_C}$$

is obtained by base-changing the curve Y_{η_C} over η_C via a morphism $\eta' \rightarrow \eta_C$. In particular, since the divisor of zeroes and poles of u on the compactification of the hyperbolic curve \mathcal{Y} descends [by our “*splitness*” assumption on D_Y] to a divisor on the compactification of the hyperbolic curve Y_{η_C} , it follows immediately that u is *constant* if and only if its *norm* relative to this covering $\mathcal{Y} \rightarrow Y_{\eta_C}$ [which forms a unit on Y_{η_C}] is *constant*. Thus, in summary, we may assume without loss of generality that $\mathcal{Y} = Y_{\eta_C}$, and that u is a *unit* on Y_{η_C} .

Now since u may be regarded as a *rational function* on \overline{Y} , the divisor of zeroes and poles of this rational function on \overline{Y} may be written in the form

$$D^{\text{cusp}} + D^{\text{base}}$$

— where D^{cusp} is a divisor supported on D_Y ; D^{base} is a divisor on \overline{Y} that arises as the pull-back to \overline{Y} of a divisor D_C^{base} on C . In particular, we obtain a relation

$$0 = c(D^{\text{cusp}}) + c(D^{\text{base}}) \in H_{\text{ét}}^2(\overline{Y}, \mathbb{Q}_l(1))$$

— where l is a prime number; we write $c(-)$ for the *Chern class* of the line bundle determined by a divisor on a k -smooth scheme [such as Y, \overline{Y}, C]. Next, let us pull-back this relation via a section $s : C \rightarrow \overline{Y}$ whose image D_s is contained in D_Y . This yields a relation

$$\lambda \cdot s^*(c(D_s)) = s^*(c(D^{\text{base}})) = c(D_C^{\text{base}}) \in H_{\text{ét}}^2(C, \mathbb{Q}_l(1))$$

for some $\lambda \in \mathbb{Q}_l$. On the other hand, let us observe that, relative to the notation of the discussion preceding Theorem 5.6, if we take $U_W \rightarrow U_V$ to be the covering $Y \rightarrow X_n$, then \overline{Y} may be regarded as an *open subscheme* of W , and $D_s \subseteq \overline{Y}$ as a *log-prime divisor* of W . From this point of view, it follows from Lemma 5.4, (ii), that $s^*(c(D_s))$ is a \mathbb{Q}_l -multiple of the pull-back to C via $C \rightarrow B = X_{n-1}$ of a *fiber*

class on $B = X_{n-1}$. In particular, we conclude that $c(D_C^{\text{base}})$ is a \mathbb{Q}_l -multiple of the pull-back to C of a fiber class on $B = X_{n-1}$, hence that $c(D^{\text{base}}) \in H_{\text{ét}}^2(\overline{Y}, \mathbb{Q}_l(1))$ is a \mathbb{Q}_l -multiple of the pull-back to $\overline{Y} \subseteq W$ of a fiber class on W .

Thus, in summary, the relation $0 = c(D^{\text{cusp}}) + c(D^{\text{base}})$ in $H_{\text{ét}}^2(\overline{Y}, \mathbb{Q}_l(1))$ constitutes a \mathbb{Q}_l -linear relation between certain log-prime classes [i.e., $c(D^{\text{cusp}})$] and certain fiber classes [i.e., $c(D^{\text{base}})$] on \overline{Y} . Since [it follows immediately from the definitions that] the complement of \overline{Y} in W is a [disjoint!] union of certain log-prime divisors on W , the description of the kernel of the restriction map $H_{\text{ét}}^2(W, \mathbb{Q}_l(1)) \rightarrow H_{\text{ét}}^2(\overline{Y}, \mathbb{Q}_l(1))$ given in the exact sequence of Proposition 5.2, (v), implies that this \mathbb{Q}_l -linear relation $0 = c(D^{\text{cusp}}) + c(D^{\text{base}})$ in $H_{\text{ét}}^2(\overline{Y}, \mathbb{Q}_l(1))$ arises from some \mathbb{Q}_l -linear relation in $H_{\text{ét}}^2(W, \mathbb{Q}_l(1))$ obtained by appending some \mathbb{Q}_l -linear combination of the log-prime classes arising from the log-prime divisors in the complement $W \setminus \overline{Y}$. On the other hand, by the \mathbb{Q}_l -linear independence asserted in Lemma 5.5, (i), the coefficients of such a \mathbb{Q}_l -linear relation necessarily vanish. In particular, since D^{cusp} is a \mathbb{Z} -linear combination of log-prime divisors of W that lie in \overline{Y} , we thus conclude that the coefficients $\in \mathbb{Z}$ of this \mathbb{Z} -linear combination vanish, i.e., that $D^{\text{cusp}} = 0$. But this amounts precisely to the assertion that the unit u is constant. This completes the proof of Corollary 5.7. \circ

Section 6: Nearly Abelian Groups

In the present §6, we discuss another approach, based on the notion of a “nearly abelian” profinite group, to verifying the group-theoreticity of the various fiber subgroups associated to a configuration space group.

Definition 6.1. We shall say that a profinite group G is *nearly abelian* if it admits a normal closed subgroup $N \subseteq G$ which is *topologically normally generated* by a single element $\in G$ such that G/N is almost abelian.

Proposition 6.2. (Nearly Abelian Surface Groups) *Let \mathcal{C} be a nontrivial full formation. Then a pro- \mathcal{C} surface group Π is nearly abelian if and only if it is a free pro- \mathcal{C} group on two generators — i.e., it arises from a hyperbolic curve which is either of type $(0, 3)$ or of type $(1, 1)$ [cf. Remark 1.2.2].*

Proof. Since the sufficiency of the condition given in the statement of Proposition 6.2 is immediate from the definitions, it suffices to verify the necessity of this condition. Thus, we suppose that Π is *nearly abelian*. Now observe that for any $l \in \Sigma_{\mathcal{C}}$, the maximal pro- l quotient of Π is again a *nearly abelian [pro- l] surface group*. Thus, we may assume without loss of generality that \mathcal{C} is *primary*. Write $\Sigma_{\mathcal{C}} = \{l\}$; $\overline{\Pi} \stackrel{\text{def}}{=} \Pi/[\Pi, [\Pi, \Pi]]$; $\Pi_1 \stackrel{\text{def}}{=} \overline{\Pi}^{\text{ab}} \cong \Pi^{\text{ab}}$; $\Pi_2 \stackrel{\text{def}}{=} [\overline{\Pi}, \overline{\Pi}] \subseteq \overline{\Pi}$. Thus, we have an exact sequence

$$1 \rightarrow \Pi_2 \rightarrow \overline{\Pi} \rightarrow \Pi_1 \rightarrow 1$$

and a surjection $\wedge^2 \Pi_1 \rightarrow \Pi_2$. Here, we regard Π_1, Π_2 as *finitely generated free \mathbb{Z}_l -modules*. Note [cf. Remark 1.2.2] that if Π_1 is of rank d , then Π_2 is of rank $\frac{1}{2}d(d-1) - \epsilon$, where $\epsilon = 0$ if Π is *open*, and $\epsilon = 1$ if Π is *closed*. Also, we observe that $d \geq 2$ if $\epsilon = 0$, and $d \geq 4$ if $\epsilon = 1$ [cf. Remark 1.2.2]. Thus, to complete the proof of Proposition 6.2, it suffices to show that $d = 2$.

Next, let us observe that $\overline{\Pi}$ is also *nearly abelian*. Thus, there exists an element $\gamma \in \overline{\Pi}$ such that if we write $N \subseteq \overline{\Pi}$ for the subgroup topologically normally generated by γ , then $\overline{\Pi}/N$ is *almost abelian*. In particular, [since the commutator subgroup of *any* open subgroup of $\overline{\Pi}$ forms an *open subgroup* of Π_2] it follows immediately that $N \cap \Pi_2 \subseteq \overline{\Pi}$ forms an *open subgroup* of Π_2 — i.e., a \mathbb{Z}_l -module of the same rank as Π_2 . If $\gamma \in \Pi_2$, then $N \cap \Pi_2 = N$ is of rank ≤ 1 , so we obtain that $\frac{1}{2}d(d-1) - \epsilon \leq 1$, i.e., $d(d-1) \leq 2(1+\epsilon) \leq 4$, so $d \leq 2$, i.e., $d = 2$, as desired. Thus, it remains to consider the case where $\gamma \notin \Pi_2$. In this case, one verifies immediately that $N \cap \Pi_2 \subseteq \Pi_2$ is given by the image of the morphism

$$[\gamma, -] : \Pi_1 \rightarrow \Pi_2$$

given by forming the *commutator* with γ . Since this morphism clearly vanishes on γ , its image is of rank $\leq d-1$. Thus, we conclude that

$$d-1 \geq \frac{1}{2}d(d-1) - \epsilon$$

— i.e., that $2 \cdot \epsilon \geq (d-1)(d-2)$, which implies that $d \leq 2$ if $\epsilon = 0$, and $d \leq 3$ if $\epsilon = 1$. But this is enough to conclude that $d = 2$ [and $\epsilon = 0$]. This completes the proof of Proposition 6.2. \circ

Corollary 6.3. (Group-theoreticity of Projections of Configuration Spaces II) *Let \mathcal{C} be a PT-formation. For $\xi = \alpha, \beta$, let X^ξ be a hyperbolic curve whose type is neither $(0, 3)$ nor $(1, 1)$ over an algebraically closed field k_ξ of characteristic zero; $n_\xi \geq 1$ an integer; $X_{n_\xi}^\xi$ the n_ξ -th configuration space associated to X^ξ ; $\Pi^\xi \stackrel{\text{def}}{=} \pi_1^{\mathcal{C}}(X_{n_\xi}^\xi)$; E_ξ the index set of $X_{n_\xi}^\xi$. Let*

$$\gamma : \Pi^\alpha \xrightarrow{\sim} \Pi^\beta$$

be an isomorphism of profinite groups. Then γ induces a bijection $\sigma : E_\alpha \xrightarrow{\sim} E_\beta$ [so $n_\alpha = n_\beta$] such that

$$\gamma(F_\alpha) = F_\beta$$

for all fiber subgroups $F_\alpha \subseteq \Pi^\alpha$, $F_\beta \subseteq \Pi^\beta$, whose respective profiles $E'_\alpha \subseteq E_\alpha$, $E'_\beta \subseteq E_\beta$ correspond via σ .

Proof. Just as in the proof of Corollary 4.8, to complete the proof of Corollary 6.3, it suffices to verify that the image via γ of any fiber subgroup of Π^α of co-length

one is contained in a fiber subgroup of Π^β of co-length one. For $j = 1, \dots, n_\xi$, let us write

$$K_j^\xi \subseteq \Pi_\xi$$

for the *fiber subgroup* $\subseteq \Pi^\xi$ of *co-length one* with co-profile given by the element of E_ξ labeled by j , and

$$J_j^\xi \subseteq \Pi_\xi$$

for the *fiber subgroup* $\subseteq \Pi^\xi$ of *length one* with profile given by the element of E_ξ labeled by j . Thus, [cf. Proposition 2.4, (vi)] to complete the proof of Corollary 6.3, it suffices to verify the following statement [in general]:

For each $i \in E_\alpha$, there exists a $j \in E_\beta$ such that $J_{j'}^\beta \subseteq \gamma(K_i^\alpha)$ for all $j' \in E_\beta$ such that $j' \neq j$.

To verify this statement, we reason as follows: Write

$$\phi : \Pi^\beta \xrightarrow{\sim} \Pi^\alpha \twoheadrightarrow G \stackrel{\text{def}}{=} \Pi^\alpha / K_i^\alpha$$

for the surjection determined by γ^{-1} . Then it suffices to show that there do not exist two distinct elements $j_1, j_2 \in E_\beta$ such that $J_{j_1}^\beta, J_{j_2}^\beta$ have *nontrivial image* under ϕ . Thus, let us suppose that the images J_1, J_2 of $J_{j_1}^\beta, J_{j_2}^\beta$ under ϕ are *nontrivial*. Since $J_{j_1}^\beta, J_{j_2}^\beta$ are *topologically finitely generated normal closed subgroups* of Π^β , it follows from Theorem 1.5 that J_1, J_2 are *open* in G [cf. Remark 3.3.2]. Moreover, by Proposition 2.4, (v), it follows that there exists a normal closed subgroup $N \subseteq G$ that is topologically normally generated by a *single element* such that the images of J_1, J_2 in G/N *commute*. Thus, G/N contains an *abelian open subgroup*, i.e., is *almost abelian*. On the other hand, since $K_i^\alpha \subseteq \Pi^\alpha$ is a fiber subgroup of co-length one, it follows that $G = \Pi^\alpha / K_i^\alpha$ is a *surface group*. Thus, in summary, we conclude that G is a *nearly abelian surface group*, which, by Proposition 6.2, contradicts our hypothesis concerning the *type of the hyperbolic curve* X^α . This completes the proof of Corollary 6.3. \circ

Remark 6.3.1. Unlike the case with Corollary 4.8, it seems *unrealistic* at the time of writing to extend the technique of the proof of Corollary 6.3 to the case of *arbitrary product-theoretic open subgroups* $\subseteq \pi_1^{\mathcal{C}}(X_{n_\xi}^\xi)$ [cf. Corollary 4.8], since this would require an analogue of Proposition 6.2 for surface groups that become almost abelian after forming the quotient by a subgroup topologically normally generated by a *very large* number of elements [roughly, on the order of the *index* of the product-theoretic open subgroups under consideration].

Section 7: A Discrete Analogue

In the present §7, we discuss various consequences of Theorems 4.7, 5.6 and Corollaries 4.8, 6.3 [cf. Corollaries 7.3, 7.4 below] for the *topological fundamental groups* of configuration spaces over the complex number field \mathbb{C} .

In the following discussion, if Z is a connected scheme of finite type over \mathbb{C} , then we shall use the notation

$$\pi_1^{\text{top}}(Z)$$

to denote the “*topological fundamental group*” [i.e., the fundamental group in the usual sense of algebraic topology], for some choice of basepoint, of the topological space of \mathbb{C} -rational points $Z(\mathbb{C})$ [equipped with the topology determined by the topology of \mathbb{C}].

Let X be a *hyperbolic curve* over \mathbb{C} , $n \geq 1$ an integer. Write $X_n \subseteq P_n$ for the n -th *configuration space* associated to X [cf. the notation of Definition 2.1, (i)]. Since the complement in a connected complex manifold of any submanifold of [complex] codimension ≥ 1 is clearly connected, it thus follows that the inclusion $X_n \hookrightarrow P_n$ induces a *natural surjection* $\pi_1^{\text{top}}(X_n) \twoheadrightarrow \pi_1^{\text{top}}(P_n)$.

The following “discrete analogue” of [a certain portion of] Proposition 2.2 is well-known:

Proposition 7.1. (Topological Fundamental Groups of Configuration Spaces) *In the notation of the above discussion:*

(i) *Any projection morphism $X_n \rightarrow X_{n-1}$ of length one determines a **natural exact sequence***

$$1 \rightarrow \pi_1^{\text{top}}((X_n)_{\bar{x}}) \rightarrow \pi_1^{\text{top}}(X_n) \rightarrow \pi_1^{\text{top}}(X_{n-1}) \rightarrow 1$$

[where we write $X_0 \stackrel{\text{def}}{=} \text{Spec}(\mathbb{C})$; \bar{x} is a \mathbb{C} -valued geometric point of X_{n-1}].

(ii) *The natural morphism*

$$\pi_1^{\text{top}}(X_n) \rightarrow \pi_1(X_n)$$

*to the étale fundamental group $\pi_1(X_n)$ is **injective**, i.e., $\pi_1^{\text{top}}(X_n)$ is **residually finite**.*

Proof. Assertion (i) is discussed, for instance, in [Birm], Theorem 1.4. To verify assertion (ii), observe that [by induction on n] it follows from the exact sequences of assertion (i) and the analogue of assertion (i) for $\pi_1(X_n)$ [cf. Proposition 2.2, (i)], that we may assume without loss of generality that $n = 1$. Now let us recall that $\pi_1^{\text{top}}(X)$ may be embedded [by considering the well-known *uniformization* of $X(\mathbb{C})$ by the *upper half-plane*] into $SL_2(\mathbb{R})/\{\pm 1\}$. That is to say, $\pi_1^{\text{top}}(X)$ is a “*finitely*

generated linear group”, so the desired *residual-finiteness* follows from a well-known theorem of Mal’cev [cf., e.g., [Wehr], Theorem 4.2]. \circ

Definition 7.2.

(i) We shall refer to a subgroup $H \subseteq \pi_1^{\text{top}}(X_n)$ as being *product-theoretic* if H arises as the inverse image via the natural surjection $\pi_1^{\text{top}}(X_n) \twoheadrightarrow \pi_1^{\text{top}}(P_n)$ of a subgroup of $\pi_1^{\text{top}}(P_n)$.

(ii) Write E for the *index set* of X_n . Let $E' \subseteq E$ be a subset of cardinality n' ; $E'' \stackrel{\text{def}}{=} E \setminus E'$; $n'' \stackrel{\text{def}}{=} n - n'$; $p_{E'} = p^{E''} : X_n \rightarrow X_{n''}$ the projection morphism of profile E' . Then we shall refer to the *kernel*

$$F \subseteq \pi_1^{\text{top}}(X_n)$$

of the induced *surjection* $\pi_1^{\text{top}}(X_n) \twoheadrightarrow \pi_1^{\text{top}}(X_{n''})$ [cf. Proposition 7.1, (i)] as the *fiber subgroup* of $\pi_1^{\text{top}}(X_n)$ of profile E' .

Remark 7.2.1. Note that by the *injectivity* of Proposition 7.1, (ii), it follows immediately that the *fiber subgroup of profile E'* of $\pi_1^{\text{top}}(X_n)$ [cf. the notation of Definition 7.2, (ii)] is equal to the *inverse image* via the natural injection $\pi_1^{\text{top}}(X_n) \hookrightarrow \pi_1(X_n)$ of Proposition 7.1, (ii), of the fiber subgroup of $\pi_1(X_n)$ of profile E' .

Corollary 7.3. (Discrete Extendability of Coverings) *Let X be a hyperbolic curve of genus ≥ 2 over \mathbb{C} ; $n \geq 1$ an integer; $X_n \subseteq P_n$ the n -th configuration space associated to X ; $(X_n^*)^{\log} \stackrel{\text{def}}{=} Z_n^{\log}$ the n -th log configuration space associated to X ;*

$$Y \rightarrow X_n$$

a finite étale morphism, where Y is connected; $Y^ \rightarrow X_n^*$ the normalization of X_n^* in Y ; G a strongly torsion-free profinite group;*

$$\phi : \pi_1^{\text{top}}(Y) \rightarrow G$$

a homomorphism [of abstract groups!]. Then:

(i) *Suppose that X is **proper**. Then the homomorphism ϕ **factors uniquely** through the natural morphism $\pi_1^{\text{top}}(Y) \rightarrow \pi_1^{\text{top}}(Y^*)$ induced by the open immersion $Y \hookrightarrow Y^*$.*

(ii) *Suppose that one of the following holds: (a) X is **proper**; (b) G is **prosolvable**. Also, let us suppose that the covering $Y \rightarrow X_n$ arises from a **product-theoretic** subgroup of finite index of $\pi_1^{\text{top}}(X_n)$. Then the kernel of ϕ is **product-theoretic**.*

Proof. Indeed, since the homomorphism $\phi : \pi_1^{\text{top}}(Y) \rightarrow G$ necessarily factors through the *profinite completion* of $\pi_1^{\text{top}}(Y)$, the conclusion of assertion (i) (respectively, assertion (ii) when (a) holds; assertion (ii) when (b) holds) follows immediately from Theorem 5.6, (i) (respectively, Theorem 5.6, (ii); Theorem 4.7). \circ

Corollary 7.4. (Group-theoreticity of Projections of Configuration Spaces III) For $\xi = \alpha, \beta$, let X^ξ be a **hyperbolic curve** over \mathbb{C} whose type is **neither (0, 3) nor (1, 1)**; $n_\xi \geq 1$ an integer; $X_{n_\xi}^\xi$ the n_ξ -th configuration space associated to X^ξ ; E_ξ the **index set** of $X_{n_\xi}^\xi$; $H_\xi \subseteq \Pi^\xi \stackrel{\text{def}}{=} \pi_1^{\text{top}}(X_{n_\xi}^\xi)$ a **product-theoretic subgroup of finite index**. Let

$$\gamma : H_\alpha \xrightarrow{\sim} H_\beta$$

be an **isomorphism of groups**. Moreover, if either $H_\alpha \neq \Pi^\alpha$ or $H_\beta \neq \Pi^\beta$, then we assume that X^ξ is of **genus** ≥ 2 , for $\xi = \alpha, \beta$. Then γ induces a **bijection** $\sigma : E_\alpha \xrightarrow{\sim} E_\beta$ [so $n_\alpha = n_\beta$] such that

$$\gamma(F_\alpha \cap H_\alpha) = F_\beta \cap H_\beta$$

for all **fiber subgroups** $F_\alpha \subseteq \Pi^\alpha$, $F_\beta \subseteq \Pi^\beta$, whose respective **profiles** $E'_\alpha \subseteq E_\alpha$, $E'_\beta \subseteq E_\beta$ correspond via σ .

Proof. Indeed, in light of Remark 7.2.1, Corollary 7.4 follows immediately by applying Corollaries 4.8, 6.3 to the isomorphism induced by γ between the *profinite completions* of H_α, H_β . \circ

Remark 7.4.1. There is a partial overlap between the content of Corollary 7.4 above and Theorems 1, 2 of [IIM].

Bibliography

- [Birm] J. Birman, *Braids, Links, and Mapping Class Groups*, Princeton University Press (1974).
- [DM] P. Deligne and D. Mumford, The Irreducibility of the Moduli Space of Curves of Given Genus, *IHES Publ. Math.* **36** (1969), pp. 75-109.
- [Fulton] W. Fulton, *Intersection Theory*, Springer-Verlag (1984).
- [EGA III] A. Grothendieck, *Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): III. Étude cohomologique des faisceaux cohérents, Première partie*, *Publ. Math. IHES* **11** (1961), pp. 5-167.

- [EGA IV] A. Grothendieck, *Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): IV. Étude locale des schémas et des morphismes des schémas, Seconde partie*, *Publ. Math. IHES* **24** (1965), pp. 5-231.
- [FJ] M. Fried and M. Jarden, *Field Arithmetic* (second edition), Springer-Verlag (2005).
- [Ill] L. Illusie, An Overview of the Work of K. Fujiwara, K. Kato and C. Nakayama on Logarithmic Etale Cohomology, in *Cohomologies p -adiques et applications arithmétiques (II)*, P. Berthelot, J.-M. Fontaine, L. Illusie, K. Kato, M. Rapoport, eds, *Asterisque* **279** (2002), pp. 271-322.
- [IIM] E. Irmak, N. Ivanov, J. D. McCarthy, *Automorphisms of surface braid groups*, preprint (arXiv:math.GT/0306069v1 3 Jun 2003).
- [Kato1] K. Kato, Logarithmic Structures of Fontaine-Illusie, *Proceedings of the First JAMI Conference*, Johns Hopkins Univ. Press (1990), pp. 191-224.
- [Kato2] K. Kato, Toric Singularities, *Amer. J. Math.* **116** (1994), pp. 1073-1099.
- [Knud] F. F. Knudsen, The Projectivity of the Moduli Space of Stable Curves, II, *Math. Scand.* **52** (1983), pp. 161-199.
- [Milne] J. S. Milne, *Étale Cohomology*, Princeton Mathematical Series **33**, Princeton University Press (1980).
- [Mzk1] S. Mochizuki, Extending Families of Curves over Log Regular Schemes, *J. reine angew. Math.* **511** (1999), pp. 43-71.
- [Mzk2] S. Mochizuki, Topics Surrounding the Anabelian Geometry of Hyperbolic Curves, *Galois Groups and Fundamental Groups, Mathematical Sciences Research Institute Publications* **41**, Cambridge University Press (2003), pp. 119-165.
- [Mzk3] S. Mochizuki, The Absolute Anabelian Geometry of Hyperbolic Curves, *Galois Theory and Modular Forms*, Kluwer Academic Publishers (2003), pp. 77-122.
- [Mzk4] S. Mochizuki, The Absolute Anabelian Geometry of Canonical Curves, *Kazuya Kato's fiftieth birthday, Doc. Math. 2003, Extra Vol.*, pp. 609-640.
- [Mzk5] S. Mochizuki, Galois Sections in Absolute Anabelian Geometry, *Nagoya Math. J.* **179** (2005), pp. 17-45.
- [Mzk6] S. Mochizuki, A combinatorial version of the Grothendieck conjecture, *Tohoku Math. J.* **59** (2007), pp. 455-479.
- [Mzk7] S. Mochizuki, Absolute anabelian cuspidalizations of proper hyperbolic curves, *J. Math. Kyoto Univ.* **47** (2007), pp. 451-539.
- [Naka] H. Nakamura, Galois rigidity of pure sphere braid groups and profinite calculus, *J. Math. Sci. Univ. Tokyo* **1** (1994), pp. 71-136.
- [NT] H. Nakamura and N. Takao, *Galois Rigidity of pro- l Pure Braid Groups of Algebraic Curves*, *Trans. Amer. Math. Soc.* **350** (1998), pp. 1079-1102.
- [RZ] L. Ribes and P. Zalesskii, *Profinite Groups, Ergebnisse der Mathematik und ihrer Grenzgebiete* **3**, Springer-Verlag (2000).

- [Stix] J. Stix, A monodromy criterion for extending curves, *Int. Math. Res. Not.* **29** (2005), pp. 1787-1802.
- [Vid] I. Vidal, Contributions à la cohomologie étale des schémas et des log schémas, *thèse* (2002).
- [Wehr] B. Wehrfritz, *Infinite Linear Groups*, *Ergebnisse der Mathematik und ihrer Grenzgebiete* **76**, Springer-Verlag (1973).